

Correção / Correction**FREQUENCY OF OBSESSIVE AND COMPULSIVE SYMPTOMS IN PATIENTS WITH BLEPHAROSPASM AND HEMIFACIAL SPASM**

R.P. Munhoz, H.A.G. Teive, M.V. DellaColletta, F.M.B. Germiniani, F.M. Iwamoto, C.H.F. Camargo, L.C. Wemeck

Arq Neuropsiquiatr 2005; 63:213-216

Tables of this article are missing*. They appear below.

Table 1. Yale-Brown scale scores in each group separated by symptom.

Symptoms	N°	Mean	Standard deviation	Minimum	Maximum	Median	p ⁽¹⁾
Obsessive							
BS	13	5,9	3,8	1,0	11,0	7,0	0,201
HFS	08	3,6	1,8	1,0	6,0	4,0	
Compulsive							
BS	03	12,7	5,5	9,0	19,0	10,0	-
HFS	04	4,8	4,1	1,0	10,0	4,0	
Both							
BS	04	14,0	14,2	4,0	35,0	8,5	0,938
HFS	09	9,4	3,5	4,0	15,0	9,0	
Total							
BS	20	8,6	7,6	1,0	35,0	7,5	0,417
HFS	21	6,3	4,0	1,0	15,0	6,0	

BS, Blepharospasm; HFS, Hemifacial spasm. (1) Mann-Whitney test.

Table 2. Patients with concomitant obsessive and compulsive symptoms with corresponding Yale-Brown scale scores.

Patient	MD	Obsessive symptoms	Compulsive symptoms	Total
1	BS	3	1	4
2	BS	3	4	7
3	BS	18	17	33
4	BS	3	7	10
5	HFS	7	5	12
6	HFS	3	1	4
7	HFS	6	3	9
8	HFS	9	3	12
9	HFS	10	4	14
10	HFS	11	2	13
11	HFS	7	4	11
12	HFS	2	4	6

MD, movement disorder; BS, blepharospasm; HFS, hemifacial spasm.

*The Editor presents to the Authors his excuses for such omission.