

Original Article

## Genetic parameters controlling the inheritance of glaucousness and yield traits in bread wheat

Parâmetros genéticos que controlam a herança de glaucosidade e características de produção em trigo mole

A. M. M. Al-Naggar<sup>a</sup> , K. F. Al-Azab<sup>b#</sup> , A. S. M. Younis<sup>c</sup>, I. O. Hassan<sup>b</sup> , M. A. E. Basyouny<sup>b</sup>  and M. Ayaad<sup>b\*</sup> 

# equal first author

<sup>a</sup>Agronomy Department, Faculty of Agriculture, Cairo University, Giza, Egypt

<sup>b</sup>Plant Research Department, Nuclear Research Center, Egyptian Atomic Energy Authority, Abo-Zaabal, Egypt

<sup>c</sup>Department of Field Crop Research, Agriculture and Biological Research Institute, National Research Center, Dokki, Cairo, Egypt

### Abstract

Wheat breeders frequently use generation mean analysis to obtain information on the type of gene action involved in inheriting a trait to choose the helpful breeding procedure for trait improvement. The present study was carried out to study the inter-allelic and intra-allelic gene action and inheritance of glaucousness, earliness and yield traits in a bread wheat cross between divergent parents in glaucousness and yield traits; namely Mut-2 ( $P_1$ ) and Sakha 93 ( $P_2$ ). The experimental material included six populations, i.e.  $P_1$ ,  $P_2$ ,  $F_1$ ,  $F_2$ ,  $BC_1$ , and  $BC_2$  for this wheat cross. A randomized complete block design with three replications was used, and a six parameters model was applied. Additive effects were generally more critical than dominance for all studied traits, except for plant height (PH) and grain yield/plant (GYPP). The duplicate epistasis was observed in spike length; SL, spikes/plant; SPP and days to heading; DTH. All six types of allelic and non-allelic interaction effects controlled SL, GYPP, DTH and glaucousness. All three types of epistasis, i.e. additive x additive, additive x dominance, and dominance x dominance, are essential in determining the inheritance of four traits (SL, GYPP, DTH and glaucousness). Dominance x dominance effects were higher in magnitude than additive x dominance and additive x additive in most traits. The average degree of dominance was minor than unity in six traits (glaucousness, grains/spike, spike weight, days to maturity, 100-grain weight and SL), indicating partial dominance and selection for these traits might be more effective in early generations. Meanwhile, the remaining traits (PH, SPP, GYPP and DTH) had a degree of dominance more than unity, indicating that overdominance gene effects control such traits and it is preferable to postpone selection to later generations. The highest values of narrow-sense heritability and genetic advance were recorded by glaucousness trait followed by SL and SPP, indicating that selection in segregating generations would be more effective than other traits.

**Keywords:** *Triticum aestivum*, generation mean analysis, gene effects, epistasis, heritability.

### Resumo

Os criadores de trigo frequentemente usam a análise da média de geração para obter informações sobre o tipo de ação do gene envolvida na herança de uma característica para escolher o procedimento de melhoramento útil para o aprimoramento da característica. O presente estudo foi conduzido para estudar a ação do gene interalélico e intraalélico e a herança de características de glaucosidade, precocidade e produção em um cruzamento de trigo mole entre pais divergentes em glaucosidade e características de produção; nomeadamente Mut-2 ( $P_1$ ) e Sakha 93 ( $P_2$ ). O material experimental incluiu seis populações, ou seja,  $P_1$ ,  $P_2$ ,  $F_1$ ,  $F_2$ ,  $BC_1$  e  $BC_2$  para este cruzamento de trigo. O delineamento experimental foi em blocos ao acaso com três repetições e aplicado um modelo de seis parâmetros. Os efeitos aditivos foram geralmente mais críticos do que a dominância para todas as características estudadas, exceto para altura da planta (AP) e rendimento de grãos / planta (GYPP). A epistasia duplicada foi observada no comprimento da ponta; SL, espigas/planta; SPP e dias para o cabeçalho; DTH. Todos os seis tipos de efeitos de interação alélica e não alélica controlaram SL, GYPP, DTH e glaucosidade. Todos os três tipos de epistasia, ou seja, aditivo x aditivo, aditivo x dominância e dominância x dominância, são essenciais na determinação da herança de quatro características (SL, GYPP, DTH e glaucosidade). Os efeitos de dominância x dominância foram maiores em magnitude do que aditivo x dominância e aditivo x aditivo na maioria das características. O grau médio de dominância foi menor do que a unidade em seis características (glaucosidade, grãos / espiga, peso da espiga, dias até a maturidade, peso de 100 grãos e SL), indicando dominância parcial, e a seleção para essas características pode ser mais eficaz nas gerações iniciais. Enquanto isso, os traços restantes (PH, SPP, GYPP e DTH) tiveram um grau de dominância maior do que a unidade, indicando que os efeitos do gene de superdominância controlam tais traços e é preferível adiar a seleção para gerações posteriores. Os maiores valores de herdabilidade no sentido restrito e avanço genético foram registrados pelo traço de glaucosidade seguido por SL e SPP, indicando que a seleção em gerações segregadas seria mais eficaz do que outros caracteres.

**Palavras-chave:** *Triticum aestivum*, análise da média de geração, efeitos gênicos, epistasia, herdabilidade.

\*e-mail: Mohammed\_ayaad@mail.hzau.edu.cn

Received: July 4, 2021 – Accepted: September 14, 2021



This is an Open Access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction

Wheat (*Triticum aestivum* L.) is the most important cereal crop in Egypt. In 2019, the wheat cultivated area in Egypt was 1.41 million hectares producing 9.0 million tons and Egypt imported 10.42 million tons of wheat in 2019 to satisfy the local consumption, costing the government about 3024 million US dollars (FAO, 2021). Therefore, increasing wheat production horizontally and vertically is the key to saving hard currency for wheat imports. Thus, all efforts of plant breeders are required to improve the productivity of wheat varieties. This development could be achieved by adding genes of traits that increase the grain yield of current varieties.

Glaucousness, the waxy bloom on the surface of leaves, leaf sheaths and spikes, has been associated with several traits and physiological processes, mainly related to an increased drought and heat tolerance and thus higher yield under dry conditions (Johnson et al., 1983; Richards et al., 1986; Febrero et al., 1998; Merah et al., 2000; Samuels et al., 2008; Bi et al., 2017). Early genetic studies in wheat revealed two loci for wax production, termed *W1* and *W2*, and two inhibitors of wax loci, *Iw1* and *Iw2*, that inhibit glaucousness (Tsunewaki and Ebana 1999). Further analyses revealed the genetic control of glaucousness to be more complex, as several quantitative trait loci (QTL) were identified (Börner et al., 2002; Kulwal et al., 2003; Mason et al., 2010; Bennett et al., 2012). Inheritance of glaucousness has been previously reported in bread wheat (Jensen and Driscoll, 1962; Stuckey, 1972; Liu et al., 2007). In addition, the evidence of additive gene action influencing glaucousness was also reported in bread wheat (Al-Bakry, 2010). The genetics of glaucousness and precise interaction between the genes is still not fully understood.

Yield has prime importance in any breeding program, but it is a much complex trait. Habib and Khan (2003), Mohammad et al. (2003) and Riaz and Chowdhry (2003) described the additive type of gene action with partial dominance controlling this trait. On the other hand, Inamullah and Hassan (2005), Dere and Yildirim (2006), and Hassan et al. (2008) showed that overdominance type of gene action control this parameter. Considerable studies in this concern were conducted by many investigators, e.g. Kearsy and Pooni (1996), Farshadfar et al. (2001), Novoselovic et al. (2004), Erkul et al. (2010), Farshadfar et al. (2013), Ljubicic et al. (2016a, b). Epistasis was reported in many studies of wheat, Przulj and Mladenov (1999) and Al-Naggar et al. (2010a, b, 2017) and (Al-Azab et al., 2017) for grain yield traits.

The choice of selection and breeding procedures for genetic improvement of any crop is mainly dependent on the knowledge of the type and relative amount of genetic components and the presence of allelic and non-allelic interaction for different traits in the plant materials under investigation. Information on the type of gene action involved in inheriting a trait helps decide the breeding procedures for plant improvement. The breeding strategy mainly depends on the different gene effects, i.e. dominance, additive, and non-allelic interaction. Plant breeders frequently use generation mean analysis to obtain information of allelic and non-allelic gene interaction

controlling the economic traits in wheat (Khattab et al., 2001; Esmail and Khattab, 2002; Akhtar and Chowdhry, 2006; Khaled, 2007 and Farag, 2009). Generation means analysis is a tool for designing the most appropriate breeding approaches to develop crop varieties with desired traits and commonly used in studying the inheritance of quantitative traits. The generation means analysis procedure is based on the hypothesis that the studied generations must arise from a cross involving two contrasting genotypes. This model is free from the limitations of other models and can estimate the genetic markers needed for each trait (Mather and Jinks, 1971). In this way, in addition to estimate additive and dominance gene effects, the effects of epistasis can also be estimated using the scaling test.

The present study was carried out to obtain information about gene action, inheritance nature, heterosis, inbreeding depression, the average degree of dominance, broad and narrow sense heritability and expected genetic advance from selection for induced glaucousness, agronomic, and yield traits in a bread wheat cross between a glaucous parent (Mut-2) and non-glaucous parent (Sakha 93). Such information would help plant breeders to implement an exact breeding procedure for developing high-yielding varieties.

## 2. Materials and Methods

### 2.1. Plant materials

Two bread wheat genotypes, i.e. glaucous wheat mutant (Mut-2) and Sakha 93 cultivar (non-glaucous), were crossed and used as parents in the present study because of their diversity for glaucousness grain yield and yield-related traits. Sakha 93 bread wheat was developed by the wheat research department, field crops research institute, Agriculture Research Center, Egypt, while Mut-2 was developed by the wheat breeding program of Atomic Energy Authority, Anshas, Egypt (Al-Naggar et al., 2013). It was selected in the  $M_2$  population resulting from Sakha 93 with 350 Gy of gamma rays. Mut-2 has high epicuticular wax content on leaf sheath, leaf blade, peduncle and spike. It has too many kernels/spike with long spike, but its tillers number is moderate. However, leaf-blades and spikes of Sakha 93 are completely non-glaucous. Also, it has a moderate number of spikelets/spike and a moderate number of kernels/spike, but it has a high number of tillers/plant.

In the first winter season (2016/2017), the mutant Mut-2 (P1) was crossed to Sakha 93 (P2), and an  $F_1$  seed was produced. Thus,  $BC_1$  ( $F_1 \times P_1$ ) and  $BC_2$  ( $F_1 \times P_2$ ) were developed in 2017/2018 season, while the remaining  $F_1$  plants were self-pollinated to produce  $F_2$  seeds. The two parents and their  $F_1$ ,  $F_2$ ,  $BC_1$ , and  $BC_2$  populations were grown during the 2018/2019 season for field evaluation.

### 2.2. Experimental design

The experiment was conducted in a randomized complete block design (RCBD) with three replications at the experimental farm of Plant Research Department,

Nuclear Research Center, Atomic Energy Authority, Egypt. Each replication composed 20 experimental rows; 2 rows for each of  $P_1$  and  $P_2$ , four rows for  $F_1$ , six rows for  $F_2$ , and three rows for  $BC_1$  and  $BC_2$  populations. Individual grains were planted in 3-meter rows. Each row included 30 plants spaced 10 cm apart. Rows were spaced 30 cm apart. The sowing date was 20<sup>th</sup> November (recommended sowing date for north Egypt). All other Agricultural practices were applied as recommended for wheat planting.

### 2.3. Data recorded in the field

Data for each replication were noted on ten guarded plants for each  $P_1$ ,  $P_2$  and  $F_1$ , 75 plants of  $F_2$ , and 20 plants of  $BC_1$  and  $BC_2$ . The following traits were recorded: 1. Days number from sowing date to 50% heading (DTH), 2. Days from sowing date to 50% physiological maturity (DTM), 3. Plant height (PH), 4. Main spike length (SL), 5. Number of fertile spikes per plant (SPP), 6. Grains number per spike (GPS), 7. Main spike weight (SW), 8. 100-grain weight (100-GW) and 9. Grain yield/plant (GYPP).

### 2.4. Epicuticular wax quantification

Thirty flag leaf blades of each parent,  $F_1$ ,  $F_2$ -glaucous,  $F_2$ -moderately glaucous,  $F_2$ -nonglaucous,  $BC_1$ -glaucous,  $BC_1$ -moderately glaucous,  $BC_2$ -moderately glaucous and  $BC_2$ -nonglaucous plants were immersed individually, each for 15 sec in 15 ml redistilled chloroform. The extracts were filtered and evaporated at 35 °C. After drying for 24 hours at room temperature, the residues were weighed (Silva Fernandes et al., 1964; Ebercon et al., 1977). The amount of wax was considered against leaf area (both leaf surfaces) as mg/dm<sup>2</sup> of each sample.

### 2.5. Statistical and genetic analysis

The analysis of variance of RCBD for each studied character was performed using SAS v9.4 (SAS Institute, 2011) software package to test the differences among the six studied populations. Scaling tests outlined by Mather (1949) and Hayman and Mather (1955) were achieved to detect the presence of epistasis (non-allelic interaction).

The significance of scaling tests was executed using a t-test by dividing the effects of A, B, C and D on their respective standard error. The significance of any scaling test implies the inadequacy of the simple additive-dominance model to explain genetic variances and thus indicates the presence of epistasis. Then the six-parameter model was performed as outlined by Hayman (1958) and Mather and Jinks (1982). The gene effects were estimated for each trait and defined as Hayman's notations; mean effects ( $m$ ) = mean of the  $F_2$  generation, the allelic gene effects ( $d$ ) = additive gene effects, ( $h$ ) = dominance gene effects, and the non-allelic gene effects ( $i$ ) = additive × additive gene effects, ( $j$ ) = additive × dominance gene effects, and ( $l$ ) = dominance × dominance gene effects.

The genetic components of variance were calculated according to Mather and Jinks (1982). Broad sense ( $h^2_{bs}$ ) and narrow sense ( $h^2_{ns}$ ) heritability were estimated according to Warner (1952). Expected genetic advance (GA%) from selection as a percent of the mean of  $F_2$  was computed according to Johnson et al. (1955) using selection intensity of 5% ( $k=2.06$ ). Heterosis was calculated as the percentage deviation of  $F_1$  generation from the mid parent (heterosis) and better parent (heterobeltiosis) according to Fonseca and Patterson (1968). Inbreeding depression was estimated as the percentage decrease of the  $F_2$  from the  $F_1$ . The average degree of dominance ( $(H/D)^{1/2}$ ) was tested by the formula suggested by Jinks and Hayman (1953).

## 3. Results and Discussion

### 3.1. Mean performance

Analysis of variance (Table 1) showed significant ( $P \leq 0.01$ ) mean squares due to genotypes (populations) for all studied traits of the studied cross, except for 100-grain weight, indicating the existence of genetic variation for these traits of the studied materials. The significance among the six populations made it possible to go further for generation mean analysis. The same results were obtained by Ataei et al. (2017) and Feltaous et al. (2020).

**Table 1.** Mean squares from the analysis of variance of parents, their  $F_1$ ,  $F_2$ , and backcrosses for studied in a cross between Sakha 93 and Mut-2 mutant lines .

S. O. V.	df	Mean Squares				
		Plant height	Spike length	Spikes/plant	100-grain weight	Grain yield/plant
Replications	2	13.72	0.06	2.67	0.48	4.29
Genotypes	5	164.98**	39.68**	18.36**	0.57	369.7**
Error	10	5.92	1.32	1.33	0.27	5.81
		50% Heading	50% Maturity	Spike weight	Grains /spike	Wax content
Replications	2	9.06	7.39	0.01	2.39	0.02
Generations	5	144.2**	127.5**	0.33**	214.8**	0.75**
Error	10	1.66	2.79	0.01	4.86	0.001

S.O.V.: Source of variation, df: Degrees of freedom, \*\* indicates significance at 0.01 probability level.,

Mean performances of  $P_1$ ,  $P_2$ ,  $F_1$ ,  $F_2$ ,  $BC_1$ , and  $BC_2$  for the ten studied traits are presented in Table 2. The difference between  $P_1$  and  $P_2$  was significant ( $P \leq 0.05$  or  $0.01$ ) for all studied traits, except 100-grain weight; such difference was more pronounced for wax content, GYPP, SL, GPS, DTH and DTM. The divergence between parents of the hybrid Mut-2  $\times$  Sakha 9 is a prerequisite for studying gene action and inheritance of glaucousness, agronomic, and yield traits using generation mean analysis. A similar conclusion was reported by Al-Naggar et al. (2017), Ataei et al. (2017), Koubisy (2019), Salmi et al. (2019) and Feltaous et al. (2020).

The  $F_1$  mean values surpassed the mid-parent values for 6 out of 10 studied traits, namely PH, SPP, GYPP, 100-grain weight, GPS and wax content, indicating the prevalence of heterotic and dominance effects in controlling these traits. The  $F_1$  mean values significantly ( $P \leq 0.01$ ) surpassed the higher parent for three traits, namely GYPP, SPP and PH, indicating that overdominance controls the inheritance of these traits. The means of the  $F_2$  population tended to be less than  $F_1$  means for all studied traits, except days to heading and days to maturity, indicating the presence

of non-additive components of genetic variance for these traits. However, means of  $BC_1$  and  $BC_2$  populations tended toward the mean of its recurrent parent. These results agreed with those reported by Tammam (2005), Kumar et al. (2017), Al-Azab et al. (2017), Abd El-Hady et al. (2018), Koubisy (2019) and Feltaous et al. (2020).

### 3.2. Adequacy of additive-dominance model

The scaling test, i.e. A, B, C, and D, were performed to determine the efficacy of the additive-dominance model for the inheritance of the studied traits. The significance of one or more of these scales indicates that the additive-dominance model is inadequate for describing the inheritance of the studied trait and the presence of epistasis (non-allelic interaction).

Estimates of A, B, C and D scaling tests and their significance are presented in Table 3. The significance of the C scale reveals dominance  $\times$  dominance type of epistasis. The significant D scale reveals additive  $\times$  additive type of gene interaction, and the significance of both C and D scales indicates additive  $\times$  additive and dominance  $\times$  dominance

**Table 2.** Mean performance of studied traits for parents, their  $F_1$ ,  $F_2$ , and backcrosses in the cross between Mut-2 ( $P_1$ ) and Sakha 93 ( $P_2$ ).

Trait	Genotypes						LSD	
	$P_1$	$P_2$	$F_1$	$F_2$	$BC_1$	$BC_2$	0.05	0.01
Plant height (cm)	91.00	86.67	105.67	99.00	97.00	93.33	4.43	6.30
Spike length(cm)	23.00	11.67	18.00	17.67	18.00	19.00	2.09	2.97
Spikes/plant	8.00	10.67	15.00	13.33	13.33	12.67	2.10	2.98
100-Grain weight (g)	4.90	5.50	6.20	5.20	5.53	5.60	0.95	1.34
Grain yield /plant (g)	43.33	67.17	76.37	68.73	64.70	66.03	4.39	6.24
50% Heading (day)	74.67	96.00	86.00	88.00	84.00	88.00	2.34	3.33
50% Maturity (day)	119.00	139.33	129.00	130.33	128.33	131.33	3.04	4.32
Spike weight (g)	4.20	5.23	4.80	4.67	4.67	4.77	0.18	0.26
Grains / spike	90.00	66.67	89.33	83.67	84.33	82.33	4.01	5.70
Wax content (mg/dm <sup>2</sup> )	2.95	1.55	2.54	2.31	2.56	1.94	0.04	0.06

**Table 3.** Estimates and significances of scaling test parameters A, B and C for all studied traits in a cross between glaucous mutant line (Mut-2) and non-glaucous wheat cultivar (Sakha 93).

Trait	A	B	C	D
Plant height	-8.88**	-7.35**	-1.21	7.67**
Spike length	-2.69**	7.11**	0.56	-1.66
Spikes/plant	2.44*	-0.21	4.84**	0.66
100 grain weight	0.27	0.78	-0.31	-0.73
Grain yield /plant	-13.57**	3.95**	2.13	6.73**
50% Heading	-7.70**	6.34**	1.00	4.00**
50% Maturity	-5.29**	9.59**	1.63	1.00
Spike weight	-0.66	0.32	-0.63	-0.10
Grains/spike	-10.08**	12.61**	3.01*	0.68
Wax content	-0.39	-0.25	-0.3	0.12**

\*, \*\* Significant at  $P \leq 0.05$  and  $P \leq 0.01$ , respectively.

types of gene interaction. The adequacy of scales is vital because, in most cases, the estimation of additive and dominance components of the variance is made assuming the absence of genes interaction.

The values of A, B, C and D scales should significantly differ from zero within the limits of their standard error. However, the results in Table 3 indicate that the scaling test values significantly differed from zero for most studied traits, indicating the presence of non-allelic gene interaction and dominance-additive model are inadequate for explaining the inheritance of these traits. The exceptions are 1000-grain weight and spike weight, which did not show significance in any one of the scaling test parameters, indicating that epistasis was not involved in the genetic control of 1000-grain weight and spike weight and the adequacy of simple additive-dominance model for estimating components of genetic variance of such traits (*i.e.* three-parameters model). Al-Bakry (2010) and Al-Azab et al. (2017) reported that no epistasis was detected for 1000-kernel weight in wheat, indicating that estimates of gene effects for this trait were free from linkage bias. These results are harmony those reported by Tammam (2005), Moussa (2010), El-Aref et al. (2011), Zaazaa et al. (2012), Amin (2013), Abd El-Rahman (2013), Hamam and Negim (2014), El-Hawary and El-Shafey (2016), Kumar et al. (2017) and Abd El-Hady et al. (2018).

### 3.3. Gene effects

The six-parameter model was used in the present study to determine the nature and magnitude of gene effects involved in the genetic control of eight studied traits; PH, SL, SPP, GYPP, DTH, DTM, GPS and wax content (Table 4).

The estimated mean effect parameter ( $m$ ), which reflects the contribution due to the overall mean plus the locus effects and interactions of the fixed loci, was significant ( $P \leq 0.01$ ) for all traits of the studied cross, indicating that these characters are quantitatively inherited and show the importance of non-allelic interactions. These results are in harmony with earlier results reported by Bilgin et al. (2016), El-Hawary and El-Shafey (2016), Al-Azab et al. (2017), Patel et al. (2018), Abd El-Hady et al. (2018), Salmi et al. (2019), Koubisy (2019) and Feltaous et al. (2020).

The additive (a) gene effects were positive significance ( $P \leq 0.05$  or  $0.01$ ) for PH, SPP, and wax content and negative significance ( $P \leq 0.01$ ) for SL, GYPP, DTH, DTM and GPS, indicating possible improvement of these traits through selection in early segregating generations. Similar results for yield and its components were reported by Ataei et al. (2017), Al-Azab et al. (2017), Patel et al. (2018), Mahpara et al. (2018), Soliman (2018), Koubisy (2019) and Feltaous et al. (2020). In addition, the evidence of additive gene action influencing glaucousness was also reported in bread wheat (Al-Bakry, 2010).

The dominance effects (d) were positive significance ( $P \leq 0.01$ ) for SL, SPP, GYPP, GPS and wax content; moreover, negative significance ( $P \leq 0.01$ ) for DTH. Therefore, to make the selection more efficient in these studied traits, it is preferable to postpone the selection to late generations to increase homozygosity.

The results indicated that both additive and dominance have important contributions in the inheritance of the studied traits, suggesting that selection for desirable traits might be effective in early generations. However, it is better to postpone it to later segregating generations. These results are in agreement with those reported by Ljubicic et al. (2016b), Al-Azab et al. (2017), Koubisy (2019) and Feltaous et al. (2020).

Additive x additive interaction (aa) was found significant ( $P \leq 0.01$ ) with a positive value (Table 4) for SL. A negative and significant (aa) interaction was found in PH, GYPP, DTH and wax content. These findings agree with previous results reported by Koubisy (2019) and Feltaous et al. (2020).

Additive x dominance interaction (ad) presented positive significance ( $P \leq 0.01$ ) (Table 4) in SL, GYPP, DTH, DTM and GPS, but found significant ( $P \leq 0.01$ ) and negative in wax content. Significance of additive x dominance interaction suggests that it is preferred to delay selection for improving such traits to later segregating generations. These results are in harmony with Patel et al. (2018), Soliman (2018), Koubisy (2019) and Feltaous et al. (2020).

Dominance x dominance interaction (dd) effects were found positive and significant ( $P \leq 0.01$ ) for PH, grain GYPP, DTH, GPS and wax content, and negative significance ( $P \leq 0.01$ ) for SL and SPP (Table 4). Dominance x dominance gene interaction was higher in magnitude

**Table 4.** Gene effects from generation mean analysis for eight traits in the wheat cross between glaucous mutant line (Mut-2) and non-glaucous wheat cultivar (sakha 93).

Trait	Mean	additive	dominance	aa	ad	dd	Epistasis
Plant height	99**	4.33*	1.495	-15.34**	1.505	23.69**	Complementary
Spike length	17.67**	-11.33**	3.985**	3.32**	4.665**	-6.65**	Duplicate
Spikes/plant	13.33**	2.67*	4.345**	-1.32	-0.675	-2.01*	Duplicate
Grain yield /plant	68.73**	-23.84**	7.66**	-13.46**	10.59**	15.24**	Complementary
50% Heading	88**	-21.33**	-7.335**	-8**	6.665**	6.67**	Duplicate
50% Maturity	130.33**	-20.33**	-2.165	-2.0	7.165**	-0.99	Complementary
Grains/spike	83.67**	-23.33**	9.635**	-1.36	9.665**	3.37*	Complementary
Wax content	2.31**	1.4**	0.05*	-0.24**	-0.08**	0.82**	Complementary

(aa) additive\*additive, (ad) additive\*dominance, (dd) dominance\*dominance\*, \*\* Significant at  $P \leq 0.05$  and  $P \leq 0.01$ , respectively.

than additive × dominance and additive × additive in most traits, indicating that these traits are greatly affected by dominance and non-allelic interactions. Therefore, it is advisable to delay selection to late segregating generations with increased homozygosity.

The present results reveal that all non-allelic interactions (epistasis), *i.e.* additive × additive, additive × dominance, and dominance × dominance, are essential in determining the inheritance of SL, GYPP, DTH and wax content. Epistasis was involved in the genetic control of kernels/spikelet, spikes/plant and GYPP in wheat by Al-Bakry et al. (2008) and Al-Bakry and Al-Naggar (2011). Similar results were reported by Ataei et al. (2017), Al-Azab et al. (2017) and Feltaous et al. (2020), who stated that all non-allelic interactions are essential factors in controlling the expression of GYPP.

The type of epistasis can be determined only when dominance (d) and dominance × dominance (dd) gene effects were significant. Consequently, the type of epistasis can be determined, as Kearsey and Pooni (1996) concluded. When these effects have the same sign, epistasis is of complementary type. The duplicate epistasis determined when the sign was different. The results in Table 4 showed that five traits were controlled by complementary epistasis (plant height, grain yield/plant, days to maturity, grains/spike and wax content). The duplicate epistasis was observed in SL, SPP and DTH. The presence of duplicate dominant epistasis in the expression of a trait would limit the range of variability (Kearsey and Pooni 1996). This type of epistasis limits the efficacy of selection in early segregating generations; therefore, delaying the selection to advanced generations will be an excellent decision to exploit the transgressive segregations. Various investigators reported similar results (Ljubicic et al., 2016b; Abdallah et al., 2019; Raikwar, 2019; Salmi et al., 2019).

Results showed that all six types of allelic (additive and dominance) and non-allelic (epistasis) interaction gene effects controlled four out of ten studied traits; SL, GYPP, DTH, and wax content.

For the traits grain yield/plant, spike length, grains/spike plant height and wax content, results showed a predominance of additive (a) followed by dominance × dominance (dd), additive × additive (aa), additive × dominance (ad) and dominance (d) gene effects in descending order, suggesting that selection for improving these traits could be started in early segregating generations ( $F_2$ ). However, it is preferable to postpone selection to later segregating generation ( $F_4$  or  $F_5$ ) to utilize the non-additive gene effects after homozygosity and release the homozygotes. This conclusion is in complete agreement with that reported by Al-Naggar et al. (2010a, b, 2015, 2017).

For SPP, it is observed from Table 4 that the dominance gene effects are the most predominated type of gene action followed by dominance × dominance gene effects, indicating that it is preferable to postpone selection to later segregating generations for this trait.

Regarding plant height, dominance × dominance was the most predominant gene effect, followed by

additive × additive, additive and dominance in descending order (Table 4), indicating that selection for this trait should be postponed to later segregating generations. This result is in agreement with that reported by Nanda et al. (1981).

For the earliness traits; DTH and DTM, it is observed from Table 4 that additive is the major type of gene action followed by dominance, dominance × dominance and additive × additive was controlling inheritance of DTH and by additive × dominance followed by dominance gene effects controlling inheritance of DTM. In general, for earliness traits, selection could be practised in the  $F_2$  generation, but it is also preferred to be postponed to later segregating generations to get more efficient selection gain.

Our results are in agreement with those reported by Tammam (2005), Hussein et al. (2009) and Al-Naggar et al. (2010a, b, 2015, 2017) for earliness and Przulj and Mladenov (1999), Al-Bakry et al. (2008) and Al-Bakry and Al-Naggar (2011) for grain yield traits. On the contrary, our results disagree with those reported by other researchers (Menshaw, 2005, 2007) for days to heading and to maturity and Darwish and El-Hosary (2003), Riaz and Chowdhry (2003) and Al-Naggar et al. (2012) for grain yield traits. The divergence in results can be attributed to the differences in the genetic materials used in different research studies.

### 3.4. Components of genetic variance

The components of genetic variance, additive and dominance variances in Table 5, revealed that the dominance variance component was higher than the additive one for PH and GYPP, indicating that the dominance gene effect plays a vital role in the inheritance of these characters and selection may be effective in later segregating generations. On the other hand, the additive variance was more important in the genetic system controlling the remaining traits, suggesting the effectiveness of selection in early segregating generations. Similar results were obtained by El-Aref et al. (2011), Amin (2013), Hamam and Negim (2014), Abd El-Hady et al. (2018) and Koubisy (2019).

The average degree of dominance  $(H/D)^{1/2}$  given in Table 5 was less than unity in six traits: wax content, grains/spike, spike weight, days to maturity, 100-grain weight, and spike length. These results confirm the role of partial dominance gene effects in controlling these traits and selection for these traits might be more effective in later segregating generations. These findings reveal that additive effects  $[(H/D)^{1/2} < 1]$  were more critical in the majority of the traits than dominance effects  $[(H/D)^{1/2} \geq 1]$  in early generations. Meanwhile, the remaining four traits (PH, SPP, GYPP and DTH) had a degree of dominance more than unity, indicating that overdominance gene effects control such traits and selection should be delayed to later generations to improve these traits. Similar results were reported by El-Aref et al. (2011), Amin (2013), Hamam and Negim (2014), Ataei et al. (2017), Soliman (2018), Abd El-Hady et al. (2018), Koubisy (2019), Koubisy (2019) and Feltaous et al. (2020).

**Table 5.** Additive, dominance, genotypic, environmental and phenotypic variances, heritability in broad ( $h^2_b$ %) and narrow ( $h^2_n$ %) sense, genetic advance (GA %), degree of dominance ("a"), heterosis%, heterobeltiosis% and inbreeding depression (ID%).

Parameter	Plant height	Spike length	Spikes/plant	100-Grain weight	Grain yield/plant	Days to heading	Days to mature	Spike weight	Grains/spike	Wax content
<b>Additive</b>	6.34	14.39	6.75	0.59	1.28	22.94	17.45	0.23	6.7	0.75
<b>Dominance</b>	18.67	6.65	5.01	0.27	8.61	13.13	6.42	0.13	2.17	0.23
<b>Genotypic</b>	25.01	21.04	11.75	0.87	9.89	36.07	23.87	0.36	8.87	0.98
<b>Environmental</b>	1.85	0.46	0.45	0.01	6.72	0.59	40.06	0.01	1.57	0.01
<b>Phenotypic</b>	26.86	21.5	12.2	0.88	16.61	36.66	63.93	0.37	10.44	0.99
$h^2_b$ %	93.08	97.88	96.31	99.1	59.55	98.38	37.34	97.97	84.95	98.89
$h^2_n$ %	23.59	66.94	55.28	68.0	7.71	62.57	27.3	62.77	64.18	75.76
<b>GA %</b>	2.54	36.19	29.84	25.27	0.94	8.87	3.45	16.84	5.11	67.22
<b>"a"</b>	2.43	0.96	1.22	0.96	3.67	1.07	0.86	0.84	0.80	0.78
<b>Heterosis%</b>	18.95**	3.84	60.69**	19.23**	38.23**	0.78	-0.13	1.80*	14.04**	12.89**
<b>Heterobeltiosis%</b>	21.92	-21.74	40.58**	12.73**	13.70**	15.17**	8.40**	-8.22**	-0.74	-13.89**
<b>ID%</b>	6.31**	1.83**	11.13**	16.13**	10.00**	-2.33	-1.03	2.71*	6.34**	9.06**

\*\* significant at 0.05 and 0.01 probability levels, respectively.

### 3.5. Heritability and genetic advance

Estimating heritability, either in broad-sense or narrow-sense, enables plant breeders to choose the right breeding strategy. The broad-sense heritability determines the heritable portion of the total variation, while the narrow-sense heritability determines the portion of an additive effect. Heritability was categorized into three groups; high ( $\geq 60\%$ ), moderate (30-60%), and low heritability (0-30%). The present results (Table 5) indicate that the broad sense heritability falls between moderate for DTM and GYPP to high for the rest of the studied traits. The broad-sense heritability ranged from 37.34% for DTM up to 99.1% for 100-grain weight. The present results revealed that a high portion of the phenotypic variation could be attributed to the genetic variation for most studied traits.

On the other side, the narrow-sense heritability ranged from 7.71% for grain yield/plant up to 75.76% for wax content. The values of narrow-sense heritability imply that the additive effects have a higher role than the dominance effect in controlling the above mentioned studied traits. The present results indicate that the selection process will be effective, which will result in a higher response. These results are in agreement with those reported previously by Farshadfar et al. (2013), Abd El-Hamid and Ghareeb (2018) and Koubisy (2019).

Genetic advance (GA %) as a percent of the mean was classified into three groups, i.e. low (less than 10%), moderate (between 10% and 20%), and high (more than 20%). The results in Table 5 showed that genetic advance values were high for SL, SPP, 100-grain weight, and wax content, moderate for spike weight and low for PH, GYPP, DTH, DTM and GPS.

The highest heritability values in narrow-sense and genetic advance were recorded by glaucousness trait followed by SL and SPP (only in GA%), indicating that selection would be more effective in segregating generations than other traits.

The high genetic advance values indicate that the direct selection is effective, while the low values reveal that indirect selection can be applied for these traits with correlated high heritability traits. These results are in agreement with those reported previously by Farshadfar et al. (2013), Said (2014), Ninghot et al. (2016) and Koubisy (2019).

### 3.6. Heterosis and inbreeding depression

Percentages of heterosis over mid-parent and better parent (heterobeltiosis) and inbreeding depression are presented in Table 5. Positive and significant ( $P \leq 0.05$  or  $0.01$ ) heterosis and heterobeltiosis values were observed for all studied traits, except DTH and DTM, and heterobeltiosis for GPS. These results are in harmony with Abd El-Rahman (2013), Hamam and Negim (2014), Abd El-Hamid and El-Hawary (2015), and Abd El-Hady et al. (2018). The positive significance of heterobeltiosis ( $P \leq 0.01$ ) for GYPP and SPP, indicates that the studied cross could be promising in wheat breeding programs to produce elite recombinant inbred lines in its early segregating generations.

Inbreeding depression measured as a reduction in performance of  $F_2$  generation relative to  $F_1$  is presented in Table 5. Results showed significant ( $P \leq 0.05$  or  $0.01$ ) and positive inbreeding depression values for all studied characters, except DTH and DTM. These results are expected because the expression of heterosis in  $F_1$  will be reduced in the  $F_2$  generation due to selfing and starting homozygosity. These results are in close agreement with Yadav and Singh (2011); Hamam and Negim (2014); Said, 2014; El-Hawary, 2016; Jaiswal et al., 2018. These results exhibit the importance of inbreeding depression and heterosis jointly in the positive selection process. These results are in good agreement with those reported by Kumar et al. (2017), Jaiswal et al. (2018) and Kumar et al. (2018).

#### 4. Conclusion

This investigation concluded that epistasis was involved in the genetic control of all studied traits, except 100-grain weight and spike weight. Dominance and dominance × dominance type of epistasis was the most dominated types of gene effects controlling SL, SPP, GYPP, DTH, GPS and wax content traits in the genetic material under study, indicating that selection for improving these traits in the future should be practised in the later segregating generations (F<sub>4</sub> or F<sub>5</sub>) in order to achieve high genetic gain. The highest values of heritability in narrow-sense and genetic advance were recorded by glaucousness trait followed by spike length and spikes/plant, indicating that selection would be more effective in segregating generations than other traits.

#### Acknowledgements

This work was partially supported by Nuclear Research Center, Egyptian Atomic Energy Authority. The authors thank Prof. Sobieh for his excellent fieldwork managing and his guidance, Prof. Hesham for his partial supported via Radiation Detection project.

#### References

- ABD EL-HADY, M.A., FERGANI, M.A., EL-TEMSAH, M.E. and ABDELKADER, M.A., 2018. Physiological response of some wheat cultivars to antitranspirant under different irrigation treatments. *Middle East Journal of Agriculture Research*, vol. 7, no. 1, pp. 100-109.
- ABD EL-HAMID, E.A.M. and EL-HAWARY, M.N.A., 2015. Generation mean analysis for some quantitative traits in bread wheat: I-Agronomic characteristics. *Egyptian Journal of Agricultural Research*, vol. 93, no. 2A, pp. 2015.
- ABD EL-HAMID, E.A.M. and GHAREEB, Z.E., 2018. Generation mean analysis for estimating some genetic parameters in four bread wheat crosses. In: *Proceedings of the 7th Field Crops Conference*, 18-19 December 2018, Giza, Egypt. Giza, pp. 17-29.
- ABD EL-RAHMAN, M.E., 2013. Estimation of some genetic parameters through generation mean analysis in three bread wheat crosses. *Alexandria Journal of Agricultural Research*, vol. 58, no. 3, pp. 183-195.
- ABDALLAH, E., SALEM, A.H., ALI, M.M.A. and KAMAL, K.Y., 2019. Genetic analysis for earliness and grain yield of bread wheat (*Triticum aestivum* L.). *Zagazig Journal of Agricultural Research*, vol. 46, no. 6, pp. 1769-1784. <http://dx.doi.org/10.21608/zjar.2019.51867>.
- AKHTAR, N. and CHOWDHRY, M.A., 2006. Genetic analysis of yield and some other quantitative characters in bread wheat. *International Journal of Agriculture and Biology*, vol. 8, pp. 523-527.
- AL NAGGAR, A.M.M., SHABANA, R., ABD-EL-ALEEM, M.M. and EL-RASHIDY, Z., 2017. Mode of inheritance of low-N tolerance adaptive traits in wheat (*Triticum aestivum* L.) under contrasting nitrogen environments. *Spanish Journal of Agricultural Research*, vol. 15, no. 2, e0702. <http://dx.doi.org/10.5424/sjar/201715210808>.
- AL-AZAB, K.F., ABDEL-LATTIF, H.M. and AL-NAGGAR, A.M.M., 2017. Allelic and non-allelic gene interactions controlling the inheritance of earliness and yield traits in three crosses of bread wheat. *Egyptian Journal of Plant Breeding*, vol. 21, no. 7, pp. 1309-1325.
- AL-BAKRY, M.R.I. and AL-NAGGAR, A.M.M., 2011. Genetic recombinants and transgressive segregants selected for grain yield and its related traits in bread wheat. *Egyptian Journal of Plant Breeding*, vol. 15, no. 4, pp. 145-154.
- AL-BAKRY, M.R.I., 2010. Inheritance of induced glaucousness, grain yield, and yield-related traits in bread wheat, *Triticum aestivum* L. *Egyptian Journal of Genetics and Cytology*, vol. 39, no. 1, pp. 15-27.
- AL-BAKRY, M.R.I., AL-NAGGAR, A.M.M. and MOUSTAFA, H.A.M., 2008. Improvement of grain yield of glaucous wheat mutant line via backcrossing. *Egyptian Journal of Plant Breeding*, vol. 12, no. 2, pp. 123-131.
- AL-NAGGAR, A., HUSSEIN, H.A.A., HUSSEIN, E.H.A., SABRY, S.R.S. and GAD, K.I.M., 2010a. Genetic diversity and bulked segregant analysis for earliness in bread wheat. *Arab Journal of Biotechnology*, vol. 13, pp. 157-172.
- AL-NAGGAR, A.M.M., SABRY, S.R.S. and GAD, K.I.M., 2010b. Epistasis and molecular markers linked to earliness in bread wheat. *Egyptian Journal of Plant Breeding*, vol. 14, no. 2, pp. 265-282.
- AL-NAGGAR, A.M.M., ABDEL-RAOUF, M.S., EL-BORHAMY, H.S. and SHEHAB-EL-DEEN, M.T., 2012. Gene effects controlling inheritance of earliness and yield traits of bread wheat under drought stress conditions. *Egyptian Journal of Plant Breeding*, vol. 16, no. 3, pp. 41-59.
- AL-NAGGAR, A.M.M., AL-AZAB, K.F., SOBIEH, S.E.S. and ATTA, M.M.M., 2015. Morphological and SSR assessment of putative drought-tolerant M<sub>3</sub> and F<sub>3</sub> families of wheat (*Triticum aestivum* L.). *British Biotechnology Journal*, vol. 6, no. 4, pp. 174-190. <http://dx.doi.org/10.9734/BBJ/2015/15898>.
- AL-NAGGAR, A.M.M., ATTA, M.M.M., SOBIEH, S.E.S. and AL-AZAB, K.F., 2013. Drought tolerant mutant induction via gamma rays in bread wheat. *Egyptian Journal of Plant Breeding*, vol. 17, no. 4, pp. 129-159.
- AMIN, I.A., 2013. Genetic behavior of some agronomic traits in two durum wheat crosses under heat stress. *Alexandria Journal of Agricultural Research*, vol. 58, no. 1, pp. 53-66.
- ATAEI, R., GHOLAMHOSEINI, M. and KAMALIZADEH, M., 2017. Genetic analysis for quantitative traits in bread wheat exposed to irrigated and drought stress conditions. *FYTON*, vol. 86, pp. 228-235.
- BENNETT, D., IZANLOO, A., REYNOLDS, M., KUCHEL, H., LANGRIDGE, P. and SCHNURBUSCH, T., 2012. Genetic dissection of grain yield and physical grain quality in bread wheat (*Triticum aestivum* L.) under water-limited environments. *Theoretical and Applied Genetics*, vol. 125, no. 2, pp. 255-271. <http://dx.doi.org/10.1007/s00122-012-1831-9>. PMID:22374139.
- BI, H., KOVALCHUK, N., LANGRIDGE, P., TRICKER, P.J., LOPATO, S. and BORISJUK, N., 2017. The impact of drought on wheat leaf cuticle properties. *BMC Plant Biology*, vol. 17, no. 1, pp. 85. <http://dx.doi.org/10.1186/s12870-017-1033-3>. PMID:28482800.
- BILGIN, O., KUTLU, Đ. and BALKAN, A., 2016. Gene effects on yield and quality traits in two bread wheat (*T. aestivum* L.) crosses. *International Journal of Crop Science and Technology*, vol. 2, no. 1, pp. 1-10.
- BÖRNER, A., SCHUMANN, E., FÜRSTE, A., CÖSTER, H., LEITHOLD, B., RÖDER, M. and WEBER, W., 2002. Mapping of quantitative trait loci determining agronomic important characters in hexaploid wheat (*Triticum aestivum* L.). *Theoretical and Applied Genetics*, vol. 105, no. 6-7, pp. 921-936. <http://dx.doi.org/10.1007/s00122-002-0994-1>. PMID:12582918.

- DARWISH, I.H.I. and EL-HOSARY, A.A., 2003. Response of wheat varieties to nitrogen under rainfed and irrigation. *Egyptian Journal of Plant Breeding*, vol. 7, no. 1, pp. 241-251.
- DERE, S. and YILDIRIM, M.B., 2006. Inheritance of grain yield per plant, flag leaf width and length in an 8 × 8 diallel cross-population of bread wheat (*Triticum aestivum* L.). *Turkish Journal of Agriculture and Forestry*, vol. 30, pp. 339-345.
- EBERCON, A., BLUM, A. and JORDAN, W.R., 1977. A rapid colorimetric method for epicuticular wax content of sorghum leaves. *Crop Science*, vol. 17, no. 1, pp. 179-180. <http://dx.doi.org/10.2135/cropsci1977.0011183X001700010047x>.
- EL-AREF, K.A., TAMMAM, A.M., IBRAHIM, M.M. and KOUBISY, Y.S., 2011. Generation mean analysis in bread wheat under drought conditions. *Egyptian Journal of Applied Science*, vol. 26, no. 2, pp. 187-208.
- EL-HAWARY, M.M. and EL-SHAFFEY, A.I., 2016. Improving rice productivity by potassium fertilization under saline soils conditions. *Journal of Plant Production, Mansoura University*, vol. 7, no. 7, pp. 677-682.
- ERKUL, A., UNAY, A. and KONAK, C., 2010. Inheritance of yield and yield components in a bread wheat (*Triticum aestivum* L.) cross. *Turkish Journal of Field Crops*, vol. 15, no. 2, pp. 137-140.
- ESMAIL, R.M. and KHATTAB, S.A.M., 2002. Genetic behavior of yield and its components in three bread wheat crosses. *Minufiya Journal of Agricultural Research*, vol. 27, no. 2, pp. 215-224.
- FARAG, H.I.A., 2009. Inheritance of yield and its components in bread wheat (*Triticum aestivum* L.) using six parameter model under Ras Sudr conditions. In: *Proceedings of the 6th International Plant Breeding Conference*, 2009, Ismailia, Egypt. Ismailia, pp. 90-112.
- FARSHADFAR, E., GHANADHA, SUTKA, J. and ZAHRAVI, M., 2001. Generations mean analysis of drought tolerance in wheat (*Triticum aestivum* L.). *Acta Agronomica Hungarica*, vol. 49, no. 1, pp. 59-66. <http://dx.doi.org/10.1556/AAgr.49.2001.1.7>.
- FARSHADFAR, E., RAFIEE, F. and HASHEMINASAB, V., 2013. Evaluation of genetic parameters of agronomic and morpho-physiological indicators of drought tolerance in bread wheat (*Triticum aestivum* L.) using diallel mating design. *AJCS*, vol. 7, no. 2, pp. 268-275.
- FEBRERO, A., FERNÁNDEZ, S., MOLINA-CANO, J.L. and ARAUS, J.L., 1998. Yield, carbon isotope discrimination, canopy reflectance and cuticular conductance of barley isolines of differing glaucousness. *Journal of Experimental Botany*, vol. 49, no. 326, pp. 1575-1581. <http://dx.doi.org/10.1093/jxb/49.326.1575>.
- FELTAOUS, Y.M., SOLIMAN, G.M.M. and EL-RAWY, M.A., 2020. Genotyping and phenotyping for some bread wheat genotypes terminal heat stress. *Egyptian Journal of Plant Breeding*, vol. 24, no. 1, pp. 195-223.
- FONSECA, S. and PATTERSON, F.L., 1968. Hybrid vigour in a seven parents diallel cross in common winter wheat (*Triticum aestivum* L.). *Crop Science*, vol. 8, no. 1, pp. 85-88. <http://dx.doi.org/10.2135/cropsci1968.0011183X000800010025x>.
- FOOD AND AGRICULTURE ORGANIZATION – FAO, 2021 [viewed 3 March 2021]. *FAOSTAT* [online]. Available from: <http://www.fao.org/faostat/en/#data/QC>
- HABIB, I. and KHAN, A.S., 2003. Genetic model of some economic traits in bread wheat (*Triticum aestivum* L.). *Asian Journal of Plant Science*, vol. 2, no. 17, pp. 1153-1154. <http://dx.doi.org/10.3923/ajps.2003.1153.1155>.
- HAMAM, K.A. and NEGIM, O., 2014. Evaluation of wheat genotypes and some soil properties under saline water irrigation. *Annals of Agricultural Science*, vol. 59, no. 2, pp. 165-176. <http://dx.doi.org/10.1016/j.aos.2014.11.002>.
- HASSAN, S.E., KHALIQ, I. and KHAN, A.S., 2008. Genetic mechanism of some physiological traits in spring wheat at two plant population regimes. *Journal of Agricultural Research*, vol. 46, pp. 395-401.
- HAYMAN, B.I. and MATHER, K., 1955. The description of gene interaction in continuous variation. *Biometrics*, vol. 11, no. 1, pp. 69-82. <http://dx.doi.org/10.2307/3001481>.
- HAYMAN, B.I., 1958. The separation of epistatic from additive and dominance variation in generation means. *Heredity*, vol. 12, no. 3, pp. 371-391. <http://dx.doi.org/10.1038/hdy.1958.36>.
- HUSSEIN, H.A., HUSSEIN, E.H.A., AL-NAGGAR, A.M.M., SABRY, S.R. and GAD, K.I.M., 2009. Genetic analysis of earliness and grain yield in two wheat crosses. *Egyptian Journal of Plant Breeding*, vol. 13, pp. 371-381.
- INAMULLAH, M.F. and HASSAN, G., 2005. Genetics of important traits in bread wheat using diallel analysis. *Sarhad Journal of Agriculture*, vol. 21, pp. 617-622.
- JAISWAL, R., GAUR, S.C. and JAISWAL, S.K., 2018. Heterosis and inbreeding depression for grain yield and yield component traits in bread wheat (*Triticum aestivum* L.). *Journal of Pharmacognosy and Phytochemistry*, vol. 7, no. 2, pp. 3586-3594.
- JENSEN, N.F. and DRISCOLL, C.J., 1962. Inheritance of the waxless character in wheat. *Crop Science*, vol. 2, no. 6, pp. 504-505. <http://dx.doi.org/10.2135/cropsci1962.0011183X000200060019x>.
- JINKS, J.L. and HAYMAN, B.I., 1953. The analysis of diallel crosses. *Maize Genetic Cooperative News*, vol. 1, pp. 48-54.
- JOHNSON, D.A., RICHARDS, R.A. and TURNER, N.C., 1983. Yield, water relations, gas exchange, and surface reflectances of near-isogenic wheat lines differing in glaucousness. *Crop Science*, vol. 23, no. 2, pp. 318-325. <http://dx.doi.org/10.2135/cropsci1983.0011183X002300020033x>.
- JOHNSON, H.W., ROBINSON, H.F. and COMSTOCK, R.E., 1955. Estimates of genetic and environmental variability in soybean. *Agronomy Journal*, vol. 47, no. 7, pp. 314-318. <http://dx.doi.org/10.2134/agronj1955.00021962004700070009x>.
- KEARSEY, M.J. and POONI, H.S., 1996. *The genetical analysis of quantitative traits*. London: Chapman & Hall, 396 p. <http://dx.doi.org/10.1007/978-1-4899-4441-2>.
- KHALED, M.A.I., 2007. Estimation of genetic variance for yield and yield components in two bread wheat (*Triticum aestivum* L.) crosses. *Journal of Agricultural Science, Mansoura University*, vol. 32, no. 10, pp. 8043-8053.
- KHATTAB, S.A.M., SHAHEEN, A.M.A. and AFIAH, S.A.N., 2001. Genetic behavior of some metric traits in four bread wheat crosses under normal and saline conditions. *Journal of Agricultural Science, Mansoura University*, vol. 26, no. 1, pp. 217-229.
- KOUBISY, Y.S.I., 2019. Generation mean analysis in two bread wheat crosses under heat stress conditions. *Egyptian Journal of Agricultural Research*, vol. 97, no. 2, pp. 589-607.
- KULWAL, P.L., ROY, J.K., BALYAN, H.S. and GUPTA, P.K., 2003. QTL mapping for growth and leaf characters in bread wheat. *Plant Science*, vol. 164, no. 2, pp. 267-277. [http://dx.doi.org/10.1016/S0168-9452\(02\)00409-0](http://dx.doi.org/10.1016/S0168-9452(02)00409-0).
- KUMAR, A., RAZDAN, A.K., SHARMA, V., KUMAR, N. and KUMAR, D., 2018. Study of heterosis and inbreeding depression for economic and biochemical traits in bread wheat (*Triticum aestivum* L.). *Journal of Pharmacognosy and Phytochemistry*, vol. 7, no. 4, pp. 558-564.
- KUMAR, J., KUMAR, A., KUMAR, M., SINGH, S.K., SINGH, L. and SINGH, G.P., 2017. Heterosis and inbreeding depression in relation to heterotic parameters in bread wheat (*Triticum aestivum* L.) under late sown condition. *Journal of Wheat Research*, vol. 9, no. 1, pp. 32-41. <http://dx.doi.org/10.25174/2249-4065/2017/67482>.

- LIU, Q., NI, Z., PENG, H., SONG, W., LIU, Z. and SUN, Q., 2007. Molecular mapping of a dominant nonglucousness gene from synthetic hexaploid wheat (*Triticum aestivum* L.). *Euphytica*, vol. 155, no. 1-2, pp. 71-78. <http://dx.doi.org/10.1007/s10681-006-9302-5>.
- LJUBICIC, N., PETROVIC, S., DIMITRIJEVIC, M. and HRISTOV, N., 2016a. Genetic analysis of some important quantitative traits in bread wheat (*Triticum aestivum* L.). *Ekin Journal*, vol. 2, no. 2, pp. 47-53.
- LJUBICIC, N., PETROVIC, S., DIMITRIJEVIC, M. and HRISTOV, N., 2016b. Gene actions involved in the inheritance of yield-related traits in bread wheat (*Triticum aestivum* L.). *Emirates Journal of Food and Agriculture*, vol. 28, no. 7, pp. 477-484. <http://dx.doi.org/10.9755/ejfa.2016-02-117>.
- MAHPARA, S., HUSSAIN, S.T., IQBAL, J., NOORKA, I.R. and SALMAN, S., 2018. Analysis of generation means for some metric plant traits in two wheat (*Triticum aestivum* L.) hybrids. *Pure and Applied Biology*, vol. 7, no. 1, pp. 93-102. <http://dx.doi.org/10.19045/bspab.2018.70012>.
- MASON, R.E., MONDAL, S., BEECHER, F.W., PACHECO, A., JAMPALA, B., IBRAHIM, A.M.H. and HAYS, D.B., 2010. QTL associated with heat susceptibility index in wheat (*Triticum aestivum* L.) under short-term reproductive stage heat stress. *Euphytica*, vol. 174, no. 3, pp. 423-436. <http://dx.doi.org/10.1007/s10681-010-0151-x>.
- MATHER, K. and JINKS, J.L., 1971. *Biometrical genetics*. 2nd ed. London: Chapman & Hall. <http://dx.doi.org/10.1007/978-1-4899-3404-8>.
- MATHER, K. and JINKS, J.L., 1982. *Biometrical genetics: the study of continuous variation*. London: Chapman & Hall, 396 p.
- MATHER, K., 1949. *Biometrical genetics: the study of continuous variation*. London: Methuen and Co. Ltd..
- MENSHAWY, A.M.M., 2005. Genetic analysis for earliness components in some wheat genotypes of different photothermal response. *Egyptian Journal of Plant Breeding*, vol. 8, no. 1, pp. 31-37.
- MENSHAWY, A.M.M., 2007. Evaluation of some early bread wheat genotypes under different sowing dates: 1. Earliness characters. *Egyptian Journal of Plant Breeding*, vol. 11, no. 1, pp. 25-40.
- MERAH, O., DELEENS, E., SOUYRIS, I. and MONNEVEUX, P., 2000. Effect of glucousness on carbon isotope discrimination and grain yield in durum wheat. *Journal Agronomy & Crop Science*, vol. 185, no. 4, pp. 259-265. <http://dx.doi.org/10.1046/j.1439-037x.2000.00434.x>.
- MOHAMMAD, W., SHAH, Z., SHAH, S.M. and IQBAL, M.M., 2003. Rotational benefits of legumes to subsequent rainfed wheat in a low N soil. *Pakistan Journal of Soil Science*, vol. 22, pp. 19-27.
- MOUSSA, A.M., 2010. Estimation of epistasis, additive and dominance variation in certain bread wheat (*Triticum aestivum*, L.) crosses. *Journal of Plant Production, Mansoura University*, vol. 1, no. 12, pp. 1707-1719.
- NANDA, G.S., HAZARIKA, G.N. and GILL, K.S., 1981. Inheritance of heading date, plant height, ear length and spikelets per spike in an intervarietal cross of wheat. *Theoretical and Applied Genetics*, vol. 60, no. 3, pp. 167-171. <http://dx.doi.org/10.1007/BF00264523>. PMID:24276689.
- NINGHOT, C.J., BORATKAR, M.V., THAWARI, S.B. and POTDUKHE, N.R., 2016. Generation mean analysis for yield and yield components in wheat (*Triticum aestivum* L.). *International Journal of Genetics*, vol. 8, no. 4, pp. 204-206.
- NOVOSELOVIC, D., BARIC, M., DREZNER, G., GUNJACA, J. and LALIC, A., 2004. Quantitative inheritance of some wheat plant traits. *Genetics and Molecular Biology*, vol. 27, no. 1, pp. 2-98. <http://dx.doi.org/10.1590/S1415-47572004000100015>.
- PATEL, H.N., ABHISHEK, D., SHRIVASTAVA, A. and PATEL, S.R., 2018. Genetic analysis for heterotic traits in bread wheat (*Triticum aestivum* L.) using six parameters model. *International Journal of Current Microbiology and Applied Sciences*, vol. 7, no. 6, pp. 239-249. <http://dx.doi.org/10.20546/ijcmas.2018.706.029>.
- PRZULJ, N. and MLADENOV, N., 1999. Inheritance of grain filling duration in spring wheat. *Plant Breeding*, vol. 118, no. 6, pp. 517-521. <http://dx.doi.org/10.1046/j.1439-0523.1999.00436.x>.
- RAIKWAR, R.S., 2019. Genetic architecture of yield and quality traits in wheat (*Triticum aestivum* L.). *Indian Journal of Genetics*, vol. 79, no. 1, pp. 100-103. <http://dx.doi.org/10.31742/IJGPB.79.1.14>.
- RIAZ, R. and CHOWDHRY, M.A., 2003. Genetic analysis of some economic traits of wheat under drought conditions. *Asian Journal of Plant Science*, vol. 2, no. 10, pp. 790-796. <http://dx.doi.org/10.3923/ajps.2003.790.796>.
- RICHARDS, R., RAWSON, H. and JOHNSON, D., 1986. Glucousness in wheat: its development and effect on water-use efficiency, gas exchange and photosynthetic tissue temperatures. *Functional Plant Biology*, vol. 13, no. 4, pp. 465-473. <http://dx.doi.org/10.1071/PP9860465>.
- SAID, A.A., 2014. Generation mean analysis in wheat (*Triticum aestivum* L.) under drought stress conditions. *Annals of Agricultural Science*, vol. 59, no. 2, pp. 177-184. <http://dx.doi.org/10.1016/j.a0as.2014.11.003>.
- SALMI, M., BENMAHAMMED, A., BENDERRADJI, L., FELLAHI, Z.E.A., BOUZERZOUR, H., OULMI, A. and BENBELKACEM, A., 2019. Generation means analysis of physiological and agronomical targeted traits in durum wheat (*Triticum durum* Desf.) cross. *Revista Facultad Nacional de Agronomía*, vol. 72, no. 3, pp. 8971-8981. <http://dx.doi.org/10.15446/rfnam.v72n3.77410>.
- SAMUELS, L., KUNST, L. and JETTER, R., 2008. Sealing plant surfaces: cuticular wax formation by epidermal cells. *Annual Review of Plant Biology*, vol. 59, no. 1, pp. 683-707. <http://dx.doi.org/10.1146/annurev.arplant.59.103006.093219>. PMID:18251711.
- SAS INSTITUTE, 2011. SAS v9.3. Cary, NC: SAS Institute Inc.
- SILVA FERNANDES, A.M.S., BAKER, E.A. and MARTIN, J.T., 1964. Studies on plant culture, VI: the isolation and fractionation of cultivar waxes. *Annals of Applied Biology*, vol. 53, pp. 43-58.
- SOLIMAN, G.M.M., 2018. Genetical analysis for yield and yield component in durum wheat under different sowing dates. In: *Proceedings of the 7th Field Crops Conference*, 18-19 December 2018, Giza, Egypt. Giza, pp. 133-148.
- STUCKEY, J.R., 1972. *Inheritance of glucousness in wheat*. Sydney: University of New South Wales. Ph. D. Thesis in Wheat Plant Breeding.
- TAMMAM, A.M., 2005. Generation mean analysis in bread wheat under different environmental conditions. *Minufiya Journal of Agricultural Research*, vol. 30, no. 3, pp. 937-956.
- TSUNEWAKI, K. and EBANA, K., 1999. Production of near-isogenic lines of common wheat for glucousness and genetic basis of this trait. *Genes & Genetic Systems*, vol. 74, no. 2, pp. 33-41. <http://dx.doi.org/10.1266/ggs.74.33>.
- WARNER, J.N., 1952. A method for estimating heritability. *Agronomy Journal*, vol. 44, no. 9, pp. 427-430. <http://dx.doi.org/10.2134/agronj1952.00021962004400080007x>.
- YADAV, H.K. and SINGH, S.P., 2011. Inheritance of quantitative traits in opium poppy (*Papaver somniferum* L.). *Genetika*, vol. 43, no. 1, pp. 113-128. <http://dx.doi.org/10.2298/GENSR1101113Y>.
- ZAAZAA, E.I., HAGER, M.A. and EL-HASHASH, E.F., 2012. Genetical analysis of some quantitative characters in wheat using six parameters genetic model. *American-Eurasian Journal of Agricultural & Environmental Sciences*, vol. 12, no. 4, pp. 456-462.