

Research Paper

## Biodegradation of cypermethrin by immobilized cells of *Micrococcus* sp. strain CPN 1

Preeti N. Tallur, Sikandar I. Mulla, Veena B. Megadi, Manjunatha P. Talwar,  
Harichandra Z. Ninnekar

Department of Biochemistry, Karnatak University, Karnataka, India.

Submitted: May 24, 2013; Approved: November 28, 2014.

---

### Abstract

Pyrethroid pesticide cypermethrin is an environmental pollutant because of its widespread use, toxicity and persistence. Biodegradation of such chemicals by microorganisms may provide a cost-effective method for their detoxification. We have investigated the degradation of cypermethrin by immobilized cells of *Micrococcus* sp. strain CPN 1 in various matrices such as, polyurethane foam (PUF), polyacrylamide, sodium alginate and agar. The optimum temperature and pH for the degradation of cypermethrin by immobilized cells of *Micrococcus* sp. were found to be 30 °C and 7.0, respectively. The rate of degradation of 10 and 20 mM of cypermethrin by freely suspended cells were compared with that of immobilized cells in batches and semi-continuous with shaken cultures. PUF-immobilized cells showed higher degradation of cypermethrin (10 mM and 20 mM) than freely suspended cells and cells immobilized in other matrices. The PUF-immobilized cells of *Micrococcus* sp. strain CPN 1 retained their degradation capacity. Thus, they can be reused for more than 32 cycles, without losing their degradation capacity. Hence, the PUF-immobilized cells of *Micrococcus* sp. could potentially be used in the bioremediation of cypermethrin contaminated water.

**Key words:** degradation, immobilization, cypermethrin, polyurethane foam, *Micrococcus* sp. strain CPN 1.

---

### Introduction

Cypermethrin is a synthetic pyrethroid pesticide widely used to control pests in cotton, fruits, and vegetable crops, (Tullur *et al.*, 2008; Chen *et al.*, 2012). In recent years, the use of cypermethrin has been increased sharply in the agricultural field for crop protection (Weston *et al.*, 2009). As a result, cypermethrin residues were found in most of the tested samples (sediment) from urban creek (Riederer *et al.*, 2010; Weston *et al.*, 2009, 2011). Hence, the presence of cypermethrin in the environment is of global concern. Cypermethrin has been shown to be genotoxic, neurotoxic, immunotoxic and carcinogenic to mammals, including humans (Ansari *et al.*, 2011; Jin *et al.*, 2011; McKinlay *et al.*, 2008; Wang *et al.*, 2011; Wolansky and Harrill, 2008; Zhang *et al.*, 2010). There are reports of degradation of cypermethrin by the free cells of microorganisms (Roberts and Standen, 1981; Jilani and Khan,

2006; Tallur *et al.*, 2008; Zhang *et al.*, 2010; Lin *et al.*, 2011; Chen *et al.*, 2012). In addition, degradation of cypermethrin by physico-chemical methods such as oxidation with ozone, photolysis, ultrasonic degradation, Fenton degradation, incineration and adsorption has also been investigated (Segal-Rosenheimer and Dubowski, 2007; Xie *et al.*, 2011). However, these methods were found to be more expensive and less-effective than biological systems used for the remediation of toxic pollutants (Yang *et al.*, 2011). In nature, cypermethrin may be degraded through several ways, including hydrolysis, volatilization, photolysis, and aerobic degradation by microorganisms. There is a report on the enhanced degradation of cypermethrin by mixed microbial culture (Chen *et al.*, 2012). The use of freely suspended cells for the degradation of various toxic/hazardous compounds for industrial applications has a number of disadvantages. It is mostly because of low mechanical strength, low density of cell population and the difficulty in

biomass effluent separation (Massalha *et al.*, 2007; Wang and Hu, 2007). Immobilization techniques have now been well established to overcome these problems (Cassidy *et al.*, 1996). Immobilized microbial cells have more advantages than freely suspended cells as well as enzymes immobilized in various matrices (Zheng *et al.*, 2009). Moreover, the immobilizing materials like polyurethane foam (PUF), agar and sodium alginate are inert, non toxic to cells, inexpensive and practical (Mulla *et al.*, 2012). Use of immobilized cells permit the operation of bioreactors at flow rates that are independent of the microorganisms, tolerate higher concentrations of toxic compounds than do their non immobilized counterparts (Westmeier *et al.*, 1987; Zheng *et al.*, 2009). The calcium alginate beads act like a slow release delivery system, where the substrate is slowly released from culture medium to the cells for microbial mineralization without a significant impact on the surrounding environment. There are several reports on the potential use of immobilized bacterial cells in different matrices for the degradation of numerous toxic/hazardous chemicals (Chen *et al.*, 2002; Tallur *et al.*, 2009; Ullah *et al.*, 2010; Mulla *et al.*, 2012; Ali *et al.*, 2013; Mulla *et al.*, 2013; Hoskeri *et al.*, 2014). However, less information is available on biodegradation of cypermethrin by immobilized cells of microorganisms

Previously, we have reported the degradative pathway of cypermethrin in *Micrococcus* sp. strain CPN 1 (Tallur *et al.*, 2008). In this paper, we describe the enhanced degradation of cypermethrin by the cells of *Micrococcus* sp. strain CPN 1 immobilized in sodium alginate (SA), agar, polyacrylamide and polyurethane foam (PUF). The rate of degradation of cypermethrin by the cells immobilized in various matrices were compared with that of freely suspended cells of *Micrococcus* sp. strain CPN 1 in batches and semi-continuous degradation.

## Materials and Methods

### Chemicals

Cypermethrin with 98% purity was purchased from Sigma-Aldrich. Sodium alginate, acrylamide, bis-acrylamide, ammonium persulphate with 99% purity were purchased from Himedia, India. Polyurethane foam (PUF) was obtained from local suppliers. All other chemicals used in this study were of analytical grade.

### Organism

*Micrococcus* sp. strain CPN 1 previously isolated and identified in our laboratory by cypermethrin enrichment culture technique was used in this study (Tallur *et al.*, 2008). The organism was maintained on the slants of cypermethrin-mineral salts medium solidified with 2% agar (wt/vol).

### Medium

Two different media were used in this study. The medium used for precultivation of *Micrococcus* sp. strain CPN 1 was a mineral salts medium (MM 1) containing (g L<sup>-1</sup>) K<sub>2</sub>HPO<sub>4</sub>, 6.30; KH<sub>2</sub>PO<sub>4</sub>, 1.82; NH<sub>4</sub>NO<sub>3</sub>, 1.00; MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.20; CaCl<sub>2</sub>·2H<sub>2</sub>O, 0.10; Na<sub>2</sub>MoO<sub>4</sub>·2H<sub>2</sub>O, 0.06; MnSO<sub>4</sub>·H<sub>2</sub>O, 0.06 and FeSO<sub>4</sub>·7H<sub>2</sub>O, 0.10 (Tallur *et al.*, 2008). The pH of the medium was adjusted to 7.0. One hundred mL aliquots of this medium were transferred into 500 mL Erlenmeyer flask and sterilized by autoclaving at 121 °C for 15 min. Cypermethrin (10 mM) dissolved in ethyl acetate (solvent) was sterilized by membrane filtration, added to the medium and kept on a rotary shaker for solvent evaporation. After two days, the organism was inoculated into the sterilized medium containing cypermethrin (10 mM) and this process was followed for all experiments.

The medium (MM 2) used for cypermethrin degradation studies contained (g L<sup>-1</sup>) K<sub>2</sub>HPO<sub>4</sub>, 6.30; MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.20; CaCl<sub>2</sub>·2H<sub>2</sub>O, 0.20; NH<sub>4</sub>NO<sub>3</sub>, 1.0 and FeCl<sub>3</sub>, 0.05 (Tallur *et al.*, 2009). The pH of the medium was adjusted to 7.0 and sterilized by autoclaving. Cypermethrin (10 mM and 20 mM) dissolved in ethyl acetate was added after sterilization of the medium. The cultures were incubated on a rotary shaker (150 rpm) at 30 °C. The bacterial cell concentrations were measured by the plate-count method (Mulla *et al.*, 2013).

### Immobilization of whole cells in various matrices

The cypermethrin-degrading organism (*Micrococcus* sp. strain CPN 1) was harvested during the mid-logarithmic growth phase from 4 L of culture medium. The cell suspension (3 × 10<sup>12</sup> cfu/mL) was obtained by centrifugation at 8,000 × g for 10 min at 4 °C and washed thrice with 50 mM phosphate buffer pH 7.0. The washed cells were immobilized in different matrices namely; Polyurethane foam (PUF), polyacrylamide, sodium alginate (SA) and agar (Mulla *et al.*, 2013).

**Preparation of PUF-entrapment:** The PUF was cut into approximately 5 mm cubes, washed thrice with sterile double distilled water and dried. 100 mL of bacterial cell suspensions (3 × 10<sup>12</sup> cfu/mL) were added to 500 mL conical flasks containing sterilized foam cubes (5 g). The content of the flasks were mixed by stirring on magnetic stirrer for 2 h and then the flasks were kept on a rotary shaker for 1 h at 150 rpm. The flasks were then allowed to stand undisturbed for additional 4 h. The medium was removed and foam cubes containing the immobilized bacteria were washed with saline. The decanted bacterial suspension and the saline wash were combined and the bacterial population in the mixture was counted by the plate-count method.

**Preparation of sodium alginate (SA) entrapment:** Sodium alginate (4%, wt/vol) was dissolved in boiling water and autoclaved at 121 °C for 15 min. 50 mL of bacterial cell suspension (18 g wet weight/ 50 mL sterilized SA solution)

was added to 200 mL sterilized sodium alginate solution (4%, wt/vol) and mixed by stirring on a magnetic stirrer. This sodium alginate cell mixture was extruded drop by drop cold, sterile  $\text{CaCl}_2$  (0.2 M) solution. Gel beads of approximately 4 mm diameter were obtained. The beads were hardened by resuspending in a fresh  $\text{CaCl}_2$  solution for 7 h with gentle agitation and then frozen at  $-18^\circ\text{C}$  for 24 h. Finally, these beads were washed several times with sterile double distilled water and stored at  $4^\circ\text{C}$  for further investigations.

**Preparation of polyacrylamide entrapment:** About 12 g wet cells were suspended in 10 mL distilled water and chilled in ice. To 10 mL of 0.2 M potassium phosphate buffer, (pH 7.0), 2.85 g acrylamide, 0.15 g bisacrylamide and 10 mg ammonium persulphate were added this buffer solution was mixed with the chilled cell suspension followed by the addition of 10  $\mu\text{L}$  of TEMED and poured into 2 or 3 glass petri dishes. It was then allowed for polymerization for 1 h. The sieved gels were suspended in 100 mL of 0.2 M potassium phosphate buffer pH 7.0 and allowed to settle.

**Preparation of agar entrapment:** 100 mg agar was dissolved in 4.5 mL of 0.9% (wt/vol) sodium chloride by heating at  $100^\circ\text{C}$  and then cooled to  $40^\circ\text{C}$ . Cell slurry was suspended in 0.9% (wt/vol) sodium chloride solution. 0.5 mL of the cell slurry was added to 4.5 mL of the agar solution and mixed. Immediately, the mixture was poured on nylon net placed on a glass plate and cooled to  $5^\circ\text{C}$ . The membrane was stored in 0.1 M phosphate buffer, pH 7.0.

## Degradation conditions

### Batch degradation experiments

The batch degradation experiments were performed to evaluate the degradation of cypermethrin by both freely suspended cells and immobilized cells in various matrices.

For freely suspended cell cultures, 10 mL of exponentially growing cells were inoculated into 90 mL of MM 2 medium in 500 mL Erlenmeyer flasks along with the same amount of heat-killed cells as controls. The cell concentration was adjusted ( $3 \times 10^{12}$  cfu/mL) and different amounts of cypermethrin (10 and 20 mM) were added. For immobilized cells, 12 g wet beads/foam cubes of the various matrices were added to a 500 mL Erlenmeyer flask containing 100 mL of mineral salts medium (MM 2) with 10 and 20 mM of cypermethrin. The cell counts in the immobilized matrices of the PUF, SA, polyacrylamide and agar were found to be  $1.7 \times 10^{12}$  cfu/g foam cubes,  $1.6 \times 10^{12}$  cfu/g,  $1.8 \times 10^{12}$  cfu/g and  $1.6 \times 10^{12}$  cfu/g beads, respectively. Both the freely suspended cell culture and the immobilized cells of *Micrococcus* sp. strain CPN 1 in various matrices were incubated at  $30^\circ\text{C}$  on a rotary shaker (150 rpm) under identical conditions along with controls. The samples from the culture broth were withdrawn under sterile conditions at different incubation periods and analyzed for residual

cypermethrin by HPLC (Metwally *et al.*, 1997). The rate of degradation of cypermethrin by freely suspended and PUF-immobilized cells of *Micrococcus* sp. strain CPN 1 at different pH (4.0-10.0) and temperature ( $25-45^\circ\text{C}$ ) were measured after 96 h of incubation. The storage stability of both freely suspended cells and PUF immobilized cells were tested up to 60 days at  $4^\circ\text{C}$ .

### Semi-continuous degradation

For establishing the longevity of degrading activity of immobilized cells in various matrices, repeated batch degradations were performed. After each cycle of incubation (96 h/cycle), the spent medium was decanted and beads/foam cubes were washed with sterile water and transferred into fresh MM2 containing cypermethrin. The degradation process was performed under identical conditions as described above and the residual cypermethrin in the spent medium was analyzed (Metwally *et al.*, 1997).

## Analytical methods

Cypermethrin concentration in the spent medium was determined by High Performance Liquid Chromatography (HPLC) as described in Metwally *et al.* (1997). At regular intervals, 5 mL samples were withdrawn and centrifuged at  $8,000 \times g$  for 10 min. The supernatants were extracted with ethyl acetate and the residue obtained after solvent evaporation was dissolved in methanol and used for HPLC analysis. 10  $\mu\text{L}$  of each residual sample of cypermethrin was analyzed by HPLC (Shimadzu, Japan) equipped with SPD-10AVP UV-Detector using shim-pack CLC- $\text{C}_8$  column (4.6 x 150 mm) of particle size (5  $\mu\text{m}$ ) (Phenomenex) and methanol-water (90:10, vol/vol) as mobile phase at the flow rate of  $1 \text{ mL min}^{-1}$ .


## Statistical analysis

All Experiments were carried out in triplicate and their results are presented as mean  $\pm$  standard deviations (SD).

## Results

### Degradation of cypermethrin of free and immobilized cells of *Micrococcus* sp. strain CPN 1 in batch cultures

The results in the enhanced degradation of 10 and 20 mM of cypermethrin in batch cultures, both by freely suspended cells and immobilized cells of *Micrococcus* sp. strain CPN 1 in polyurethane foam (PUF), sodium alginate, polyacrylamide and agar are shown in Figure 1A and B. With the initial concentration of 10 mM cypermethrin, the freely suspended cells degraded 9 mM of cypermethrin after 168 h incubation period, whereas immobilized cells degraded the same concentration within 96 h of incubation (Figure 1A). But with the increased initial concentration to 20 mM, the rate of degradation freely suspended cells was




**Figure 1** - Degradation of cypermethrin at 10 mM (A) and 20 mM (B) in batch cultures by cells of *Micrococcus* sp. strain CPN 1 immobilized on PUF (-♦-), polyacrylamide (-●-), sodium alginate (-▲-), agar (-■-), freely suspended cells (-+-) and uninoculated control (-x-). Data values represent means of triplicate and error bars indicate 95% confidence intervals.

decreased, whereas immobilized cells degraded the same concentration within 120 h (Figure 1B). These results suggest that higher concentration of cypermethrin was better tolerated and more rapidly degraded by immobilized cells than freely suspended cells.

#### Semi-continuous degradation of cypermethrin by immobilized cells of strain CPN 1


The results of the semi-continuous degradation of 10 and 20 mM cypermethrin by cells of *Micrococcus* sp. strain CPN 1 immobilized in PUF, alginate, polyacrylamide and agar are shown in Figure 2A and B. The PUF-immobilized cells can be reused for up to 32 cycles without losing their ability to degrade cypermethrin at the initial concentration of 10 and 20 mM. In contrast, agar, sodium alginate and polyacrylamide immobilized cells could be reused for 16, 20 and 24 cycles, respectively (Figure 2A). However, when the initial concentration of cypermethrin was increased to 20 mM, these immobilized cells could be reused with a decreased rate of degradation of cypermethrin (Figure 2B). These observations suggest that lower concentration of cypermethrin (10 mM) could be fed at much higher frequency than higher concentration of cypermethrin (20 mM).




**Figure 2** - Semi-continuous degradation of cypermethrin at 10 mM (A) and 20 mM (B) by cells of *Micrococcus* sp. strain CPN 1 immobilized on PUF (-♦-), polyacrylamide (-■-), sodium alginate (-▲-) and agar (-●-). Data values represent means of triplicate and error bars indicate 95% confidence intervals.

#### Effect of pH, temperature and storage stability on degradation capacity of PUF-immobilized cells of *Micrococcus* sp. strain CPN 1


The effect of pH on the degradation of cypermethrin by freely suspended cells and PUF immobilized cells shows that variations of the initial pH between 5.0 and 9.0 (Figure 3) had no effect on cypermethrin degradation (Figure 3). In contrast, the freely suspended cells were able to degrade cypermethrin at narrow range pH between 6.5 to 7.5. The effect of temperatures on the degradation of cypermethrin by PUF-immobilized cells showed higher activity at the temperature between 25 and 40 °C (Figure 4). Whereas the freely suspended cells showed activity at the temperature between 30 and 35 °C (Figure 4). However, the optimal pH and temperature were found to be 7.0 and 30 °C both for PUF-immobilized cells and freely suspended cells in culture medium. The PUF-immobilized cells can be stored for 60 days at 4 °C without loss of its degradation capacity while the freely suspended cells lost their degrading capacity after 60 days at 4 °C (Figure 5).



**Figure 3** - Effect of pH on the degradation of cypermethrin (10 mM) by freely suspended cells and PUF-immobilized cells of *Micrococcus* sp. strain CPN 1. Data values represent means of triplicate and error bars indicate 95% confidence intervals.



**Figure 4** - Effect of Temperature on the degradation of cypermethrin (10 mM) by freely suspended cells and PUF-immobilized cells of *Micrococcus* sp. strain CPN 1. Data values represent averages of three replicate determinations.



**Figure 5** - Storage stability of freely suspended cells (-◆-) and PUF-immobilized (-▲-) cells of *Micrococcus* sp. strain CPN 1 grown on cypermethrin (10 mM). Data values represent means of triplicate and error bars indicate 95% confidence intervals.

## Discussion

The degradation of cypermethrin by immobilized cells of *Micrococcus* sp. strain CPN 1 in various matrices such as PUF, sodium alginate, polyacrylamide and agar were compared with that of freely suspended cells in batches and semi-continuous shaken cultures. The results obtained from cells immobilized in various matrices with batch cultures suggested that the rate of degradation of cypermethrin, even at a higher concentration (20 mM), was much higher than that with freely suspended cells. The enhanced degradation of cypermethrin by immobilized cells may be due to availability of a high density of cells in or on immobilized matrices. Immobilization of cells may also lead to a stabilization of membrane permeability and protect against the toxicity of high substrate concentration, thus leading to enhanced degradation rate (Cassidy *et al.*, 1996).

The results of semi-continuous degradation suggests that the PUF and polyacrylamide immobilized cells retained the cypermethrin degradation capacity for a longer period and they could be reused for 32 and 24 cycles, respectively. When the initial concentration of cypermethrin (10 mM) was increased to 20 mM, the PUF immobilized cells could be reused without losing their degrading capacity. The immobilized cells in other matrices could also be reused but with decreased rate of degradation at higher concentration (20 mM). The storage stability and activity of cells entrapped in PUF were better than those cells entrapped in other matrices. The alginate and agar-immobilized cells showed lower degradability of cypermethrin with increased cycle numbers. The mechanical instability and gradual cell leakage from these beads decreased the degradation rate with an increasing cycle number (Trevors *et al.*, 1992; Tallur *et al.*, 2009; Mulla *et al.*, 2012). The PUF-immobilized cells showed more tolerance to pH and temperature changes than freely suspended cells. The advantages of PUF in chemical and physical properties compared to other matrices are its high porosity, mechanical strength, stability and adsorbing capacity. PUF is an ideal support of cell growth (Romaskevicius *et al.*, 2006). The cells immobilized in PUF showed a better and faster degradation rate even at higher initial concentration of substrate. These immobilized cells could be stored for longer periods without losing their degradation ability. Furthermore, the longevity of cells immobilized in PUF and their operational stability is better than those of other matrices determined.

The present study have revealed that the more effective degradation of cypermethrin at higher concentration could be achieved by immobilized cells of *Micrococcus* sp. strain CPN 1 than freely suspended cells. The immobilized microbial system has an advantage of enhanced rate of degradation, tolerance to higher substrate concentrations and their reuseability. Thus, the immobilized microbial technology provides a highly versatile and cost-effective ap-

proach that can be used for degradation of pesticide contaminated wastewater.

## References

- Ali O, Namane A, Hellal A (2013) Use and recycling of Calcium alginate biocatalyst for removal of phenol from wastewater. *J Ind Eng Chem* 19:1384-1390.
- Ansari AR, Rahman S, Kaur M *et al.* (2011) *In vivo* cytogenetic and oxidative stress-inducing effects of cypermethrin in freshwater fish, *Channa punctata* Bloch. *Ecotoxicol Environ Saf* 74:150-156.
- Cassidy MB, Lee H, Trevors JT (1996) Environmental applications of immobilized microbial cells: a review. *J Ind Microbiol* 16:79-101.
- Chen KC, Lin YH, Chen WH *et al.* (2002) Degradation of phenol by immobilized *Candida tropicalis*. *Enzyme Microbiol Technol* 31:490-497.
- Chen S, Luo J, Hu M *et al.* (2012) Enhancement of cypermethrin degradation by a coculture of *Bacillus cereus* ZH-3 and *Streptomyces aureus* HP-S-01. *Bioresour Technol* 110:97-104.
- Hall DO, Rao KK (1989) Immobilized photosynthetic membranes and cells for the production of fuels and chemicals. *Chemosci* 1:41-47.
- Hoskeri RS, Mulla SI, Ninnekar HZ (2014) Biodegradation of chloroaromatic pollutants by bacterial consortium immobilized in polyurethane foam and other matrices. *Biocatal Agric Biotechnol* 3 (4): 390-396 <http://dx.doi.org/10.1016/j.bcab.2014.03.001i>.
- Jilani S, Khan A (2006) Biodegradation of Cypermethrin by *Pseudomonas* in a batch activated sludge process. *Int J Environ Sci Tech* 3:371-380. Bottom of Form
- Jun YX, Zheng SS, Fu ZW (2011) Embryonic exposure to cypermethrin induces apoptosis and immunotoxicity in zebrafish (*Danio rerio*). *Fish Shellfish Immunol* 30:1049-1054.
- Lin QS, Chen SH, Hu MY *et al.* (2011) Biodegradation of cypermethrin by a newly isolated actinomycetes HU-S-01 from wastewater sludge. *Int J Environ Sci Technol* 8:45-56.
- McKinlay R, Plant JA, Bell JN *et al.* (2008) Endocrine disrupting pesticides: implications for risk assessment. *Environ Int* 34:168-183.
- Massalha N, Basheer S, Sabbah I (2007). Effect of adsorption and bead size of immobilized biomass on the rate of biodegradation of phenol at high concentration levels. *Ind Eng Chem Res* 46:6820-6824.
- Metwally ME-S, Osman MS, Al-Rushaid R (1997) A high-performance liquid chromatographic method for the determination of cypermethrin in vegetables and its application to kinetic studies after greenhouse treatment. *Food Chemistry* 59:283-290.
- Mulla SI, Talwar MP, Hoskeri RS *et al.* (2012) Enhanced Degradation of 3-Nitrobenzoate by Immobilized Cells of *Bacillus flexus* strain XJU-4. *Biotech Bioprocess Eng* 17:1294-1299.
- Mulla SI, Talwar MP, Bagewadi ZK *et al.* (2013) Enhanced degradation of 2-nitrotoluene by immobilized cells of *Micrococcus* sp. strain SMN-1. *Chemosphere* 90:1920-1924.
- Romaskevicius T, Budriene S, Pielichowski K *et al.* (2006) Application of polyurethane-based materials for immobilization of enzymes and cells: A review. *CHEMJA* 17:74-89.
- Riederer AM, Hunter RE, Hayden SW *et al.* (2010) Pyrethroid and organophosphorus pesticides in composite diet samples from Atlanta, USA adults. *Environ Sci Technol* 44:483-490.
- Roberts TR, Standen ME (1981) Further studies of the degradation of the pyrethroid insecticide cypermethrin in soils. *Pest Science* 12:285-296.
- Segal-Rosenheimer M, Dubowski Y (2007) Heterogeneous ozonolysis of cypermethrin using real-time monitoring FTIR techniques. *J Phys Chem C* 111:11682-11691.
- Tallur PN, Megadi VB, Ninnekar HZ (2008) Biodegradation of cypermethrin by *Micrococcus* sp. strain CPN1. *Biodegradation* 19:77-82.
- Tallur PN, Megadi VB, Ninnekar HZ (2009) Biodegradation of p-cresol by immobilized cells of *Bacillus* sp. strain PHN 1. *Biodegradation* 20:79-83.
- Trevors JT, VanElsa JD, Lee H *et al.* (1992) Use of alginate and other carriers for encapsulation of microbial cells for use in soil. *Microb Release* 1:61-69.
- Ullah H, Shah AA, Hasan F *et al.* (2010) Biodegradation of trinitrotoluene by immobilized *Bacillus* sp. YRE1. *Pak J Bot* 42:3357-3367.
- Wang BE, Hu YY (2007) Comparison of four supports for adsorption of reactive dyes by immobilized *Aspergillus fumigatus* beads. *J Environ Sci* 19:451-457.
- Wang H, Wang SF, Ning H *et al.* (2011) Maternal cypermethrin exposure during lactation impairs testicular development and spermatogenesis in male mouse offspring. *Inc Environ Toxicol* 26:382-394.
- Westmeier F, Rehm HJ (1987) Degradation of 4-chlorophenol in municipal wastewater by adsorptive immobilized *Alcaligenes* sp. A7-2. *Appl Microbiol Biotechnol* 26:78-83.
- Weston DP, Holmes RW, Lydy MJ (2009) Residential runoff as a source of pyrethroid pesticides to urban creeks. *Environ Pollut* 157:287-294.
- Weston DP, Asbell AM, Hecht SA *et al.* (2011) Pyrethroid insecticides in urban salmon streams of the Pacific Northwest. *Environ Pollut* 159:3051-3056.
- Wolansky MJ, Harrill JA (2008) Neurobehavioral toxicology of pyrethroid insecticides in adult animals: a critical review. *Neurotoxicol Teratol* 30:55-78.
- Xie JM, Wang PL, Liu J *et al.* (2011) Photodegradation of lambda-cyhalothrin and cypermethrin in aqueous solution as affected by humic acid and/or copper: intermediates and degradation pathways. *Environ Toxicol Chem* 30:2440-2448.
- Yang L, Chen SH, Hu MY *et al.* (2011) Biodegradation of carbofuran by *Pichia anomala* strain HQ-C-01 and its application for bioremediation of contaminated soils. *Biol Fertil Soils* 47:917-923.
- Zhang C, Jia L, Wang S *et al.* (2010) Biodegradation of Beta-cypermethrin by two *Serratia* spp. with different cell surface hydrophobicity. *Bioresour Technol* 101:3423-3429.
- Zheng C, Zhou J, Wang J *et al.* (2009) Aerobic degradation of nitrobenzene by immobilized *Rhodotorula mucilaginosa* in polyurethane foam. *J Haz Mat* 168:298-303.

Associate Editor: Rodrigo da Silva Costa