RETRACTION

The Editorial Board of **Radiologia Brasileira** is officially communicating the Retraction for extraction of the following article:

Lee JC, Teles MS. Prevalence of additional primary malignancies detected incidentally on PET/CT. Radiol Bras. 2020;53(1):69. https://dx.doi.org/10.1590/0100-3984.2019.0097.

Reason: Duplicate publication of an article published in a previous issue of Radiologia Brasileira:

Lee JC, Teles MS. Prevalence of additional primary malignancies detected incidentally on PET/CT. Radiol Bras. 2019;52(5):342. https://dx.doi.org/10.1590/0100-3984.2019.0097.

Prof. Edson Marchiori Editor-in-Chief, **Radiologia Brasileira**

Prevalence of additional primary malignancies detected incidentally on PET/CT

Dear Editor,

We entirely agree with the conclusions of Tibana et al.⁽¹⁾. Although false-positives can occur with ¹⁸F-fluorodeoxyglucose positron emission tomography/computed tomography (PET/CT), the prevalence of true-positives cannot be underestimated. Additional primary malignancies may often be identified by this means and the likelihood of cure is much increased if such malignancies are treated promptly and aggressively it was unclear what is the actual prevalence of malignancies discovered with PET/CT in this study.

The prevalence of additional malignancies discovered with PET/CT was highlighted in the various systems for breas (2) bowel⁽³⁾, prostate⁽⁴⁾ and thyroid⁽⁵⁾ lesions. For example, in study of 1665 patients, 70 incidentally detected lesions in the colon were identified and eventually 10 were diagnosed with bowel carcinoma⁽³⁾. There were additional pre-malignant lesions identified and early treatment was warranted in this scenario. Thus, it was found the prevalence of malignant and premalignant findings in the PET population was about 1.3%. A similar series showed a lower number of carcinomas but more adenomas (which would, of course, be considered pre-malignant) in a slightly larger series⁽⁶⁾. Hence, the prevalence was similar (at 1.1%). Knowing the prevalence may give a better idea of the importance of the incidental findings. This would be very helpful for both managing the individual patient as well as for public health and population health purposes.

REFERENCES

- Tibana TK, Santos RFT, Arão Filho A, et al. Detection of additional primary malignancies: the role of CT and PET/CT combined with multiple percutaneous biopsy. Radiol Bras. 2019;52:166–71.
- Bertagna F, Evangelista L, Piccardo A, et al. Multicentric study on ¹⁸F-FDG-PET/CT breast incidental uptake in patients studied for nonbreast malignant purposes. Rev Esp Med Nucl Imagen Mol. 2015;34: 24–9
- Lee JC, Hartnett GF, Hughes BG, et al. The segmental distribution and clinical significance of colorectal fluorodeoxyglucose uptake incidentally detected on PET-CT. Nucl Med Commun. 2009;30:333–7.
- Yang Z, Hu S, Cheng J, et al. Prevalence and risk of cancer of incidental uptake in prostate identified by fluorine-18 fluorodeoxyglucose positron emission tomography/computed tomography. Clin Imaging. 2014;38: 470–4.
- Adas M, Adas G, Koc B, et al. Incidental thyroid lesions on FDG-PET/CT: a prevalence study and proposition of management. Minerva Endocrinol. 2015;40:169–75.
- Kamel EM, Thumshirn M, Truninger K, et al. Significance of incidental 18F-FDG accumulations in the gastrointestinal tract in PET/CT: correlation with endoscopic and histopathologic results. J Nucl Med. 2004;5:1804–10.

Joseph C. Lee^{1,2,a}, Marcelo Santos Teles^{1,b}

 Department of Medical Imaging, The Prince Charles Hospital, Chermside, Queensland, 4032 Australia.
 Faculty of Medicine, University of Queensland, Herston Rd, Herston, Queensland, 4006 Australia.

Correspondence: Joseph C. Lee. Department of Medical Imaging, The Prince Charles Hospital, Chermside, Queensland, 4032 Australia. Email: Joseph.Lee@health.qld.gov.au.

https://orcid.org/0000-0002-7683-8825; b. https://orcid.org/0000-0002-2003-7279.

Received 25 July 2019. Accepted after revision 1 August 2019.

http://dx.doi.org/10.1590/0100.3984.2019.0097

(cc)) BY