



Original Article

Leaf venation pattern to recognize austral South American medicinal species of “cow's hoof” (*Bauhinia* L., Fabaceae)



Renée H. Fortunato^a, Beatriz G. Varela^b, María A. Castro^c, María J. Nores^{d,*}

^a Instituto de Recursos Biológicos (Instituto Nacional de Tecnología Agropecuaria – Consejo Nacional de Investigaciones Científicas y Técnicas) and Facultad de Agronomía y Ciencias Agroalimentarias, Universidad de Morón, Buenos Aires, Argentina

^b Facultad de Farmacia y Bioquímica, Universidad Nacional de Buenos Aires, Buenos Aires, Argentina

^c Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Buenos Aires, Buenos Aires, Argentina

^d Instituto Multidisciplinario de Biología Vegetal (Universidad Nacional de Córdoba – Consejo Nacional de Investigaciones Científicas y Técnicas) and Facultad de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de Córdoba, Córdoba, Argentina

ARTICLE INFO

Article history:

Received 6 September 2016

Accepted 28 October 2016

Available online 8 December 2016

Keywords:

Arealation

Botanical identification

Freely ending veinlets

Leaf venation pattern

Quality control

Schnella

ABSTRACT

The leaves extracts of some species of *Bauhinia* L. s.l. are consumed to treat diabetes, inflammation, pains and several disorders in traditional medicine in austral South America. Despite its wide use and commercialization, sale is not controlled, and botanical quality of samples is not always adequate because of plant misidentification and adulteration. Here, we characterized leaf vein pattern in nineteen taxa to contribute to the recognition and commercial quality control of plant material commercially available. The vein characters intercostal tertiary and quaternary vein fabric, areole development and shape, free ending veinlet branching and marginal ultimate venation allowed to distinguish the main medicinal species in the region.

© 2016 Sociedade Brasileira de Farmacognosia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The leaves extracts of some species of *Bauhinia* L., Fabaceae, so called cow's hoof, “pata-de-vaca”, cow's paw or orchid trees, are consumed to treat diabetes, inflammation, pains and several disorders in traditional medicine in austral South America (Oliveira and Saito, 1989; Bortoluzzi et al., 2006; Barboza et al., 2009; Pizzioli et al., 2011; Fortunato, 2012). With 300 species worldwide, *Bauhinia* s.l. is represented by fifteen species in Argentina, Paraguay, Uruguay and Santa Catarina, Paraná and Rio Grande do Sul states of Brazil (Fortunato, 1986, 1996; Fortunato et al., 2008; Vaz et al., 2010).

Bauhinia bauhinioides (Mart.) J.F.Macbr., *B. microstachya* (Raddi) J.F.Macbr., *B. unguis* L., *B. holophylla* (Bong.) Steud., *B. forficata* Link, *B. cheilantha* (Bong.) Steud. and *B. rufa* (Bong.) Steud. are the main species for which pharmacological activities have been reported. Hypoglycemic, antioxidant, analgesic, antiinflammatory and larvicidal activities along with antitumor potentiality let them auspicious targets for new bioactive substances researching and phytotherapy (Silva and Cechinel Filho, 2002; Sousa et al., 2004;

Gadotti et al., 2005; Luna et al., 2005; Nakahata et al., 2006; Cechinel Filho, 2009; Paula et al., 2014; Rozza et al., 2015; Martins-Olivera et al., 2016).

These taxa are extensively employed and sold in native American, rural and urban communities in the region and surrounding areas (Bortoluzzi et al., 2006; Barboza et al., 2009; Hurrell et al., 2011; Ibarrola and Degen de Arrúa, 2011). Despite its wide use and commercialization, sale is not controlled, and botanical quality of samples is not always adequate because of plant misidentification and adulteration. Commercial samples – crude herbs or industrialized herbal medicines – include entire or broken dried leaves, stems and often pods. In general, plants are harvested from their natural habitats, being popularly recognized by their bilobed or bifoliolate leaves. Species are difficult to identify, especially from vegetative stages or in fragmented material, and sample labeling and trading are mostly accomplished by using common names, generic name or incorrect names (Melo et al., 2004). These often result in adulterated, falsified or substitute samples, which would interfere with the effectiveness and even the safety of the product (Ferreres et al., 2012). Leaf anatomical features are useful for identification and adequate sample botanical quality, in *Bauhinia* s.l., venation pattern has been utilized in taxonomy, phylogenetic analyses and even in palaeobotany (Vaz, 1979; Fortunato, 1986; Zhang, 1994; Lin et al., 2015).

* Corresponding author.

E-mail: jnores@imbiv.unc.edu.ar (M.J. Nores).




Fig. 1. Leaf venation in southern South American *Bauhinia*. (A and B) General vein pattern. (C–G, I–P, R and S) Areolation. (H and Q) Looped marginal ultimate venation. (A) *B. bauhinioides*. (B) *B. hagenbeckii*. (C) *Bauhinia affinis*. (D) *B. angulosa*. (E) *B. argentinensis*. (F) *B. bauhinioides*. (G and H) *B. campestris*. (I) *B. cheilantha*. (J) *B. forficata* ssp. *pruinosa*. (K) *B. hagenbeckii*. (L) *B. holophylla*. (M) *B. microstachya*. (N) *B. mollis* var. *mollis*. (O) *B. mollis* var. *notophila*. (P and Q) *B. rufa*. (R) *B. unguulata* var. *ungulata*. (S) *B. uruguayensis*. Arrowhead: basal veins. Arrow: agrographic veins. Scale bar: A and B: 3 mm; C–G, I–P, R and S: 300 µm; H, Q: 1 mm.

In this work, we contribute to the recognition and commercial quality control of austral South American medicinal *Bauhinia* through a study of leaf vein pattern.

Materials and methods

We analyzed 19 austral South American taxa (Appendix 1). The study of leaf venation pattern was carried out following Ellis et al. (2009). For this, fresh or herbarium leaves were cleared according to Dizeo de Strittmatter (1973) and stained in safranin/80% ethanol. Samples were mounted in DPX (Aldrich Chemical Company, Gillingham, UK) and examined using Zeiss Stereo Microscope a Zeiss Axiolab light microscope (Carl Zeiss, Germany) with a trinocular phototube and a color digital imaging camera. Some venation data were extracted from previous work (Fortunato, 1986, 1996).

Results and discussion

All taxa analyzed present basal actinodromous primary vein framework, with no naked basal veins, 3–5 (7) basal veins per lobe or leaflet and agrophic veins simple (Fig. 1A and B). Previously, some of these taxa have been described as acrodromous (Fortunato, 1986, 1996) or campylodromous (Vaz, 1979), and recently both American *Bauhinia* s. s. and *Schnella* are considered actinodromous or acrodromous by Lin et al. (2015). Main vein characters (Fig. 1, Table 1) provide a criterion to identification of medicinal species and detection of falsifications or adulteration.

The areole development allows differentiating three main groups, some of them, but not all, taxonomically related.

The first group is characterized by a good areole development that is found in some medicinal members of sect. *Pauletia* ser. *Cansenia*, in the liana *Bauhinia angulosa* (= *S. angulosa*) and in the spiny tree *B. uruguayensis* (Fig. 1D, L, P–S). The medicinal *B. holophylla* and *B. rufa* both display quadrangular and pentagonal, rarely triangular areoles, with absent and unbranched freely ending veinlets (FEV), the first one can be further recognized since FEV in general cross the areola and marginal ultimate venation and quaternary vein fabric venation are absent (Fig. 1L). *B. rufa*, instead, presents looped marginal ultimate venation and quaternary vein fabric present (Fig. 1P and Q). The medicinal *B. unguulata* has absent, unbranched and one branched FEV and areola shape more variable (Fig. 1R). Some similar features have been described in others taxa from ser. *Cansenia*, and Zhang (1994) named it as the *Cansenia* venation type. *B. angulosa* is easily distinguished because of the very small areoles lacking FEV and marginal ultimate vein absent (Fig. 1D). *B. uruguayensis* is the only taxon that presents paxillate areole development without FEV, or just very few FEV (Fig. 1S), these features are similar to the *Elayuna* type described by Zhang (1994).

By the other side, the second group, composed by the prickly trees and shrubs from sect. *Pauletia* ser. *Aculeatae* and the suffrutescences and shrubs from ser. *Pentandrae*, possesses moderate areole development along with marginal ultimate venation absent and quaternary vein fabric present (Fig. 1B–E, J–K, M–O). Comparatively, areoles are larger and more variable in size than in previous

Table 1

Leaf venation pattern in austral South American *Bauhinia*. FEV, freely ending veinlets; P, pentagonal; PE, percurrent; Q, quadrangular; R, reticulate; T, triangular.

Species	Intercostal tertiary vein fabric	Quinternary vein fabric	Areole development	Free ending veinlet (FEV) branching	Areole shape	Marginal ultimate venation	Fig. 1
<i>B. affinis</i>	PE	Present	Moderate	Unbranched and one or two branched; rarely absent	variable	Absent	C
<i>B. amambayensis</i>	R at apex, PE at base	Present	Moderate	Unbranched and with one or more branches; sometimes absent	variable	Absent	
<i>B. angulosa</i> (= <i>S. angulosa</i>)	R at apex, PE at base	Absent	Good	Absent; rarely unbranched	Q and P, rarely T	Absent	D
<i>B. argentinensis</i>	R at apex, PE at base	Present	Moderate	Mostly with one or more branches; sometimes unbranched	Variable	Absent	E
<i>B. bauhinoides</i>	R	Absent, to poor distinct	Moderate to good	Mostly unbranched and one branched; rarely two branched or absent	Variable	Absent	A, F
<i>B. campestris</i>	PE to R	Absent	Moderate, rarely good	Mostly with one or more branches; sometimes unbranched; rarely absent	Variable	Looped	G and H
<i>B. cheilantha</i>	PE	Absent	Good to moderate	Absent, unbranched and one branched; rarely two branched.	Variable	Absent	I
<i>B. forficata</i>	PE, notably at base	Present	Moderate	Mostly with one or more branches; sometimes unbranched	Variable	Absent	J
<i>B. hagenbeckii</i>	R at apex, PE at base	Present	Moderate	Mostly with one and more branches; rarely unbranched	Variable	Absent	B, K
<i>B. holophylla</i>	PE	Absent	Good	Absent and unbranched (FEVs mostly crossing areoles); rarely one branched	Q and P, rarely T	Absent	L
<i>B. microstachya</i> (= <i>S. microstachya</i>)	R at apex, PE at base	Absent, to poor distinct	Moderate	Mostly unbranched and one branched; sometimes with more branches	Variable	Looped and absent	M
<i>B. mollis</i> var. <i>mollis</i>	PE to R	Present	Moderate	Mostly unbranched and one branched; sometimes with more branches or absent	Variable	Absent	N
<i>B. mollis</i> var. <i>notophila</i>	PE	Poor distinct	Moderate	Mostly unbranched and with one or more branches; rarely absent	Variable	Absent	O
<i>B. rufa</i>	PE	Present	Good	Absent and unbranched	Q and P, rarely T	Looped	P and Q
<i>B. ungulata</i> var. <i>cuiabensis</i>	PE to R at apex	Absent	Good	Absent, unbranched and one branched	Variable to Q, P, T	Looped and absent	
<i>B. ungulata</i> var. <i>ungulata</i>	PE to R at apex	Absent, to poor distinct	Good	Absent, unbranched and one branched	Variable to Q, P, T	Looped	R
<i>B. uruguayensis</i>	R at apex, PE at base	Present	Good/paxillate	Absent; rarely unbranched and one branched	Q and P, rarely T	Looped and in part absent	S

group. The medicinal *B. forficata* (both subspecies), *B. argentinensis* and *B. hagenbeckii* show almost the same pattern, with mostly branching FEV (one or more branches) and less frequently, unbranched FEV (Fig. 1J, E, K). *B. forficata* can be distinguished by the percurrent tertiary vein fabric, while the others are percurrent at the base and reticulate at apex. *B. affinis* (Fig. 1C), *B. amambayensis* (not shown) and *B. mollis* (Fig. 1N, O) have in general both unbranched and branched FEV, in different frequency. The medicinal liana *B. microstachya* (= *S. microstachya*) possesses also moderate areole development, mostly unbranched and one branched FEV, but is recognized because of the presence of both looped and absent marginal ultimate venation (Fig. 1M), this report is similar to the one by Fortunato (1986) but differs from previous (Zhang, 1994).

Finally, the medicinal *B. cheilantha* and *B. campestris* (sect. *Pauletia* ser. *Cansenia*) present areole development intermediate between good and moderate. *B. cheilantha* alternates between absent, unbranched and one branched FEV and ultimate venation absent while *B. campestris* can be distinguished because FEV are in general branched and ultimate venation is looped (Fig. 1I, G and H). The medicinal *B. bauhinoides* (sect. *Pauletia* ser. *Perlebia*) also possess areole development moderate to good but areoles are mostly unbranched and one branched (Fig. 1F).

Interestingly, all species present prismatic crystals of calcium oxalate associated to vascular bundles and often druses in mesophyll cells, more notable in *B. affinis*, *B. argentinensis*, *B. mollis* and *B. ungulata* var. *cuiabensis*.

Here we provide a helpful tool for the recognition of the main regional medicinal plants through the analysis of the leaf vein pattern based on the intercostal tertiary and quaternary vein fabric, areole development and shape, free ending veinlet branching and marginal ultimate venation characters.

Authors' contributions

RHF collected and identified the plants, performed the laboratory work and data analysis and participated in the results discussion and the manuscript final writing. MAC participated in the laboratory work and data analysis. BGV helped perform the analysis with constructive discussions. MJN drafted the manuscript.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

We thank Cristo S. for technical assistance. CONICET (PIP 112-201101-00250), INTA (PNHFA-1106094) and SECyT-UNC (32720160200161CB) are acknowledged for financial support.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:[10.1016/j.bjph.2016.10.007](https://doi.org/10.1016/j.bjph.2016.10.007).

References

- Barboza, G.E., Cantero, J.J., Núñez, C., Pacciaroni, A., Espinar, L.A., 2009. *Medicinal plants: a general review and a phytochemical and ethnopharmacological screening of the native Argentine Flora*. *Kurtziana* 34, 7–365.
- Bortoluzzi, R.L.C., Miotto, S.T.S., Reis, A., 2006. Bauhinia. In: Reis, A. (org.), *Flora Ilustrada Catarinense*, vol. 2. Herbário Barbosa Rodrigues, Itajaí, pp. 29–67.
- Cechinel Filho, V., 2009. *Chemical composition and biological potential of plants from the genus Bauhinia*. *Phytother. Res.* 23, 1347–1354.
- Dízeo de Strittmatter, C.G., 1973. *Nueva técnica de diafanización*. *Bol. Soc. Argent. Bot.* 15, 126–129.
- Ellis, B., Daily, D.C., Hickey, L.J., Johnson, K.R., Mitchell, J.D., Wilf, P., Wing, S.L., 2009. *Manual of Leaf Architecture*. Cornell University Press, New York.
- Ferrerres, F., Gil-Izquierdo, A., Vinholes, J., Silva, S.T., Valentão, P., Andrade, P.B., 2012. *Bauhinia forficata* Link authenticity using flavonoids profile: relation with their biological properties. *Food Chem.* 134, 894–904.
- Fortunato, R.H., 1986. Revisión del género *Bauhinia* (Cercideae, Caesalpinoideae, Fabaceae). *Darwiniana* 27, 527–557.
- Fortunato, R.H., 1996. *Bauhinia affinis* (Fabaceae) una nueva cita para la Flora Argentina. *Darwiniana* 34, 405–409.
- Fortunato, R.H., (Ph.D. thesis) 2012. *Género Bauhinia* (Caesalpinoideae, Leguminosae) en *Astroamérica*: caracterización taxonómica, diferenciación morfológica y usos populares. Universidad Nacional de Buenos Aires, Buenos Aires, 238 p.
- Fortunato, R.H., Miotto, S., Izaguirre, P., Beyhaut, R., Bortoluzzi, R.L.C., Ulibarri, E., Gómez-Sosa, E., 2008. Fabaceae. In: Zuloaga, F.O., Morrone, O., Belgrano, M.J. (org.), *Catálogo de las Plantas Vasculares del Cono Sur*. Monog. Syst. Bot. 107, 2078–2319.
- Gadotti, V.M., Santos, A.R., Meyre-Silva, C., Schmeling, L.O., Machado, C., Liz, F.H., 2005. Antinociceptive action of the extract and the flavonoid quercitrin isolated from *Bauhinia microstachya* leaves. *J. Pharm. Pharmacol.* 57, 1345–1351.
- Hurrell, J.A., Ulibarri, E.A., Puentes, J.P., Buet Costantino, F., Arenas, P.M., Pochettino, M.L., 2011. *Leguminosas medicinales y alimenticias utilizadas en la conurbación Buenos Aires – La Plata, Argentina*. Bol. Latinoam. Caribe 10, 443–455.
- Ibarrola, D.A., Degen de Arrúa, R.L., 2011. *Catálogo Ilustrado de 80 Plantas Medicinales del Paraguay*. Facultad de Ciencias Químicas – UNA, Agencia de Cooperación International del Japón (JICA), Asunción.
- Lin, Y., Wong, W.O., Shi, G., Shen, S., Li, Z., 2015. Bilobate leaves of *Bauhinia* (Leguminosae, Caesalpinoideae, Cercideae) from the middle Miocene of Fujian Province, southeastern China and their biogeographic implications. *BMC Evol. Biol.* 15, <http://dx.doi.org/10.1186/s12862-015-0540-9>.
- Luna, J.D.S., Santos, A.F., Lima, M.R.F., Omena, M.C., Mendonça, F.A.C., Bieber, L.W., Sant'Ana, A.E.G., 2005. A study of the larvicidal and molluscicidal activities of some medicinal plants from northeast Brazil. *J. Ethnopharmacol.* 97, 199–206.
- Martins-Olivera, B.T., Almeida-Reis, R., Theodoro-Júnior, O.A., Oliva, L.V., Santos Nunes, N.N., Olivo, C.R., Brito, M.V., Prado, C.M., Leick, E.A., Arruda Martins, M., Oliva, M.L.V., Righetti, R.F., Tibério, I.F.L.C., 2016. The plant-derived *Bauhinia bauhinoides* kallikrein proteinase inhibitor (rBbKI) attenuates elastase-induced emphysema in mice. *Mediat. Inflamm.*, <http://dx.doi.org/10.1155/2016/534654>.
- Melo, J.G., Nascimento, V.T., Amorim, E.L.C., Andrade Lima, C.S., Albuquerque, U.P., 2004. Avaliação da qualidade de amostras comerciais de boldo (*Peumus boldus Molina*), pata-de-vaca (*Bauhinia spp.*) e ginkgo (*Ginkgo biloba* L.). *Rev. Bras. Farmacogn.* 14, 111–120.
- Oliveira, F., Saito, M.L., 1989. *Alguns vegetais brasileiros empregados no tratamento da diabetes*. *Rev. Bras. Farmacogn.* 2–4, 170–196.
- Nakahata, A.M., Bueno, N.R., Rocha, H.A., Franco, C.R., Chammás, R., Nakae, C.R., Jasulionis, M.G., Nader, H.B., Santana, L.A., Sampaio, M.U., Oliva, M.L.V., 2006. Structural and inhibitory properties of a plant proteinase inhibitor containing the RGD motif. *Int. J. Biol. Macromol.* 40, 22–29.
- Paula, C.D.S., Canteli, V.C.D., Hirota, B.C.K., Campos, R., Oliveira, V.B.D., Kalegari, M., Silva, C.B., Silva, G.M., Miguel, O.G., Miguel, M.D., 2014. Potencial antioxidante *in vitro* das folhas da *Bauhinia ungulata* L. *Rev. Cien. Farm. Bas.* 35, 217–222.
- Pizzoli, V.R., Brasileiro, B.G., Oliveira, T.T., Nagem, T.J., 2011. *Plantas com possível atividade hipolipidêmica: uma revisão bibliográfica de livros editados no Brasil entre 1998 e 2008*. *Rev. Bras. Pl. Med.* 13, 98–109.
- Rozza, A.L., Cesar, D.A.S., Pieroni, L.G., Salданha, L.L., Dokkedal, A.L., De-Faria, F.M., Souza-Brito, A.R.M., Vilegas, W., Takahira, R.K., Pellizzon, C.H., 2015. Antiulcerogenic activity and toxicity of *Bauhinia holophylla* hydroalcoholic extract. *Evid.-Based Complement. Alternat. Med.*, <http://dx.doi.org/10.1155/2015/439506>.
- Silva, K.L., Cechinel Filho, V., 2002. *Plantas do gênero Bauhinia: composição química e potencial farmacológico*. Quim. Nova 25, 449–454.
- Sousa, E., Zanatta, L., Seifriz, I., Creczynski-Pasa, T.B., Pizzolatti, M.G., Szpoganicz, B., Silva, F.R.M.B., 2004. Hypoglycemic effect and antioxidant potential of kaempferol-3,7-O-(α)-dirhamnoside from *Bauhinia forficata* leaves. *J. Nat. Prod.* 67, 829–832.
- Trethowan, L., Clark, R., Mackinder, B., 2015. A synopsis of the neotropical genus *Schnella* (Cercideae: Caesalpinoideae: Leguminosae) including 12 new combinations. *Phytotaxa* 204, 237–252.
- Vaz, A.M.S.F., 1979. Considerações sobre a taxonomia do gênero *Bauhinia* L. sect. *Tylotaea* Vogel (Leguminosae – Caesalpinoideae) do Brasil. Rodriguésia 31, 127–234.
- Vaz, A.M.S.F., Bortoluzzi, R.L.C., Silva, L.A.E., 2010. Checklist of *Bauhinia sensu stricto* (Caesalpiniaceae) in Brazil. *Plant Ecol. Evol.* 143, 212–221.
- Wunderlin, R.P., 2010a. Reorganization of the Cercideae (Fabaceae: Caesalpinoideae). *Phytoneuron* 48, 1–5.
- Wunderlin, R.P., 2010b. New combinations in *Schnella* (Fabaceae: Caesalpinoideae: Cercideae). *Phytoneuron* 49, 1–5.
- Zhang, D., 1994. Leaf venation of Cercideae. *J. Trop. Subtrop. Bot.* 2, 45–57.