

# All about neosporosis in Brazil

Tudo sobre neosporose no Brasil

Camila Koutsodontis Cerqueira-Cézar<sup>1</sup>; Rafael Calero-Bernal<sup>1</sup>; Jitender Prakash Dubey<sup>1</sup>; Solange Maria Gennari<sup>2\*</sup>

<sup>1</sup> Animal Parasitic Diseases Laboratory, Beltsville Agricultural Research Center, United States Department of Agriculture, Agricultural Research Service, Beltsville, MD, United States of America

<sup>2</sup> Departamento de Medicina Veterinária Preventiva e Saúde Animal, Faculdade de Medicina Veterinária e Zootecnia, Universidade de São Paulo – USP, São Paulo, SP, Brasil

Received January 22, 2017

Accepted June 28, 2017

## Abstract

*Neospora caninum* is protozoan parasite with domestic and wild dogs, coyotes and grey wolves as the definitive hosts and many warm-blooded animals as intermediate hosts. It was cultivated and named in 1988. Neosporosis is a major disease of cattle and has no public health significance. Since 1990's *N. caninum* has emerged as a major cause of abortion in cattle worldwide, including in Brazil. *N. caninum* also causes clinical infections in several other animal species. Considerable progress has been made in understanding the biology of *N. caninum* and there are more than 200 papers on this subject from Brazil. However, most of the reports on neosporosis from Brazil are serological surveys. Overall, little is known of clinical neosporosis in Brazil, particularly cattle. The few reports pertain to sporadic cases of abortion with no information on epidemics or storms of abortion. The objective of the present review is to summarize all reports from Brazil and suggest topic for further research, including prevalence of *N. caninum* oocysts in soil or in canine feces, and determining if there are additional definitive hosts, other than the domestic dog. There is need for a national survey in cattle using defined parameters. Future researches should focus on molecular characterization of *N. caninum* strains, possibility of vaccine production and relationship between wildlife and livestock epidemiology.

**Keywords:** *Neospora caninum*, neosporosis, domestic animals, wild animals, Brazil.

## Resumo

*Neospora caninum* é um protozoário parasita que possui os canídeos domésticos e selvagens, coiotes e lobos cinzentos como hospedeiros definitivos e vários animais de sangue quente como hospedeiros intermediários. Foi cultivado e nomeado em 1988. A neosporose é uma das principais doenças em bovinos e não tem significância em saúde pública. Desde 1990, *N. caninum* tem emergido como uma das principais causas de aborto em bovinos em todo o mundo, inclusive no Brasil. *N. caninum* também causa infecções clínicas em várias outras espécies animais. Consideráveis avanços foram feitos na compreensão da biologia desse parasita e há mais de 200 trabalhos sobre o assunto no Brasil. No entanto, a maioria dos relatos de neosporose do Brasil são relacionados a sorologia. Em geral, pouco se sabe sobre a neosporose clínica no Brasil, particularmente em bovinos. Os poucos relatos referem-se a casos esporádicos de aborto sem informações sobre epidemias ou surtos de aborto. O objetivo da presente revisão é resumir todos os relatos sobre *N. caninum* no Brasil e sugerir tópicos para pesquisas futuras, incluindo a prevalência de oocistos de *N. caninum* no solo ou em fezes caninas e determinar se há hospedeiros definitivos adicionais, exceto o cão doméstico no país. Uma pesquisa nacional em bovinos usando parâmetros definidos seria de grande importância. Pesquisas futuras deveriam ser focadas na caracterização de cepas de *N. caninum*, possibilidade de produção de vacinas e a relação epidemiológica entre a vida selvagem e o gado.

**Palavras-chave:** *Neospora caninum*, neosporose, animais domésticos, animais selvagens, Brasil.

\*Corresponding author: Solange Maria Gennari. Departamento de Medicina Veterinária Preventiva e Saúde Animal, Faculdade de Medicina Veterinária e Zootecnia, Universidade de São Paulo – USP, Av. Prof. Orlando Marques de Paiva, 87, Cidade Universitária, CEP 05508-270, São Paulo, SP, Brasil.  
e-mail: sgennari@usp.br

## Introduction

Neosporosis is relatively a newly recognized disease. In 1988, the etiologic agent of neosporosis was cultivated and named, *Neospora caninum*. It is ancestrally and morphologically related to *Toxoplasma gondii*. Since 1990's *N. caninum* has emerged as a major cause of abortion in cattle worldwide, including in Brazil. *N. caninum* also causes clinical infections in several other animal species. Considerable progress has been made in understanding the biology of *N. caninum*. A recent book on neosporosis (DUBEY et al., 2017) listed more than 2100 citations and most reports (>200) from Brazil. Many of the reports from Brazil are scattered in local journals. The objective of the present review is to summarize reports from Brazil and suggest topic for further research. To minimize citations, only references from Brazil are listed in bibliography.

## Basic Biology

As stated earlier, *N. caninum* and *T. gondii* are morphologically similar but biologically different coccidians. Both parasites have a wide host range but unlike *T. gondii*, *N. caninum* is not considered zoonotic. Canids (domestic and wild canids dogs, coyote, and wolf) are the definitive hosts of *N. caninum* whereas felids (domestic and wild Felidae) are definitive hosts for *T. gondii*. Neosporosis is a major disease of cattle whereas cattle are considered resistant to *T. gondii*. Transplacental transmission is a major route of propagation of *N. caninum* in cattle while, although medically important, is not the major route of transmission for *T. gondii*. There is only one species of *Toxoplasma*, *T. gondii*, but the genus *Neospora* has two species, *N. caninum* and *N. hughesi*; only horses are reported as an intermediate host for *N. hughesi*.

## History of Neosporosis in Brazil

The *in vitro* cultivation of *N. caninum* in 1988 in USA made it possible to develop diagnostic tests for neosporosis (DUBEY et al., 2017). However, unlike *T. gondii*, it is difficult to culture *N. caninum* (see later discussion) and only one isolate (NC-1) was available initially at the USDA laboratory in Beltsville, Maryland, USA. Thus, commercial tests were not developed for the diagnosis of neosporosis for several years. Therefore, no research was performed on this subject in Brazil until mid 1990's.

In BA, Gondim et al. (1999b, 2001) first recognized clinical neosporosis in an aborted bovine fetus and first isolated viable *N. caninum* (NC-Bahia) from the brain of a naturally-infected adult dog presenting incoordination and hind-limb paresis. Corbellini et al. (2002) first recognized neosporosis as an important cause of bovine abortion. They documented lesions consistent with protozoal infection in 22 (47.8%) of 46 fetuses submitted from 12 farms in RS. From those, 18 specimens of fetuses with encephalitis reacted to *N. caninum* antisera.

## Brazilian Contributions to the General Biology of *N. caninum*

### Viable *N. caninum* isolates and genetic diversity

Unlike *T. gondii*, little is known of genetic diversity of *N. caninum* mainly because there are only few viable isolates available worldwide. One reason for this is the difficulty to isolate viable *N. caninum* from animal tissues, especially from latently infected animals. Table 1 summarizes all viable isolates of *N. caninum* from animals in Brazil. The greatest success of isolating *Neospora* from asymptomatic animals has been achieved by feeding brain tissue from naturally-infected animals to dogs and then examining dog feces for oocyst excretion. By doing so, *N. caninum* was isolated from buffaloes, sheep and cattle.

In addition, to provide these *Neospora* isolates for biological and genetic studies to researchers in other countries (DUBEY et al., 2017), García-Melo in association with researchers from Spain did the first microsatellite typing data for Nc-Goiás isolated from clinically healthy cattle. Although the isolate had most of the alleles already identified, unique alleles were described for this strain at the MS5 and MS10 loci, using 12 microsatellite markers (GARCÍA-MELO et al., 2009). The recent isolate from brain tissue from a naturally-infected cattle (OLIVEIRA et al., 2017) was also found to have unique microsatellite alleles as MS5, MS6a, MS8 and MS10. A dog fed brain tissue from a naturally-infected cattle excreted *N. caninum* oocysts for 14 days starting seven days post infection (dpi), with an average number of 102 oocysts/g of feces. DNA isolated from cell culture derived tachyzoites was characterized using microsatellites. No new alleles were found and comparison showed closest relation to multilocus genotyping with strains from Spain and Argentina. Comparison with Brazilian strains (NC-Bahia; NC-Goiás) revealed variation in three and four of the nine markers used.

Additionally, the Brazilian NC-Bahia strain isolated from the dog was included among the *N. caninum* strains characterized by researchers outside Brazil (AL-QASSAB et al., 2009, 2010; REGIDOR-CERRILLO et al., 2013).

### Transmission

Tachyzoites (in groups), bradyzoites (in tissue cysts) and sporozoites (in oocysts) are the three infectious stages of *N. caninum*. Tachyzoites and bradyzoites occur in tissues of intermediate hosts whereas oocysts are excreted in feces of certain canids. Unlike *T. gondii*, relatively few oocysts are excreted by canids after ingesting infected tissues. However, oocysts were excreted with a high frequency by two-three-month old dogs fed brains from six naturally infected water buffaloes (RODRIGUES et al., 2004). An other study showed that oocyst excretion is more efficient by feeding brains than other tissues to dogs (CAVALCANTE et al., 2011). The authors fed 17 two-three month old dogs with different tissues from four cattle naturally infected with *N. caninum*. Each group of dogs received: masseter (n=5), heart (n=5), brain

**Table 1.** Isolation of *N. caninum* from naturally infected animals from Brazil.

Host	Source	Bioassay		Isolate designation	Genetic data	Remarks	Reference
		Animals	Cell culture				
Cattle	Clinical, blind, neurological calf, 3-months old	SW mice,	Vero	BCN/PR3	No	Tissue cysts were found in the brains of SW mice that were immunocompetent	Locatelli-Dittrich et al. (2003)
	Aborted fetus, 7 months of gestation	Mice, gerbils	Vero	BCN/PR1	Yes	Initial isolation in immunosuppressed mice. Tachyzoites infective to immunocompetent mice and gerbils	Locatelli-Dittrich et al. (2004)
	Asymptomatic, 4 months old	KO mice	MARC-145	Nc-Goiás 1b	Yes	Not pathogenic to BALB/c mice	García-Melo et al. (2009)
Dog	Adult slaughtered in abattoir	Dog, SW mice, gerbils	Vero	NC-SP1	Yes	Isolated from dog bioassay	Oliveira et al. (2017)
	Collie, 7 years old, clinical	Gerbil	Vero (COS-1)	NC-Bahia	Yes	Isolate pathogenic to gerbils	Gondim et al. (2001, 2004)
Sheep	Two 4 months old, clinically normal	Gerbil, SW and vesper mice, dogs	Vero	Not stated	No	Mice remained healthy. Necropsied 60 dpi. Tissue cysts in brains of both gerbils.	Pena et al. (2007)
Water buffalo	6 seropositive from abattoir	Dogs, gerbil, KO mice	CV1	NCBrBuf-1	No	Both gerbils and KO remained asymptomatic	Rodrigues et al. (2004)
				NCBrBuf-2			
				NCBrBuf-3			
				NCBrBuf-4			
				NCBrBuf-5			

KO=Gamma interferon gene knockout; SW=Swiss Webster.

(n=4), liver (n=3), and a group remaining as control (n=3), no infected. None of the control dogs excreted oocysts and three dogs that received brain, two that received masseter, two that received heart and one that received liver excreted *N. caninum*-like oocysts, from day seven to day 17 after ingestion of tissues. All dogs that received brain and liver excreted only *N. caninum* oocysts. The results were confirmed by polymerase chain reaction (PCR) using Nc5 and ITS-1 amplification and indicated that a variety of visceral, neural, and muscular tissues are infected naturally with *N. caninum*.

How dogs become infected with *N. caninum* in nature is not fully understood. Fecal transmission of *N. caninum* in dogs appears to be less important than carnivorism. Until the study by Bandini et al. (2011) it was unknown if the dogs could be infected via ingestion of oocysts. They fed four dogs with 1,000, 5,000 or 10,000 *N. caninum* oocysts; none of the four dogs excreted *N. caninum*-like oocysts in their feces during the observation period of 30 days. However, the two dogs fed with 10,000 oocysts seroconverted and the two dogs fed with 1,000 or 5,000 oocysts did not. Neither parasite DNA nor parasite stages were demonstrable in tissues of the seropositive dogs euthanized six months after feeding oocysts. These findings suggest that fecal transmission may not be an important mode of transmission of the parasite for the definitive host but results need confirmation.

Tachyzoites are important in transplacental transmission of *N. caninum* infection. Cavalcante et al. (2012) confirmed transplacental transmission of *N. caninum* in dogs. Six bitches in two groups were inoculated with a very high dose ( $10^8$  tachyzoites). In group I, three bitches were inoculated during the third week of

gestation, and in group II, three bitches were inoculated at the sixth week of gestation. The bitches were allowed to whelp naturally. Dams and their pups were tested by immunohistochemistry (IHC), serology, and PCR. In group I, six of the ten pups died within 48 hour of birth. In group II, seven of the 13 pups died between five and ten days of birth. *N. caninum* DNA was detected by nested PCR in two pups (hearts of both and liver of one) from group I, and one pup in central nervous system (CNS) and lymph node from group II. The dams and the pups that survived were clinically normal. *N. caninum* was not demonstrable in tissues of any of pups and their dams.

Studies in Brazil showed that *N. caninum* can be transmitted transplacentally in water buffaloes (CHRYSSAFIDIS et al., 2014, 2015). The authors conducted an important experiment in six buffaloes and seven cows inoculated intravenously with *N. caninum* tachyzoites at 70 day of gestation. Three buffaloes and five cows were inoculated with a Brazilian *N. caninum* strain (NC-Bahia); only one cow aborted but all fetuses became infected. The other two cows and three buffaloes were inoculated with the NC-1 strain; all fetuses died by 35 dpi as determined by ultrasound and *N. caninum* DNA was detected in fetal tissues.

Chryssafidis et al. (2011) also showed, for the first time, that *N. caninum* can be transmitted transplacentally in naturally infected buffaloes; they found *N. caninum* DNA in one brain of the nine fetuses from buffaloes slaughtered in an abattoir, aiming Nc5 and ITS-1 DNA regions. Although, viable or intact parasite has not been demonstrated in naturally infected fetuses from buffaloes, this is the first indication of transplacental transmission in this host.

In addition to bradyzoites and oocysts, tachyzoites can also be infectious orally. Four to five day old gerbils were successfully infected by oral inoculation of tachyzoites (FERREIRA et al., 2016). All 17 gerbils died of neosporosis between eight and 17 dpi (one died on each of days 8, 9, 15, 16, and 17, four died on day 12, and five died on day 15 (personal communication to JPD on October 24, 2016) with  $4 \times 10^5$  NC-1 tachyzoites. *N. caninum* DNA was found in heart, lung, spleen, kidney, liver and CNS of gerbils.

### *Environmental resistance of oocysts*

Oocyst is the environmentally resistant stage. Researchers in Brazil found that *N. caninum* oocysts were rendered noninfectious by heating to 100 °C for one minute, and 10% sodium hypochlorite for one hour but not at 60 °C for one minute, and by other commonly used disinfectants (ALVES et al., 2011). Aqueous 2% sulfuric acid has been commonly used to store *N. caninum* oocysts at 4 °C; how long oocysts remain viable under these conditions is not known. *N. caninum* oocysts remained viable for 108 days but not for 46 months when stored in 2% sulfuric acid at 4 °C (UZÊDA et al., 2007a).

### *Diagnostic tests*

Recently, Fehlberg et al. (2017) reported on successful development of a high resolution melting PCR method to distinguish *Neospora*, *Sarcocystis* and *Toxoplasma* using a single pair of primers targeting 18S rDNA.

### *Epidemiology*

More serological studies for *N. caninum* infection in animals have been conducted in Brazil than rest of the world.

The results, however, are not comparable because of different serological assays used different cut-offs employed, different antigens used. For example, in the indirect fluorescent antibody test (IFAT) and the *Neospora* agglutination test (NAT) whole tachyzoites are used detecting antibodies to surface proteins whereas in the enzyme-linked immunoabsorbent assay (ELISA) different antigens are used, some of them using crude tachyzoite extract while others used soluble antigens (Table 2a). The standardization of serological tests was based on studies in other countries and a full discussion of these is beyond the scope of this review—this subject was recently reviewed (DUBEY et al., 2017). One of the problems with serological testing is the availability of standardized sera from experimentally infected animals. Immunoblotting (IB) was employed in some studies where no such sera were available (Table 2b).

Little has been done in Brazil to characterize *Neospora* recombinant antigens for the serological diagnosis or vaccine development, except the report of Bezerra et al. (2017) who characterized one surface protein, by cloning the sequence and named it as NCsRS67, which has no orthologue with *Toxoplasma gondii*, only with *Hammondia hammondi*.

In Tables 3-12 serological reports in different hosts are summarized. We listed all reports that we found.

Salient features are commented for some of these surveys.

**Dogs:** Numerous surveys from different regions of Brazil are summarized in Table 3. As stated earlier results of these types of surveys are not strictly comparable. However, 45 of 49 surveys used IFAT as a diagnostic technique, and most of them employed the cut-off value of 1:50, facilitating comparisons of occurrence values. In three surveys, IFAT and ELISA were compared for serological diagnosis of *N. caninum* in dog sera and IFAT was superior to indirect ELISA used (SILVA et al., 2007; HIGA et al., 2000; RAIMUNDO et al., 2015).

In some studies, risk factors were evaluated (Table 3). The age of dogs was a statistically significant factor in nine reports; older dogs were more likely to be seropositive (OLIVEIRA et al., 2004; FERNANDES et al., 2004; AZEVEDO et al., 2005; ANDREOTTI et al., 2006; CUNHA et al., 2008; MINERVINO et al., 2012; NOGUEIRA et al., 2013; BALTHAZAR et al., 2013; RAIMUNDO et al., 2015). Gender and breed were not associated with presence of antibodies. In two surveys mixed breed dogs constituted a risk factor for the infection (BRUHN et al., 2012; RAIMUNDO et al., 2015). The diet in some studies was related with access to street and prevalence was higher in dogs that had outdoor access than in pets with little or no outdoor scavenging (GENNARI et al., 2002; CAÑÓN-FRANCO et al., 2003; BENETTI et al., 2008; SICUPIRA et al., 2012). However, in some studies such an association was not found (MINEO et al., 2004; JESUS et al., 2006; PLUGGE et al., 2011). Vertical transmission was also studied (TAQUES et al., 2016), with 41 stillborn puppies from 23 bitches. By PCR and IFAT, five (21.7%) bitches were positive and 22 (53.6%) stillborn were positive by PCR, utilizing ITS-1 DNA region/locus, being 17 from positive bitches and five from negative ones. Although the prevalence of positive stillborn was higher from positive bitches, no conclusions were made.

Epidemiologically, contact between cattle and dogs has been identified as a possible risk factor that deserves attention (Table 3). Dogs from peri-urban or rural areas had more chance to be infected by *N. caninum* (IFAT 1:50) than urban dogs in the surveys from MG (FERNANDES et al., 2004), PR (PLUGGE et al., 2008), RS (CUNHA et al., 2008), and BA (SICUPIRA et al., 2012), and this risk factor is normally associated with proximity to cattle; access and ingestion of fetal membranes, carcasses, and prey.

*Leishmania infantum chagasi* is an important parasite of dogs in Brazil; and immunosuppression caused by *Leishmania* spp. may enhance the susceptibility of dogs to *N. caninum* infection (CRINGOLI et al., 2002). Serological surveys correlating *N. caninum* and *Leishmania* spp., had been conducted (GENNARI et al., 2006; ANDREOTTI et al., 2006; GUIMARÃES et al., 2009; GRECA et al., 2010; VALADAS et al., 2010a; LOPES et al., 2011; PAULAN et al., 2013; SEABRA et al., 2015; CONSTANTINO et al., 2016). In dogs from endemic cities of visceral leishmaniasis such as Araçatuba, SP (GENNARI et al., 2006), Campo Grande, MS (ANDREOTTI et al., 2006), and Teresina, PI (LOPES et al., 2011), positive association was found (Table 3).

**Cattle:** Seroprevalence of *N. caninum* varied with the type of cattle (beef, dairy), different regions, within region and with the type of serological tests used (Table 4).

**Table 2a.** ELISA techniques used to confirm presence of *N. caninum* antibodies.

Sera source	Antigen	ELISA type*	Results in table	Reference
Dogs	Antigen incorporated into iscoms	A, C, I		Mineo et al. (2004)
	NC-1 strain maintained in bovine monocytes cells; soluble antigen	B, C	3	Silva et al. (2007)
	NC-1 tachyzoites strain in Vero cells	A, C		Raimundo et al. (2015)
	NC-1 tachyzoites strain in Vero cells, lysed in SDS buffer	J		Chahan et al. (2003)
Cattle	Commercial	E		Andreotti et al. (2004)
	Commercial	F		Paz et al. (2007)
	Commercial	E		Melo et al. (2001)
	Commercial	E		Melo et al. (2004)
	Antigen incorporated into iscoms	I		Mineo et al. (2006)
	Commercial	E	4	Locatelli-Dittrich et al. (2001)
	Commercial	E		Locatelli-Dittrich et al. (2008)
	Commercial	E		Marques et al. (2011)
	Commercial	E		Nascimento et al. (2014)
	NC-1 strain in Vero cells; soluble antigen	B, C		Ramos et al. (2016)
Buffalo	Commercial	E		Munhoz et al. (2006, 2009)
	Commercial	D		Vogel et al. (2006)
	Commercial	E		Sartor et al. (2003)
	Commercial	M		Sartor et al. (2005)
	Commercial	E		Viana et al. (2009)
Sheep	Commercial	D	5	Vogel et al. (2006)
	Commercial kit	G, H		Silva et al. (2014)
Captive deer	NC-1 strain in Vero cells	A, L		Andreotti et al. (2009)
	NC-1 strain in Vero cells, soluble antigen	A, C	6	Rossi et al. (2011)
	Commercial	D		Vogel et al. (2006)
Horse	Commercial	E	11	Zimpel et al. (2015)
	NC-1 strain in HeLa cells	A, C		Pivoto et al. (2014)
	NC-1 strain in bovine turbinate cell monolayers	C, K	12	Hoane et al. (2006)

\*A = WT = Whole tachyzoite extract; B = SA = soluble antigen; C = IH = In house; D = CHEKIT = CHEKIT *Neospora*, indirect ELISA, detergent lysate of tachyzoites, IDEXX Laboratories, The Netherlands; E = IDEXX = IDEXX HerdChek *Neospora caninum* antibody, indirect ELISA, sonicate lysate of tachyzoites, IDEXX Laboratories, USA; F = VMRD = *Neospora caninum* cELISA competitive ELISA GP65 surface antigen of tachyzoites VMRD, USA; G = Horse IgG = Horse IgG(T) ELISA Quantitation Set; Bethyl Laboratories; H = Horse IgM = Horse IgM ELISA, Kamiya Biomedical Company, Seattle; I = ISCOM = Detergent extracted tachyzoite antigen incorporated in immune stimulating complex particles; J = NcSAG1 = Recombinant surface antigen; K = NhSAG1 = Recombinant NhSAG1; L = rNcSRS2 = Recombinant antigen protein; M = NS/ND = not stated/ not done.

**Table 2b.** Immunoblotting technique used to confirm presence of *N. caninum* antibodies.

Sera source	Antigen*	Sera tested previously	Results in Table	Reference
Dog	NC-1 tachyzoites lysed in SDDS-PAGE buffer	IFAT, 1:25; indirect ELISA	3	Silva et al. (2007)
Cattle	NC-1 tachyzoites lysed in SDDS-PAGE buffer	NcSAG1 ELISA, 60/66 sera positive for ELISA	4	Chahan et al. (2003)
Sheep	NC-1 tachyzoites lysed in 4% sodium dodecyl sulfate buffer	IFAT, 1:50; discordant results	6	Rossi et al. (2011)
Cat	Detergent extracted whole NC-1 tachyzoites	NAT, 1:80	9	Dubey et al. (2002)
Donkey	Whole NC-1 tachyzoites	Sera positive by IFAT, 1:100	12	Galvão et al. (2015)
Human	NC-1 tachyzoites lysed in SDDS-PAGE buffer	Sera with discordant results by IFAT and ELISA	-	Lobato et al. (2006)

\*Produced in non reducing conditions.

**Table 3.** Serological studies of *N. caninum* in dogs from Brazil.

State	Type	No. tested	No. positive	% Positive	Test	Cut-off	Remarks	Reference
Alagoas	Urban	128	5	3.8	IFAT	1:50	Age, sex, breed, area, habitat	Sousa et al. (2012)
Alagoas	Rural	99	5	4.8	IFAT	1:50	<b>Age, sex, <i>T. gondii</i>, others</b>	Minervino et al. (2012)
<b>Amazon region</b> (MT and TO)	Indian communities	325	32	9.8	IFAT	1:50	<b>Age, sex, <i>T. gondii</i>, others</b>	Jesus et al. (2006)
Bahia	Stray	250	28	11.2	IFAT	1:50	Age, sex, others	Sicupira et al. (2012)
Bahia	Owned	165	22	13.3	IFAT	1:50	<b>Area, contact with other dogs, feeding habits, others</b>	Acosta et al. (2016)
Bahia	Urban	156	4	2.6	IFAT	1:50	<b>Area, contact with other dogs, feeding habits, others</b>	Boaventura et al. (2008)
Bahia	Rural	41	6	14.6	IFAT	1:50	<b>Area, contact with other dogs, feeding habits, others</b>	Téixeira et al. (2006)
Bahia	Peri-urban	214	28	13.1	IFAT	1:50	<b>Area, contact with other dogs, feeding habits, others</b>	Benefit et al. (2008)
Esírito Santo	Rural	187	22	11.7	IFAT	1:50	<b>Sex, <i>T. gondii</i></b>	Benetti et al. (2009)
Goiás	Zoonosis Center	72	26	36.1	IFAT	1:50	Sex, origin	Oliveira et al. (2004)
Maranhão	Hospital (owned)	125	39	31.2	IFAT	1:50	Sex, origin	Andreotti et al. (2004)
Matto Grosso	Stray	100	45	45.0	IFAT	1:50	Age, sex, diet, access to streets	Mineo et al. (2001)
Matto Grosso	Clinics	60	27	45.0	IFAT	1:50	Age, sex, diet, access to streets	Fernandes et al. (2004)
Matto Grosso do Sul	Rural	37	25	67.6	IFAT	1:50	Age, sex	Téixeira et al. (2006)
Matto Grosso do Sul	Pet	245	65	26.5	IFAT	1:50	<b>Age, sex</b>	Benefit et al. (2008)
Matto Grosso do Sul	Rural	40	12	30.0	IFAT	1:100	<b>Age, sex, <i>T. gondii</i></b>	Benetti et al. (2009)
Matto Grosso do Sul	Urban	345	93	27.2	IFAT	1:50	<b>Age, sex, <i>Leishmania</i></b>	Oliveira et al. (2004)
Minas Gerais	Clinical	163	11	6.7	IFAT	1:25	<b>Age, sex, <i>T. gondii</i></b>	Andreotti et al. (2004)
Minas Gerais	Urban	300	32	10.7	IFAT	1:50	<b>Age, breed, sex, area</b>	Andreotti et al. (2006)
Minas Gerais	Peri urban	58	11	18.9	IFAT	1:50	<b>Age, breed, sex, area</b>	Mineo et al. (2001)
Minas Gerais	Rural	92	20	21.7	IFAT	1:50	<b>Age, breed, sex, habitat, <i>T. gondii</i>, others</b>	Fernandes et al. (2004)
Minas Gerais	Clinic	275	22	7.9	ELISA	A,C	<b>Age, breed, sex, habitat, <i>T. gondii</i>, others</b>	Mineo et al. (2004)
Minas Gerais	Stray	94	12	12.8	ELISA	I	<b>Age, breed, sex, <i>T. gondii</i></b>	Mineo et al. (2004)
Minas Gerais	Clinic, stray	300	32	10.7	IFAT	1:50	<b>Age, breed, sex, <i>T. gondii</i></b>	Silva et al. (2007)
Minas Gerais	Clinic, stray	300	105	35.0	ELISA	B,C	<b>Age, breed, sex <i>T. gondii</i>, <i>Leishmania</i>, <i>Babesia canis</i></b>	Guimarães et al. (2009)
Minas Gerais	Clinics	228	7	3.1	IFAT	1:50	<b>Age, breed, origin, sex, others</b>	Bruhn et al. (2012)
Minas Gerais	Rural	240	36	15.0	IFAT	1:50	<b>Age, diet, hunting, area</b>	Nogueira et al. (2013)
Minas Gerais	Urban	182	15	8.2	IFAT	1:50	<b>Bovine abortion, area</b>	Valadas et al. (2010b)
Minas Gerais	Rural	421	58	13.7	IFAT	1:50	Sex, area <i>T. gondii</i> , <i>Leishmania</i>	Azevedo et al. (2005)
Pará	Rural	72	8	11.1	IFAT	1:50	<b>Age, sex, breed, others, habitat, <i>T. gondii</i></b>	Souza et al. (2002)
Pará	Urban-stray	57	8	14.0	IFAT	1:50	<b>Age, sex, breed</b>	Giraldi et al. (2002)
Parába	Urban Domestic	286	24	8.4	IFAT	1:50	<b>Age, sex, breed, others, habitat, <i>T. gondii</i></b>	Romanelli et al. (2007)
Paraná	Dairy farms	134	29	21.6	IFAT	1:50	<b>Age, sex, breed</b>	Plugge et al. (2008)
Paraná	Neurological	98	0	0.0	IFAT	1:50	<b><i>T. gondii</i></b>	Locatelli-Dittrich et al. (2008)
Paraná	Sheep farms	24	7	29.1	IFAT	1:50	<b>Age, sex, breed, <i>T. gondii</i></b>	
Paraná	Urban	181	23	12.7	IFAT	1:50	<b>Area, habitat</b>	
Paraná	Peri urban	178	28	15.7	IFAT	1:50		
Paraná	Rural	197	50	25.3	IFAT	1:50		
Paraná	Rural	129	32	25.0	IFAT	1:50	Dogs and cattle	

Bold=statistically significant risk factor; Area = urban, peri urban, rural; Habitat=stray, domiciled.

Table 3. Continued...

State	Type	No. tested	No. positive	% Positive	Test	Cut-off	Remarks	Reference
<b>Paraná</b>	Owned	127	14	11.0	IFAT	1:50	<i>T. gondii</i> , neurologic signs	Plugge et al. (2011)
	Stray	20	3	15.0	IFAT	1:50		
<b>Paraná</b>	Stray	26	3	11.5	IFAT	1:25	<i>Leishmania</i> spp., <i>T. gondii</i> , <i>Trypanosoma cruzi</i>	Constantino et al. (2016)
<b>Pernambuco</b>	Domiciled	289	75	26.0	IFAT	1:50	<i>T. gondii</i> , origin	Figueiredo et al. (2008)
Paulista	Domiciled	168	44	26.2	IFAT	1:50		
Amaraji	Domiciled	168	58	34.5	IFAT	1:50		
Garanhuns	Rural villages	56	0	0	IFAT	1:50	<i>T. gondii</i> antibodies surveyed	Araujo-Santos et al. (2016)
<b>Pernambuco</b>	Rural villages	530	17	3.2	IFAT	1:50	Age, sex, breed, <i>T. gondii</i> , <i>Leishmania</i>	Lopes et al. (2011)
<b>Piauí</b>	Rural villages	71	5	7.0	IFAT	1:50	<i>T. gondii</i> antibodies surveyed	Araujo-Santos et al. (2016)
<b>Piauí</b>	Urban - Clinic	402	34	8.4	IFAT	1:50	Age, neurological signs, others	Balthazar et al. (2013)
<b>Rio de Janeiro</b>	Rural	230	47	20.4	IFAT	1:50	Age, sex, area, type of farm, carcasses and abortions not removed, others	Cunha et al. (2008)
<b>Rio Grande do Sul</b>	Urban	109	6	5.5	IFAT	1:50		
	Domiciled (Street access)	157	13	8.3	IFAT	1:50		
<b>Rondônia</b>	Farms	174	22	12.6	IFAT	1:50	Age, diet, abortion, stillbirth, others	Cañón-Franco et al. (2003)
<b>São Paulo</b>	Hospital	203	44	21.6	IFAT	1:50	<i>T. gondii</i> , neurological signs	Aguiar et al. (2006)
<b>São Paulo</b>	Rural and urban	295	25	8.4	IFAT	1:50	Age, sex, <i>T. gondii</i> , neurological signs	Higa et al. (2000)
	Domiciled	500	49	10.0	NAT	1:25	Age, sex, breed, habitat	Varandas et al. (2001)
<b>São Paulo</b>	Street	611	151	25.0	NAT	1:25		Gennari et al. (2002)
<b>São Paulo</b>	Urban	204	36	17.6	IFAT	1:50	Age, sex, <i>T. gondii</i> , <i>Leishmania</i> spp.	Gennari et al. (2006)
<b>São Paulo</b>	Urban	108	17	15.7	IFAT	1:50	Age, sex, diet, others, <i>T. gondii</i>	Bresciani et al. (2007b)
<b>São Paulo</b>	Urban	865	223	25.8	IFAT	1:50	Age, sex, area	Moraes et al. (2008)
	Rural	65	11	16.9	IFAT	1:50		
	Peri urban	33	11	33.3	IFAT	1:50		
<b>São Paulo</b>	Rural (stray)	100	14	14.0	IFAT	1:25	<i>Leishmania</i> spp.	Greca et al. (2010)
<b>São Paulo</b>	Sheep farms	42	2	4.8	IFAT	1:25	Type of raising (chained or free), access to raw meat or offal	Machado et al. (2011)
<b>São Paulo</b>	Neurologic	50	7	14%	IFAT	1:25	<i>T. gondii</i>	Langoni et al. (2012)
<b>São Paulo</b>	Domiciled	342	17	4.9	IFAT	1:25	Age, sex, breed, <i>T. gondii</i>	Langoni et al. (2014)
<b>São Paulo</b>	Rural	93	6	6.5	IFAT	1:50	<i>Leishmania</i> , <i>T. gondii</i> , <i>Ehrlichia</i> spp., <i>Babesia canis</i>	Paulan et al. (2013)
<b>São Paulo</b>	Kennel	167	37	22.1	IFAT	1:25	<i>T. gondii</i> , <i>Leishmania</i>	
	Clinic	133	36	27.0	IFAT	1:25	<i>T. gondii</i> , <i>Leishmania</i>	Seabra et al. (2015)
<b>Tocantins</b>	Rural	99	43	43.4	ELISA	A,C	<i>Age</i> , <i>breed</i> , area, <i>T. gondii</i>	Raimundo et al. (2015)
	Urban	105	45	31.3	IFAT	1:25		
			31	42.9	ELISA	A,C		
			31	29.5	IFAT	1:25		

**Bold=**statistically significant risk factor, Area = urban, peri urban, rural; Habitat=stray, domiciled.

Table 4. Serologic studies of *N. caninum* antibodies in cattle from Brazil.

State	No. tested	Type	No. herds	No. positive	% positive	Test	Cut-off	Remarks <sup>a</sup>	Reference
Bahia	447	Dairy	14	63	14.0	IFAT	1:200	-	Gondim et al. (1999a)
Goiás	444	Dairy	11	135	30.4	IFAT	1:250	-	Melo et al. (2006)
Goiás	30	Mixed	1	13	43.3	IFAT	1:250	-	Melo et al. (2006)
Goiás	456	Beef	9	135	29.6	IFAT	1:250	-	Melo et al. (2006)
Maranhão	812	Dairy	27	412	50.7	IFAT	1:200	-	Teixeira et al. (2010)
Matto Grosso	932	Dairy	24 farms	499	53.5	IFAT	1:200	-	Benetti et al. (2009)
Matto Grosso do Sul	197	NS	6	66	33.5	ELISA	J	-	Chahan et al. (2003)
Matto Grosso do Sul	87	Beef	NS	26	29.9	IFAT	1:25	-	Ragozo et al. (2003)
Matto Grosso do Sul	23	Dairy	NS	5	21.7	IFAT	1:25	-	Ragozo et al. (2003)
Matto Grosso do Sul	90	Beef (history of abortions)	NS	38	43.0	ELISA	E	-	Andreotti et al. (2004)
Matto Grosso do Sul	60	Heifers	18	30.0	ELISA	E	-	Andreotti et al. (2004)	
Matto Grosso do Sul	91	Healthy	7	7.7	ELISA	E	-	Andreotti et al. (2004)	
Matto Grosso do Sul	2448	NS	205	449	14.9	IFAT	1:50	Production system (Dairy/beef)	Oshiro et al. (2007)
Matto Grosso do Sul	275	Beef	2 farms	81	29.5	eELISA	F	-	Paz et al. (2007)
Matto Grosso do Sul	392	Dairy/beef	4 farms	43	9.1	IFAT	1:50	-	Mello et al. (2008)
Matto Grosso do Sul	1098	Beef	1 farm	687	62.5	IFAT	1:50	Reproductive failure, 15% higher in seropositive cows.	Andreotti et al. (2010)
Minas Gerais	584	Dairy	18	109	18.7	ELISA	E	-	Melo et al. (2001)
Minas Gerais	126	Dairy	2	43	34.4	IFAT	1:25	-	Ragozo et al. (2003)
Minas Gerais	36	Beef	NS	4	11.1	IFAT	1:25	-	Ragozo et al. (2003)
Minas Gerais	576	Dairy	18	106	18.4	ELISA	E	-	Melo et al. (2004)
Minas Gerais	243	Dairy	2	41	16.8	ELISA	I	-	Mineo et al. (2006)
Minas Gerais	559	Dairy	18	510	91.2	IFAT	1:200	Farm size, number of cows lactating, milk production per day	Guedes et al. (2008)
Minas Gerais	575	Dairy	Abattoir	559	97.2	IFAT	1:200	Farm size, number of cows lactating, milk production per day	Guedes et al. (2008)
Minas Gerais	503	Dairy	Abattoir (fetus)	64	12.7	IFAT	1:25	Farm size, number of cows lactating, milk production per day	Guedes et al. (2008)
Minas Gerais	1,204	Dairy	40 farms	260	21.6	IFAT	1:200	Reproductive failure	Bruhn et al. (2013)
Pará	500	beef	500	52	IFAT	1:128	Region, <i>T. gondii</i>	Silva et al. (2017)	
Pará	40	Dairy	4 farms	7	17.5	IFAT	1:100	-	Minervino et al. (2008)
Pará	120	Beef	12	23	19.2	IFAT	1:100	-	Minervino et al. (2008)
Paraná	172	Dairy	1	60	34.8	ELISA	E	Abortion	Locatelli-Dittrich et al. (2001)
Paraná	15	Beef	NS	4	26.7	IFAT	1:25	-	Ragozo et al. (2003)
Paraná	75	Dairy	NS	16	21.3	IFAT	1:25	-	Ragozo et al. (2003)
Paraná	623	Dairy	23 farms	89	14.3	IFAT	1:25	Breed, presence of dogs, age, feed	Guimarães et al. (2004)
Paraná	385	Dairy	90 farms	45	12.0	IFAT	1:200	-	Ogawa et al. (2005)

<sup>a</sup>Statistically significant risk factors.

Table 4. Continued...

State	No. tested	Type	No. herds	No. positive	% positive	Test	Cut-off	Remarks <sup>a</sup>	Reference
Paraná	1263	NS	77 farms	423	33.0	ELISA	E	-	Locatelli-Dittrich et al. (2008)
Paraná	159	Beef	Abattoir	24	15.1	ELISA	E	-	Marques et al. (2011)
Paraná	309	Dairy	15 farms	63	20.4	IFAT	1:100	Feed, wild animal access, artificial insemination	Martins et al. (2012)
Paraná	76	Beef	4	23	30.3	ELISA	E	-	Nascimento et al. (2014)
Pernambuco	469	Dairy	20 farms	163	31.7	IFAT	1:200	Veterinary assistance, nutritional condition, presence of wetlands, manipulation of newborn calves, destination of cows that had aborted, abortion history, abortions period.	Silva et al. (2008)
Pernambuco	316	Dairy	25 municipalities	31	19.6	IFAT	1:200	Transplacental transmission	Ramos et al. (2016)
Rio Grande do Sul	223 (abortion history)	Dairy	5	25	11.2	IFAT	1:200	Reproductive failure	Corbellini et al. (2002)
Rio de Janeiro	75	Dairy	NS	17	22.7	IFAT	1:25	-	Ragozo et al. (2003)
Rio de Janeiro	75	Beef	NS	5	6.7	IFAT	1:25	-	Ragozo et al. (2003)
Rio de Janeiro	563	Dairy	57 farms	131	23.2	ELISA	E	Breed	Munhoz et al. (2006, 2009)
Rio Grande do Sul	70	Dairy	NS	13	18.6	IFAT	1:25	-	Ragozo et al. (2003)
Rio Grande do Sul	70	Beef	NS	15	21.4	IFAT	1:25	-	Ragozo et al. (2003)
Rio Grande do Sul	1,549	Dairy	60 farms	276	17.8	IFAT	1:200	several	Corbellini et al. (2006)
Rio Grande do Sul	781	Dairy/beef	NS	89	11.4	ELISA	D	-	Vogel et al. (2006)
Rondônia	1011	Dairy	50	114	11.2	IFAT	1:25	Farm size, number of cows	Aguiar et al. (2006)
Rondônia	584	Beef	11 farms	56	9.5	IFAT	1:25	-	Aguiar et al. (2006)
Rondônia	514	Mixed	25 farms	50	9.7	IFAT	1:25	-	Aguiar et al. (2006)
Rondônia	621	Dairy	63 farms	66	10.6	IFAT	1:100	Abortion, birth of weak calves	Boas et al. (2015)
Santa Catarina	1518	Dairy	72 farms	466	30.6	IFAT	1:100	Presence of dogs	Fávero et al. (2017)
Santa Catarina	130	Dairy	29 farms	57	43.8	IFAT	1:200	Age, no. of pregnancies	Klauck et al. (2016)
São Paulo and Minas Gerais	600	NS	NS	101	16.8	IFAT	1:200	Area	Costa et al. (2001)
São Paulo	150	NS	NS	41	27.3	IFAT	1:25	-	Ragozo et al. (2003)
São Paulo	521	Dairy	NS	82	15.9	IFAT	1:200	-	Sartor et al. (2003)
São Paulo	521	Dairy	NS	159	30.5	ELISA	E	-	Sartor et al. (2003)
São Paulo	777	Beef	8 Farms	121	15.5	IFAT	1:200	-	Hasegawa et al. (2004)
São Paulo	505	Beef	11 Herds	101	20.0	ELISA	M	Production system	Sartor et al. (2005)
São Paulo	408	Dairy	6 herds	145	35.5	ELISA	M	<i>N. caninum</i> significantly higher in dairy cattle	Sartor et al. (2005)
São Paulo	1027	Dairy	3 farms	107	10.4	IFAT	1:100	High degree of association between <i>N. caninum</i> serological status of dams and daughter	Cardoso et al. (2012a)
Tocantins	192	Dairy	10 farms	48	25.0	IFAT	1:200	-	Martins et al. (2011)

<sup>a</sup>Statistically significant risk factors.

**Table 5.** Serological studies of *N. caninum* in buffaloes from Brazil.

State	No. of farms	No. tested	No. positive	% positive	Test	Cut-off	References
Bahia	4	117	42	35.9	IFAT	1:200	Gondim et al. (2007)
Pará	3	196	139	70.9	IFAT	1:25	Gennari et al. (2005)
Pará	26	500	195	39	IFAT	1:128	Silva et al. (2017)
Pará	4	212	187	88.2	ELISA	E	Viana et al. (2009)
Paraíba	14	136	26	19.1	IFAT	1:200	Brasil et al. (2015)
Rio Grande do Sul		164	24	14.6	ELISA	D	Vogel et al. (2006)
São Paulo	11	222	117	53.0	NAT	1:40	Fujii et al. (2001a, b)
São Paulo			142	64.0	IFAT	1:25	
São Paulo	12	411	230	56.0	IFAT	1:200	Souza et al. (2001)
São Paulo	5	192	169	88.0	IFAT	1:50	Chryssafidis et al. (2015)
<b>Northern Brazil-13 provinces</b>		4,796	2,665	55.5	ELISA	G, H	Silva et al. (2014)
			2,345	48.8	IFAT	1:40	

**Table 6.** Serological studies of *N. caninum* in sheep from Brazil.

State	Type	No. tested	No. positive	% positive	Test	Cut-off	Remarks	Reference
Alagoas	26 farms	343	33	9.6	IFAT	1:50	<b>Small farms, water supply</b>	Faria et al. (2010)
Federal District	32 farms	1028	90	8.8	IFAT	1:50	Titers up to 51,200	Ueno et al. (2009)
Maranhão	5 farms	64	3	4.7	IFAT	1:25	<b>Food supplement, reproductive failure</b>	Moraes et al. (2011)
Mato Grosso do Sul	1 farm	441	136	30.8	IFAT	1:50		Andreotti et al. (2009)
			141	32.0	ELISA	A, L	Comparison of techniques	
Minas Gerais	12 farms	334	27	8.1	IFAT	1:50	<b>Abortion history</b>	Salaberry et al. (2010)
Minas Gerais	2 farms	155	73	47.1	IFAT	1:50		Rossi et al. (2011)
			41	26.4	ELISA	A, C	Age	
Minas Gerais	63 farms	488	64	13.1	IFAT	1:50	Age, area	Andrade et al. (2012)
Paraná	9 farms	305	29	9.5	IFAT	1:50	Sex, age, breed, reproductive fails, presence of dogs	Romanelli et al. (2007)
Pernambuco	23 farms	81	52	64.2	IFAT	1:50	<b>Age</b>	Tembue et al. (2011)
Pernambuco	Rural villages	179	39	21.8	IFAT	1:50	<b>Region, age, sex, breed</b>	Arraes-Santos et al. (2016)
Piauí	Rural villages	153	8	5.2	IFAT	1:50	<b>Region, age, sex, breed</b>	Arraes-Santos et al. (2016)
Rio Grande do Norte	35 farms	409	7	1.8	IFAT	1:50	Sex, presence of dogs, reproductive fails	Soares et al. (2009a)
Rio Grande do Sul	4 counties	62	2	3.2	ELISA	D		Vogel et al. (2006)
Rondônia	15 farms	141	41	29.0	IFAT	1:50	Titers up to 1:25,600; Reproductive problems, presence of dogs, source of water	Aguiar et al. (2004)
São Paulo	Meat breeds	597	55	9.2	IFAT	1:50	Age and presence of dogs	Figliuolo et al. (2004a)
São Paulo	16 farms	1497	120	8.0	IFAT	1:25	<b>Water supply, presence of dogs, reproductive problems</b>	Machado et al. (2011)
São Paulo	8 farms	382	49	12.8	IFAT	1:25	Tested for <i>T. gondii</i>	Langoni et al. (2011)
São Paulo/Rio Grande do Sul	Abattoir	596	353	59.2	IFAT	1:25	<b>Sex, breeding system, breed, area, age</b>	Paiz et al. (2015)

Bold=statistically significant risk factors.

**Table 7.** Serological studies of antibodies to *N. caninum* in goats from Brazil.

State	Type	No. tested	No. positive	% positive	Test	Cut-off	Remarks	Reference
<b>Bahia</b>	9 herds	384	58	15.1	IFAT	1:100	<b>Breed</b>	Uzêda et al. (2007b)
<b>Maranhão</b>	5 farms	46	8	17.4	IFAT	1:25	Reproductive failure	Moraes et al. (2011)
<b>Minas Gerais</b>	90 herds	667	71	10.7	IFAT	1:50	Maximum titer 1:3,200	Andrade et al. (2013)
<b>Paraíba</b>	Abattoir	306	10	3.3	IFAT	1:50	Gender	Faria et al. (2007)
<b>Pernambuco</b>	23 farms	319	85	26.6	IFAT	1:50	<b>Age</b>	Tembue et al. (2011)
<b>Pernambuco</b>	Rural Villages	174	5	2.9	IFAT	1:50	Region, breed, age, sex	Arraes-Santos et al. (2016)
<b>Piauí</b>	Rural Villages	202	4	2.0	IFAT	1:50	Region, breed, age, sex	Arraes-Santos et al. (2016)
<b>Rio Grande do Norte</b>	14 farms	381	4	1.0	IFAT	1:50	Gender, reproductive fails, presence of dogs	Lima et al. (2008)
<b>Santa Catarina</b>	57 cities	654	30	4.6	IFAT	1:50	<b>Age, abortion, diet</b>	Topazio et al. (2014)
<b>São Paulo</b>	19 farms	394	25	6.4	IFAT	1:50	Maximum titer 1:12,800; Age, presence of dogs	Figliuolo et al. (2004b)
<b>São Paulo</b>	17 farms	923	161	17.7	NAT	1:25	<b>Presence of dogs</b> , Age, gender, reproductive problems	Modolo et al. (2008)

**Bold**=statistically significant risk factors.

**Table 8.** Serological studies of *N. caninum* antibodies in swine from Brazil.

State	Type	No. tested	No. positive	% positive	Test	Cut-off	Remarks	Reference
<b>Paraíba</b>	Abattoir	130	4	3.1	IFAT	1:50	Sex, Also tested for <i>T. gondii</i>	Azevedo et al. (2010)
<b>Paraíba</b>	Abattoir	190	6	3.2	IFAT	1:50	Also tested for <i>T. gondii</i>	Feitosa et al. (2014b)
<b>Mato Grosso do Sul</b>	Free living, wild	83	9	10.8	IFAT	1:50	<b>Sex, age</b>	Soares et al. (2016)

**Bold**=statistically significant risk factor.

**Table 9.** Serological studies of *N. caninum* in cats from Brazil.

State	Type	No. tested	No. positive	% positive	Test	Cut-off	Remarks	Reference
<b>Bahia</b>	Indoor, outdoor	272	8	2.9	IFAT	1:50	Also tested <i>Sarcocystis neurona</i>	Meneses et al. (2014)
<b>Maranhão</b>	Outdoor access	200	54	27.0	IFAT	1:25	Also tested <i>T. gondii</i>	Braga et al. (2012)
<b>Mato Grosso do Sul</b>	Free roaming and domiciled	151	10	6.6	IFAT	1:50	Also tested <i>T. gondii</i> and <i>L. infantum</i>	Sousa et al. (2014)
<b>Paraíba</b>	Free roaming and domiciled	201	0	0.0	IFAT	1:50	-	Feitosa et al. (2014a)
<b>Pernambuco</b>	Rural villages	32	2	6.2	IFAT	1:50	Region, breed, age, sex	Arraes-Santos et al. (2016)
<b>Piauí</b>	Rural villages	3	0	0.0	IFAT	1:50	Region, breed, age, sex	Arraes-Santos et al. (2016)
<b>São Paulo</b>	Free roaming and domiciled	502	60	11.9	NAT	1:40	NAT + (1+ at 1:800)	Dubey et al. (2002)
<b>São Paulo</b>	Domiciled	400	98	24.5	IB	B, C	10 IB +	Bresciani et al. (2007a)
<b>São Paulo</b>	Domiciled	70	0	0.0	IFAT	1:16	Also tested <i>S. neurona</i> 4 IFAT + 1:256	Coelho et al. (2011)

**Table 10.** Serological studies of *N. caninum* in wild animals from Brazil.

Host species	Type	No. tested	No. positive	% positive	Test	Cut-off	Reference
<b>Fam. Felidae</b>							
Caracal ( <i>Caracal caracal</i> )	Zoo	1	1	100.0	IFAT	1:25	André et al. (2010)
Fishing cat ( <i>Prionailurus viverrinus</i> )	Zoo	1	1	100.0	IFAT	1:25	André et al. (2010)
Jaguar ( <i>Panthera onca</i> )	Wild	11	7	63.6	IFAT	1:25	Onuma et al. (2014)
Jaguarundi ( <i>Puma yagouaroundi</i> syn. <i>Herpailurus yagouaroundi</i> )	Zoo	13	8	61.5	IFAT	1:25	André et al. (2010)
Lion ( <i>Panthera leo</i> )	Zoo	25	5	20.0	IFAT	1:25	André et al. (2010)
Little-spotted-cat ( <i>Leopardus tigrinus</i> )	Zoo	35	11	31.4	IFAT	1:25	André et al. (2010)
Ocelot ( <i>Leopardus pardalis</i> )	Zoo	42	30	71.4	IFAT	1:25	André et al. (2010)
Pampas cat ( <i>Oncifelis colocolo</i> )	Zoo	3	3	100.0	IFAT	1:25	André et al. (2010)
Puma-cougar ( <i>Puma concolor</i> )	Zoo	18	5	27.8	IFAT	1:25	André et al. (2010)
Serval ( <i>Leptailurus serval</i> )	Zoo	1	1	100.0	IFAT	1:25	André et al. (2010)
Tiger ( <i>Panthera tigris</i> )	Zoo	6	4	66.7	IFAT	1:25	André et al. (2010)
<b>Fam. Canidae</b>							
Crab-eating fox ( <i>Cerdocyon thous</i> )	Wild	15	4	26.6	IFAT	1:50	Cañón -Franco et al. (2004)
			1	6.6	NAT	1:40	
	Captivity	2	0	0.0	IFAT	1:50	Melo et al. (2002)
Hoary fox ( <i>Pseudalopex vetulus</i> )	Wild	30	0	0.0	IFAT	1:50	Cañón-Franco et al. (2004)
					NAT	1:40	
Maned wolf ( <i>Chrysocyon brachyurus</i> )	Captivity and wild	48	0	0.0	IFAT	1:50	Melo et al. (2002)
	Zoo and preserves	59	5	8.5	IFAT	1:25	Vitaliano et al. (2004)
Pampas fox ( <i>Lycalopex gymnocercus</i> )	Wild	12	5	41.6	IFAT	1:50	Cañón-Franco et al. (2004)
					NAT	1:40	
<b>Fam. Didelphidae</b>							
Opossum ( <i>Didelphis marsupialis</i> )	Feral	396	84	21.2	IFAT	1:25	Yai et al. (2003)
<b>Fam. Bovidae</b>							
Barbary sheep ( <i>Ammotragus lervia</i> )	Zoo	17	4	23.5	IFAT	1:50	Morikawa et al. (2014)
<b>Fam. Cervidae</b>							
Brazilian dwarf brocket ( <i>Mazama nana</i> )	Captive and zoo	40	7	17.5	IFAT	1:50	Tiemann et al. (2005b)
	Captive	22	1	4.5	ELISA	E	Zimpel et al. (2015)
Brown brocket deer ( <i>Mazama gouazoubira</i> )	Captive and zoo	66	29	43.9	IFAT	1:50	Tiemann et al. (2005b)
Marsh deer ( <i>Blastoceros dichotomus</i> )	Captive	6	1	16.7	ELISA	E	Zimpel et al. (2015)
Pampas deer ( <i>Ozotoceros bezoarticus</i> )	Wild National Park	23	3	13.0	IFAT	1:50	Tiemann et al. (2005a)
	Pantanal	16	12	75.0	IFAT	1:50	
Red brocket deer ( <i>Mazama americana</i> )	Captive PR	4	0	0.0	ELISA	E	Zimpel et al. (2015)
	Captive and zoo	29	18	62.0	IFAT	1:50	Tiemann et al. (2005b)
Rodon ( <i>Mazama rondoni</i> )	Captive and zoo	8	3	37.5	IFAT	1:50	Tiemann et al. (2005b)
Small red brocket ( <i>Mazama bororo</i> )	Captive and zoo	3	2	66.6	IFAT	1:50	Tiemann et al. (2005b)
<b>Fam. Caviidae</b>							
Capybaras ( <i>Hydrochaeris hydrochaeris</i> )	Wild	213	20	9.4	IFAT	1:25	Yai et al. (2008)
	Wild	63	1	1.5	IFAT	1:25	Valadas et al. (2010a)
	Wild	170	0	0.0	IFAT	1:50	Abreu et al. (2016)

**Table 11.** Serological studies of *N. caninum* antibodies in horses from Brazil.

State	No. tested	Type	No. positive	% positive	Assay	Cut-off titer or test	Remarks	Reference
<b>Mato Grosso</b>	200	Healthy	30	15.0	IFAT	1:50	Highest titer 1:400 in 1 horse.	Laskoski et al. (2015)
<b>Pará</b>	411	Healthy horses	28	6.8	IFAT	1:50	No risk factors detected.	Norlander (2014)
<b>Paraná</b>	72	Mares	28	38.8	IFAT	1:50	2 foals had precolostral antibodies	Locatelli-Dittrich et al. (2006)
<b>Paraná</b>	14	Pregnant mares	12	85.7	IFAT	1:50	Highest titer 1:400	Hoffmann Kormann et al. (2008)
<b>Paraná</b>	97	Healthy horses	14	14.4	IFAT	1:50	Highest titer 1:200 in 2 horses 25.7% (9/35) prevalence in mares with reproductive problem versus 6.4% (5/77) without problems. Highest titer only 1:50.	Villalobos et al. (2012)
<b>Paraná and Santa Catarina</b>	112	Mares from 5 breeding farms	14	12.5	IFAT	1:50		Abreu et al. (2014)
<b>Rio Grande do Sul</b>	241	Cart horses and Crioula breed	34	15.9	IFAT	1:50	-	Toscan et al. (2011)
<b>Rio Grande do Sul</b>	181	Pregnant mares	39	21.5	ELISA	A, C	9.3% of their paired foals had precolostral anti- <i>Neospora</i> antibodies.	Pivoto et al. (2014)
<b>Rio Grande do Sul</b>	197	Abattoir	77	39.1	IFAT	1:50	Tested for <i>Sarcocystis</i> spp. and <i>T. gondii</i>	Portella et al. (2017)
<b>Rio Grande do Sul, São Paulo, Rio de Janeiro</b>	101	Race horses	0	0.0	NAT	1:25	-	Dubey et al. (1999)
<b>Santa Catarina</b>	615	Healthy	25	4.1	IFAT	1:50	72 with history of neurological and reproductive problems.	Moura et al. (2013)
<b>São Paulo</b>	325	Healthy	19	5.8	IFAT	1:50	Highest titer 1:400	Villalobos et al. (2006)
<b>São Paulo</b>	483	Diseased	73	15.1	IFAT	1:50		
<b>São Paulo</b>	26	History of ataxia.	15	57.6	IFAT	1:2	26 cerebrospinal fluids negative.	Stelmann et al. (2011)
<b>South</b>	203	Mares	129	63.3	IFAT	1:50	Of 129, 34.8% gave birth to seropositive foals.	Antonello et al. (2012)
<b>10 states</b>	961	Old horses from abattoirs	24	2.5	ELISA	C, K	-	Hoane et al. (2006)

**Table 12.** Detection of *N. caninum* antibodies in avian species from Brazil.

Host	Tested	%positive	DNA or Antibodies	Test	Cut-off / Method	Reference
<i>Chicken (<i>Gallus gallus domesticus</i>)</i>	200 (outdoor)	47 (23.5)	Antibodies	IFAT	1:50	Costa et al. (2008)
	200 (indoor)	3 (1.5)	Antibodies	IFAT	1:50	
	10 positive	6 (60.0)	DNA	PCR	3/4 of brain Direct PCR for Nc5	
	100 farm chickens	17 (17.0) 6 (6.0)	Antibodies DNA	IFAT PCR	1:50 Np21/Np6	
<i>Sparrow (<i>Passer domesticus</i>)</i>	40	3 (7.5)	DNA	PCR	Heart and brain. Nested PCR of Nc5 and sequencing of ITS	Gondim et al. (2010)
Several species	294	0 (0.0)	Antibodies	IFAT	1:50	Mineo et al. (2011)

In one survey, 802 serum samples of female cattle from 55 dairy and beef farms from six Brazilian states (SP, RJ, MG, PR, RS, MS) were assayed by IFAT (cut-off 1:25) and 23.6% were seropositive; association between positivity to *N. caninum* and state of origin, age and production purpose was analyzed using uniform methodology (RAGOZO et al., 2003). Although seroprevalence was higher in animals older than 24 months, this difference was not statistically significant. Conclusion is not definitive because of the selection of low cut-off of 1:25. Among the six states, RJ had the lowest prevalence and MG the highest and dairy cattle had higher prevalence than beef cattle. In another large study from MS, *N. caninum* antibodies were found in cattle from 143 of 205 herds (IFAT 1:50); the cows were older than two years (OSHIRO et al., 2007). Overall seroprevalence was 14.9% (Table 4).

Even after using a higher cut-off titer (IFAT 1:200) than used most of the surveys listed in Table 4, high prevalence were recorded from MG, where *N. caninum* antibodies were detected in 23 of 24 herds with individual seroprevalence of 21.6% (BRUHN et al., 2013); and MT, where antibodies to *N. caninum* were found in 499 (53.5%) of 932 cattle samples, with at least one positive in each farm (BENETTI et al., 2009). In another report from MG, very high (91.2%; 510/559) prevalence of antibodies to *N. caninum* was found among 18 farms (GUEDES et al., 2008), revealing that *N. caninum* is well spread in the southern region of the state. They also recorded 97.2% (559/575) prevalence in cows from an abattoir.

There is an unconfirmed report that the quality of milk might be affected by neosporosis (MELO et al., 2001); an association was reported between the type of milk, classified as A, B and C according to its quality, produced in the farms and positivity to *N. caninum* antibodies, with higher occurrence in farms of MG that produce milk grade A/B than grade C; and the authors discussed about the production technology used in different farms, animals stress and commercialization. One possible cause is that the production of types A/B milk is higher and requests more from the animals, generating stress, which could be responsible for the higher prevalence.

An association between seroprevalence and age of cattle (older animals presenting higher prevalence) was detected, while feed produced on the farm was negatively associated with *N. caninum* infection in PR (GUIMARÃES et al., 2004). Also in PR, production of food in the farm, absence of artificial insemination and access of domestic and wild animals to feed facilities were associated with infection (MARTINS et al., 2012).

A strong association of *N. caninum* infection was found in farms of RS, which had presence of dogs close to the livestock and which also fed calves with colostrum pooled from several cows (CORBELLINI et al., 2006). In this study, the size of the farm was inversely associated with the presence of positive animals.

An association between beef herds and infection by *N. caninum* was reported in the Amazon region, RO, while comparing beef, dairy and mixed herds (AGUIAR et al., 2006). In that region, farms with more than 25 animals were also a risk factor. However, reproductive problems, contact with forest areas and presence of dogs were not associated with the coccidian infection.

Significant association related to animal management was found in PE (SILVA et al., 2008). Veterinary assistance, nutritional status, presence of wetlands, manipulation of the newborn calves (use of gloves when handling aborted fetuses), and destination of cows that aborted (higher in the ones that were treated with antibiotics than the ones that were discarded) were risk factors for the infection.

Pure bred Holstein cows had a higher exposure rate than mixed breeds (MUNHOZ et al., 2009) and an association between reproductive abnormalities (Table 4) (repeated estrus, repeated miscarriages and temporary anestrus) and seropositivity to *N. caninum* has been reported (BRUHN et al., 2013).

*N. caninum* is considered a primary pathogen and not influenced by concurrent BHV1, BVDV infections in dairy herds (MELO et al., 2004; MINEO et al., 2006).

Antibodies can fluctuate during pregnancy as reported, and increase in titer is not always associated with abortion (CARDOSO et al., 2009). A prospective longitudinal study was carried out in three farms in SP, during two consecutive years and the reproductive parameters were analyzed in those herds (CARDOSO et al., 2012a). In only one of the three herds the relative risk of abortion between *N. caninum* positive and negative cows was higher in the positive animals. No difference was observed regarding gestational age at abortion, repeated abortion, number of inseminations, and calving interval. A high association between *N. caninum* serological status of dams and daughters were observed in a longitudinal study carried out in three farms from SP, confirming the importance of vertical transmission, but there was no difference in the culling rate between positive and negative cows (CARDOSO et al., 2012b). A significant relationship between seropositivity of cattle and their offspring was also found in PE (RAMOS et al., 2016), which had a rate of transplacental transmission of 72.2% (13/18) for adults and 69.2% (9/13) for heifers by IFAT and 43.5% (17/39) for adults and 50.0% (9/18) for heifers by ELISA, concluding that vertical transmission is the major form of infection in this region.

**Buffaloes:** Studies with serum samples from Brazilian buffaloes (Table 5) showed occurrence of *N. caninum* antibodies varying from 14.6% to 88%. However, despite the high occurrence values, no reproductive disorders were reported in those groups (FUJII et al., 2001a, b; GENNARI et al., 2005; GONDIM et al., 2007)

**Sheep:** Seroprevalences ranged from 1.8% to 64.2% (Table 6). Some surveys stated as risk factors: abortion in the flock, presence of dogs, extensive husbandry systems and breed of the sheep. Great part of the studies also evaluated the presence of *T. gondii* infection, which usually showed a higher prevalence. In 2017, Filho et al. (2017) studied the vertical transmission rate of 50 naturally infected sheep, analysed by in house ELISA for six months. The initial prevalence of infection was 26.0% (13/50) and by the end of the study it had increased to 72% (36/50), being the vertical transmission rate 11%, which one sheep out of nine from a group gave birth to two infected ewes (IFAT 1:25).

**Goats:** In goats, serological surveys found rates between 1.0% (LIMA et al., 2008) to 26.6% (TEMBUE et al., 2011) of prevalence (Table 7). A study to evaluate infections by *T. gondii*, *N. caninum* and caprine arthritis-encephalitis virus (CAEV) was conducted, finding a prevalence of 37.81%, 23.62% and 17.23%,

respectively (COSTA et al., 2012). The results indicate that CAEV does not predispose goats to infection by *T. gondii* or *N. caninum*. However, when CAEV/*T. gondii* or CAEV/*N. caninum* infection were detected, occurrence of reproductive failure was higher, maybe related to poor husbandry conditions. Differential diagnosis in cases of abortions in small ruminants is highly desirable.

Also, *N. caninum* DNA was found in brains of goats from BA. Silva et al. (2009) analyzed brains, hearts and tongues of 102 goats from slaughterhouses, and found a frequency of 1, 96% (2/102), using primers for ITS-1 region (JS4/CT2b; CT1/CT2).

**Pigs:** The prevalence of anti-*N. caninum* antibodies is low in pigs, and further studies are needed to evaluate the role of this species in the epidemiology of the parasite, including attempts to isolate viable parasites. Table 8 summarizes the studies with pigs from Brazil.

*N. caninum* antibodies were also found in feral pigs (*Sus scrofa*) in 10.8% (9/83) of samples from Pantanal, MS, with titers up to 1:800 by IFAT (SOARES et al., 2016).

**Cats:** In Brazil, studies determining the prevalence of antibodies in serum of cats (Table 9) found results that range from 0%, in cats from Andradina, SP, and Patos, PB (COELHO et al., 2011; FEITOSA et al., 2014a) to 27% in cats from MA (BRAGA et al., 2012). In animals from SP, NAT antibodies were found in 60 of 502 (12%) (DUBEY et al., 2002). The samples with titers greater than 1:80 were also examined by IB, as a confirmatory test, being ten of the 24 cats (41.6%) positive for both tests.

**Wild animals:** *N. caninum* antibodies were detected in sera of wild animals kept in captivity or trapped in the wild of the Families: Canidae, Felidae, Didelphidae, Bovidae, Caviidae and Cervidae (Table 10).

Among wild herbivores, *Neospora* has been better documented in cervids from Pantanal region (MS) (TIEMANN et al., 2005a). In that study, serum samples from 23 pampas deer (*Ozotoceros bezoarticus*) from the National Park of Emas, in GO, and 16 captured in bovine's farms, from Pantanal region, in MT, were tested for the presence of *N. caninum* antibodies. They found 13% and 75% positivity, respectively, for the deer that live inside the park and the ones from Pantanal, which is close to farms indicating that this proximity of wild and domestic animals could increase the occurrence of *N. caninum* infection among deer. From the Zoo of Curitiba, PR, 17 samples from Barbary sheep (*Ammotragus lervia*), from the Bovidae Family, were examined and four (23.5%) presented *N. caninum* antibodies (MORIKAWA et al., 2014).

Yai et al. (2003) tested 396 feral opossums (*Didelphis marsupialis*) samples from different regions of the city of São Paulo, and 21.2% (84/396) were positive.

Sera from 14 species of wild felids from zoos were tested for the presence of *N. caninum* (Table 10) antibodies and 12 species had at least one positive animal (ANDRÉ et al., 2010). In addition, 11 serum samples from free range jaguars (*Panthera onca*) from Pantanal were examined and seven (63.6%) were positive (ONUMA et al., 2014). In both studies, IFAT with a cut-off of 1:25 and anti-cat commercial conjugate was used for IgG antibody determination.

Despite the importance of wild canids in the epidemiology of *N. caninum*, few studies are available in Brazil. Crab-eating fox (*Cerdocyon thous*) is one of most common canids of South America,

(Gennari, own observations) and four of these foxes were fed with masseter muscle and brain of two *N. caninum* seropositive bovines (IFAT >200). The foxes received the inoculum in two consecutive days. Two animals excreted *Hammonia heydorni* oocysts on eight and nine dpi but not *Neospora* oocysts (SOARES et al., 2009b). By means of molecular techniques, Nascimento et al. (2015) identified *N. caninum* in brain of six from 49 (12.2%) hoary foxes (*Pseudalopex vetulus* syn. *Lycalopex vetulus*) from PB. The molecular identification of the amplified products by sequencing reaction, using Nc-5 gene, presented 99% similarity with *N. caninum*.

Similarly, Muradian et al. (2012) tested wild urban rodents tissues (Family Muridae), but did not detect *N. caninum* DNA in four mice (*Mus musculus*), 20 brown rats (*Rattus norvegicus*), and 193 black rats (*Rattus rattus*) from São Paulo city. Regarding capybaras, the first study done (YAI et al., 2008) tested animals from 11 counties in SP by IFAT (1:25) and found a prevalence of 9.4% (20/213), suggesting that they can serve as a source of *N. caninum* infection for wild canids. Also in SP, 63 capybaras were examined for *N. caninum* by IFAT (1:25) and other diseases and found two positive animals and one of them was positive for both *T. cruzi* and *N. caninum*, but no association was observed (VALADAS et al., 2010a). Recently, 170 samples of capybaras from SP were analyzed, but none were positive, although 17 (10%) were positive for *T. gondii* (ABREU et al., 2016). DNA of *N. caninum* was found in capybaras from PR (TRUPPEL et al., 2010), in 23% (6/26) of the studied animals. Parasite DNA, aiming the Nc5 gene was found in the liver and lymph nodes and ITS-1 was found in blood, liver, heart and lymph nodes.

**Horses:** At present, it is uncertain if horses are infected with both *N. caninum* and *N. hughesi* (DUBEY et al., 2017). Serosurveys on horses conducted in Brazil are summarized in Table 11. *N. caninum* antigen was used in all studies with exception of Hoane et al. (2006), that used *N. hughesi* SAG1 (NhSAG1) in an in-house ELISA.

A low rate of infection (0.4%, IFAT 1:100) was found among 500 donkeys (*Equus asinus*) sampled in BA; positive cases were confirmed by IB (GALVÃO et al., 2015). In a previous limited survey carried out in PA, no seropositive donkeys (n=6) or mules (n=9) were found (NORLANDER, 2014). Recently, a 2% seroprevalence was reported in donkeys (n=333) from five northeastern states (AL, PB, PE, PI and RN) using IFAT (1:50). In all these studies, *N. caninum* NC-1 strain was used as antigen source (GENNARI et al., 2016).

In PR, antibodies to *Neospora* were detected in two foals (LOCATELLI-DITTRICH et al., 2006), but the information is not definitive because of the low titer (1:50) detected. In addition, Antonello et al. (2012), by IFAT (1:50, *N. caninum* antigen), reported a high prevalence (63.3%) of *Neospora* antibodies in sera of 203 thoroughbred mares and their foals before suckling in two farms from Southern Brazil. A high percentage (34.8%) of seropositive mares gave birth to seropositive foals. Additionally, 8% of seronegative mares gave birth to seropositive foals. Mare sera were titrated further to 1:200 dilution and the seropositivity decreased to 33%, but foal sera were not titrated. In another study performed in RS, *Neospora* antibodies were found in 21.5% of 181 mares and in 9.3% of their foals in pre-suckling sera (PIVOTO et al., 2014); in this case, antibodies were assayed using an in-house indirect

ELISA, with NC-1 strain and soluble protein from tachyzoites maintained in CV-1 cells as antigen. Low levels of maternal IgG can cross the placenta in mares. Therefore, further studies are needed to confirm results of these investigations from Brazil (DUBEY et al., 2017).

**Avian species:** Table 12 summarizes the surveys of detection of *N. caninum* antibodies detection in avian species from Brazil. Seroprevalence was higher in free range chickens; 23.5% of 200 outdoors and 1.5% of 200 indoors chickens (*Gallus gallus*), in BA were seropositive (COSTA et al., 2008). In the same study, the authors found positive results for PCR from 6/10 (60%) seropositive chickens. 40 of 293 wild sparrows (*Passer domesticus*) from BA and PE were seropositive and *N. caninum* DNA by PCR was detected in three (7.5%) of brain and heart from 40 animals (GONDIM et al., 2010).

In a study by serological and histological methods none of the 294 wild and captive birds from nine avian orders had *N. caninum* antibodies. However in two psittacine birds, Apicomplexa-like tissue cysts were found and were immunostained positive with *N. caninum* antisera (MINEO et al., 2011).

Results of the experimental infection in birds from Brazil are summarized in Table 13. In the first Brazilian study (FURUTA et al., 2007) 50 chickens were inoculated with *N. caninum* tachyzoites, using different doses; chickens seroconverted but remained healthy. In 15 euthanized chickens, *N. caninum* tachyzoites were reported to be present in different tissues by IHC, at 15 dpi; however, no illustrations were provided. At the termination of the experiment (60 dpi), all chickens were seronegative by IFAT (<1:20) *N. caninum* was not found by IHC. In laying hens, no evidence of vertical transmission was found. However, 50% of embryonated chicken eggs inoculated with *N. caninum* died and

chickens that hatched 21 dpi had neurological signs. Dogs fed chorioallantoic membranes and whole infected eggs seroconverted and excreted *N. caninum* oocysts in their feces, as confirmed by PCR (Nc5) (FURUTA et al., 2007).

In an unconfirmed report, four littermate two-month old dogs fed with chorioallantoic membranes previously infected *in ovo* with  $10^6$ NC-1 strain tachyzoites were euthanized 3, 4, 5 and 6 dpi (MUNHOZ et al., 2013). The authors reported immunoreactivity to *N. caninum* in lesions in lungs, spleen, and small and large intestine but strangely *N. caninum* DNA was not detected in affected tissues by conventional PCR. The gross lesions depicted resemble bacterial septicemia and the results need confirmation. The dogs did not excrete oocysts. In a follow up study, the authors, infected 90-days old chickens and embryonated eggs with  $10^6$  *N. caninum* tachyzoites using NC-1 strain and fed three dogs with infected organs but the dog did not excrete oocysts (MUNHOZ et al., 2014). Although there is no confirmation by IHC and microscopy, *N. caninum* DNA was found in the spleen and pectoral muscles of one of the birds born from the inoculated eggs (MUNHOZ et al., 2014).

An attempt to infect quails (n=58) with doses of  $3.5 \times 10^6$  and  $5 \times 10^6$  of *N. caninum* tachyzoites (NC-Bahia) was largely unsuccessful (OLIVEIRA et al., 2013). Although there was evidence of transitory infection (seroconversion and finding of parasite DNA) at 14 dpi, the quails became seronegative at 30 dpi with no demonstrable parasite DNA or antigen in their tissues; two dogs fed quail tissues did not excrete oocysts.

Experimental infection was also conducted in pigeons (*Columba livia*) inoculated with the  $10^7$  NC-1 tachyzoites (MINEO et al., 2009). The pigeons developed transitory *N. caninum* antibodies starting at 5 dpi but at the end of the experimental period of

**Table 13.** Detection of *N. caninum* in avian from experimental studies in Brazil.

Host	No. animals	Dose	Tests	Results	Reference
<b>Chickens</b>					
( <i>Gallus gallus domesticus</i> )	7-days old	$10^3, 10^4, 10^5, 10^6$	Killed 15,30,45,60 dpi	No mortality; Dpi 15:IFAT+ IHC+	Furuta et al. (2007)
	Laying eggs	$10^8$	Bioassay, PCR, Histopathology	Dpi 60: IFAT - IHC -	
	Embrionated eggs	$10^3, 10^4, 10^5, 10^6$	Histopathology Dogs bioassay	Eggs- Mortality: 18-21 days incubation Dogs shed oocyst	
	90-days old	$3 \times 10^6$	IFAT, IHC, Histopathology, Bioassay	IFAT+	Munhoz et al. (2014)
	Embrionated eggs	$1 \times 10^2$	2 embryos died	PCR+ spleen, muscles (1bird)	
<b>Pigeons</b>					
( <i>Columba livia</i> )	4	$1 \times 10^7$ Blood samples each 5 days	IFAT	1:20; peak 10-20dpi	Mineo et al. (2009)
	4	control	IHC +	Lungs, heart, CNS	
<b>Quails</b>					
( <i>Coturnix coturnis japonica</i> )	40	$3.5 \times 10^6$	Histopathology	+	Oliveira et al. (2013)
	8	$5.0 \times 10^6$	IHC	+	
	10	control	PCR	+	

45 days, all birds were seronegative. One infected pigeon died at 25 dpi and *N. caninum* and lesions were found by IHC in lungs, heart, central nervous system, liver, spleen and kidney.

Rezende-Gondim et al. (2017) cultivated *N. caninum*, using a chicken cell line, and temperatures between 39 °C and 41.5 °C. Multiplication of *N. caninum* tachyzoites in vitro was inhibited at temperatures similar to those of chickens. The authors concluded that the avian body temperatures may be one of the reasons that isolation of the parasite is difficult in avian species.

In summary, avians are a poor host for *N. caninum*, based on failure to isolate viable parasite from naturally infected tissues and failure to induce chronic infection in experimentally infected birds.

### Clinical infections

**Dogs:** There are two confirmed reports of clinical neosporosis in adult dogs. A seven-year-old male Collie from Salvador, BA developed incoordination and rear limb paralysis (GONDIM et al., 2001). The dog was found to have a *N. caninum* IFAT titer of 1:1600. In spite of medication with Clindamycin (22 mg/kg) for 14 days the dog died. A necropsy was performed. Histologically there was encephalitis associated with tachyzoites and tissue cysts and the diagnosis was confirmed by IHC examination. Live tissue cysts were found in squash preparations from the dog brain. Tissue cysts were found in the brains of gerbils inoculated with brain homogenate of the dog. Viable *N. caninum* was propagated in cell cultures seeded with infected gerbil brain.

The second case was from a 10 year old dog from RS (MANN et al., 2016). Persistent dermal lesions with multifocal ulcerative nodules on the neck and pelvic limbs were observed. The dog had an IFAT *N. caninum* antibody titer of 1:6400. Cytological examination of the exudates from nodules showed pyogranulomatous inflammation with tachyzoites and *N. caninum* was identified by PCR. The dog was medicated with Clindamycin (6 mg/kg) for 28 days with resolution of lesions. However, the dermal lesions with identifiable tachyzoites reappeared 12 days after the cessation of treatment, perhaps due to a very low dose of Clindamycin used; the usual treatment is 20-25 mg/kg.

In addition to these confirmed reports, Langoni et al. (2012) isolated *N. caninum* by bioassay in gerbils inoculated with brains of two of seven dogs with neurological disorders. No other details were given about these dogs or the strain of *Neospora*, therefore, not included in Table 1.

**Cattle:** Reports of confirmed neosporosis abortion from six states are summarized in Table 14. Apparently, all of these cases were sporadic abortion. Corbellini et al. (2006) investigated 161 bovine abortions from 149 farms bovine abortions during 2001 and 2003 from RS. Causes of abortion were identified in 83 (51.5%) cases. *N. caninum* was the most important cause, and identified in 37 fetuses; in 34 fetuses, the diagnosis was confirmed by IHC examination. Overall, 37 of 161 (23%) fetuses were infected solely with *N. caninum*. In six cases, there was concurrent *Leptospira* sp. infection. Most aborted fetuses were 4.4 months gestational age. Cows aborting a *Neospora* infected fetuses were 2.4 times likely to have aborted previously.

In a follow up publication, more detailed investigation involved 258 aborted fetuses from RS and SC. Lesions indicative of neosporosis were found in 89 (34%) of these 258 submissions based on histopathology. The diagnosis was confirmed by IHC in 55 of these 89 cases (PESCADOR et al., 2007). A striking feature was the distribution of lesions; myositis in 92%, myocarditis in 76% and pneumonitis and encephalitis in 75%. Two of these fetuses had grossly visible pale white foci in lungs, indicative of necrosis. Overall, *N. caninum* was the predominant (21.3% of 258) cause of abortion in this investigation. A similar conclusion was reached by Cabral et al. (2009), who combined histopathology, IHC, and PCR to diagnose *N. caninum* in 24.8% of 105 fetal samples from the state of SP. They detected *N. caninum* DNA in the brains of 22% (16 of 72), placenta of 20.0% (4 of 20), heart and liver of 16.3% (8 of 49) and pool of kidney, lungs, and spleen of 10.9% (7 of 64).

Additional provisional evidence for neosporosis was based on higher seropositivity in aborting versus non-aborting cows on a given farm (Table 15). However, most of these reports were provisional and not a case controlled study. In PR, samples from a herd collected over a nine-yr follow-up period were analyzed for *Neospora* infections (LOCATELLI-DITTRICH et al., 2001). They found that the proportion of abortions was 20% and 8%, respectively for the seropositive and seronegative animals.

As mentioned in Table 1, Locatelli-Dittrich et al. (2003, 2004) isolated viable *N. caninum* from a seven-month gestational age fetus and a three-month old blind calf. However, the etiology was not confirmed because histological examination was not performed.

**Goats:** Fatal neosporosis has been reported from MG and RS (Table 16). Hydrocephalus was detected in a day old goat kid, which had a high titer of antibodies on IFAT (1:400) for

**Table 14.** Reports of *N. caninum*-associated abortion in cattle from Brazil.

State	No. aborted	No. positive (%)	Diagnosis			Reference
			Histo	IHC	PCR	
Bahia	1	1	Yes	Yes	ND	Gondim et al. (1999b)
Goiás	195 dead fetuses from abattoir	40 9(20.5)	No	No	Yes	Brom et al. (2014)
Paraná	34	8 (23.5)	Yes	Yes	ND	Santos et al. (2005)
Rio Grande do Sul	30	1 (3.3)	Yes	Yes	ND	Corbellini et al. (2000)
Rio Grande do Sul	46	22 (47.8)	Yes	Yes	ND	Corbellini et al. (2002)
Rio Grande do Sul	161	37 (17.4)	Yes	Yes	ND	Corbellini et al. (2006)
Rio Grande do Sul, Santa Catarina	258	55 (21.3)	Yes	Yes	ND	Pescador et al. (2007)
São Paulo	105	26 (24.7)	Yes	Yes	Yes	Cabral et al. (2009)

**Table 15.** Seropositivity as evidence of abortion in cattle from Brazil.

Total number of cattle/ farms examined	Aborting			Non aborting or control			Reference
	Test cut-off	No. tested	No. seropositive (%)	No. tested	No. seropositive (%)	Risk of abortion indicated by odd ratio (OR), significant association (SA), remarks	
223 dairy cows, 5 herds	IFAT, 1:200	NS	NS (23.3)	NS	NS (8.3)	OR 3.3	Corbellini et al. (2002)
2448 cattle from 205 herds, beef, dairy	IFAT, 1:50	-	55/68 herds (80.9)	-	84/134 herds (62.7)	OR 2.5	Oshiro et al. (2007)
1256, 41 aborted	IFAT, 1:200	41	24 (58.5)	1215	199 (16.4)	OR 7.2	Hein et al. (2012)
1204 dairy cows from 40 farms	IFAT, 1:200	NS	NS (31.1)	NS	NS (17.7)	OR 1.98	Bruhn et al. (2013)
621 cattle, 63 farms, 36 farms with abortion	IFAT, 1:100	-	26/36 farms (72.2)	-	12/27 farms (44.4)	SA	Boas et al. (2015)
3428 cattle from 174 herds	IFAT, 1:100	-	99/108 herds (91.7)	-	11/52 herds (21.1)	SA	Chiebalo et al. (2015)
1273 cattle from 6 dairy herds	IFAT, 1:200	305	122 (40.0)	968	40 (4.1)	SA	Pessoa et al. (2016)

**Table 16.** Fatal neosporosis in goats from Brazil.

State	Case no.	History	Diagnosis				Reference
			Serology	Histology	Immuno	PCR	
Minas Gerais	1- day old	Born weak, unable to nurse	Doe, IFAT 1:400	White matter absent, mild necrosis, perivasculitis, only tissue cysts 9.8-20.5 µm in diameter.	Positive	Not done	Varaschin et al. (2012)
	Day of birth	New born kid, late term	IFAT doe 1:800, presuckling kid 1:400	No lesions in placenta	Negative	Positive in placenta.	
	Abortion	A chronically infected aborted 4 fetuses, 87 days after mating.	IFAT doe 1:6,400 at abortion day.	Positive in 1.	Brain positive in 1 fetus	PCR positive brain in the first and heart in the second fetus	Mesquita et al. (2013)
	2 stillborn	Stillborn on 148 days after mating of a chronically infected doe	IFAT doe 1:3,200 at parturition.	Positive in 2.	Positive in both.	Positive in CNS of both.	
Rio Grande do Sul	1- day old	Born weak, unable to nurse, ataxic, euthanized day 3	No data	Encephalitis, more severe in mid brain, many intact and degenerating tissue cysts of 12.4-32.2 µm in diameter.	Positive	Not done	Corbellini et al. (2001)

*N. caninum* and no antibodies for *T. gondii* (VARASCHIN et al., 2012). The cerebral hemispheres of the animal were asymmetrical, the gyri were swollen, and ventricles were expanded. Histologically, there were only tissue cysts in the brain, no tachyzoites were observed and no lesions were observed in histological examination of placenta. Corbellini et al. (2001) described another case in southern Brazil, with a kid presenting neurological signs as ataxia and opistothonus, which got more severe when it was three days old, and was euthanized. Brain, heart, lungs and liver had microscopic lesions, but no grossly lesions were not observed.

**Humans:** Currently, there is no evidence of the zoonotic character of *N. caninum*, although antibodies in humans were

found in immune compromised and normal patients in different parts of the world (DUBEY et al., 2017).

In Brazil, three studies were conducted. In MG, serum samples from HIV-infected patients (Group I), patients with neurological disorders (Group II), newborns (Group III) and a healthy control (Group IV) by IFAT (1:50) were assayed for *N. caninum* antibodies by indirect ELISA to detect immunoglobulin G (IgG), utilizing soluble antigen and IB as a confirmatory test, with lysed tachyzoites as antigen (LOBATO et al., 2006). They found IB positive patients in all groups, and HIV-infected patients and those with neurological disorders patients presented significantly higher prevalence. Combining all the tests, 37.7% (23/61) of group I,

18% (9/50) of group II, and 5% (5/91) from groups III had antibodies to *N. caninum*.

In the second study, antibodies to *N. caninum* (IFAT, 1:50) were sought in Human Immunodeficiency Virus (HIV) patients; which 26.1% (81/310) patients from MS and 31.2% (10/31) patients from PR were seropositive and one patient had a titer of 1:400 (OSHIRO et al., 2015). In the third report, *N. caninum* antibodies (IFAT 1:100) were detected in 7 of 67 (10.5%) humans from 24 farms of MT (BENETTI et al., 2009).

### Prospective and areas of future research

It is evident from this review that *N. caninum* infection is widely prevalent throughout Brazil. However, nothing is known of the prevalence of *N. caninum* oocysts in soil or in canine feces. It is also uncertain if there are additional definitive hosts, other than the domestic dog and some wild canids, as stated before. Overall, little is known of clinical neosporosis, particularly in cattle. The few reports pertain to sporadic cases of abortion with no information on epidemics or storms of abortion. There is need for a national survey in cattle using defined parameters. Future researches should focus on molecular characterization of *N. caninum* strains, possibility of vaccine production and relationship between wildlife and livestock epidemiology.

## References

- Abreu JAP, Krawczak FS, Nunes FP, Labruna MB, Pena HFJ. Anti-*Toxoplasma gondii* and anti-*Neospora caninum* antibodies in capybaras (*Hydrochoerus hydrochaeris*) from Itu Municipality, São Paulo. *Rev Bras Parasitol Vet* 2016; 25(1): 116-118. PMid:26982562. <http://dx.doi.org/10.1590/S1984-29612016002>.
- Abreu RA, Weiss RR, Thomaz-Soccol V, Locatelli-Dittrich R, Laskoski LM, Bertol MAF, et al. Association of antibodies against *Neospora caninum* in mares with reproductive problems and presence of seropositive dogs as a risk factor. *Vet Parasitol* 2014; 202(3-4): 128-131. PMid:24731383. <http://dx.doi.org/10.1016/j.vetpar.2014.03.022>.
- Acosta ICL, Centoducate LD, Soares HS, Marcili A, Gondim MFN, Rossi JL, et al. Occurrence of *Neospora caninum* and *Toxoplasma gondii* antibodies in dogs from rural properties surrounding a biological reserve, Espírito Santo, Brazil. *Rev Bras Parasitol Vet* 2016; 25(4): 536-539. PMid:27925068. <http://dx.doi.org/10.1590/s1984-29612016075>.
- Aguiar DM, Cavalcante GT, Rodrigues AAR, Labruna MB, Camargo LMA, Camargo EP, et al. Prevalence of anti-*Neospora caninum* antibodies in cattle and dogs from Western Amazon, Brazil, in association with some possible risk factors. *Vet Parasitol* 2006; 142(1-2): 71-77. PMid:16857319. <http://dx.doi.org/10.1016/j.vetpar.2006.06.014>.
- Aguiar DM, Chiebano DP, Rodrigues AAR, Cavalcante GT, Labruna MB, Gennari SM. Prevalência de anticorpos anti-*Neospora caninum* em ovinos do município de Monte Negro, RO, Amazônia Ocidental Brasileira. *Arg Inst Biol* 2004; 71: 616-618.
- Al-Qassab S, Reichel MP, Ellis J. A second generation multiplex PCR for typing strains of *Neospora caninum* using six DNA targets. *Mol Cell Probes* 2010; 24(1): 20-26. PMid:19683051. <http://dx.doi.org/10.1016/j.mcp.2009.08.002>.
- Al-Qassab S, Reichel MP, Ivens A, Ellis JT. Genetic diversity amongst isolates of *Neospora caninum*, and the development of a multiplex assay for the detection of distinct strains. *Mol Cell Probes* 2009; 23(3-4): 132-139. PMid:19496247. <http://dx.doi.org/10.1016/j.mcp.2009.01.006>.
- Alves AF No, Bandini LA, Nishi SM, Soares RM, Driemeier D, Antoniassi NAB, et al. Viability of sporulated oocysts of *Neospora caninum* after exposure to different physical and chemical treatments. *J Parasitol* 2011; 97(1): 135-139. PMid:21348620. <http://dx.doi.org/10.1645/GE-2571.1>.
- Andrade GS, Bruhn FRP, Rocha CMBM, Guimarães AS, Gouveia AMG, Guimarães AM. Seroprevalence and risk factors for *Neospora caninum* in sheep in the state Minas Gerais, southeastern Brazil. *Vet Parasitol* 2012; 188(1-2): 168-171. PMid:22475416. <http://dx.doi.org/10.1016/j.vetpar.2012.03.006>.
- Andrade GS, Bruhn FRP, Rocha CMBM, Guimarães AS, Gouveia AMG, Guimarães AM. Seroprevalence for *Neospora caninum* in goats of Minas Gerais state, Brazil. *Res Vet Sci* 2013; 94(3): 584-586. PMid:23200513. <http://dx.doi.org/10.1016/j.rvsc.2012.09.026>.
- André MR, Adania CH, Teixeira RHF, Silva KF, Jusi MMG, Machado STZ, et al. Antibodies to *Toxoplasma gondii* and *Neospora caninum* in captive neotropical and exotic wild canids and felids. *J Parasitol* 2010; 96(5): 1007-1009. PMid:20950109. <http://dx.doi.org/10.1645/GE-2502.1>.
- Andreotti R, Barros JC, Pereira AR, Oshiro LM, Cunha RC, Figueiredo LF No. Association between seropositivity for *Neospora caninum* and reproductive performance of beef heifers in the Pantanal of Mato Grosso do Sul, Brazil. *Rev Bras Parasitol Vet* 2010; 19(2): 119-123. PMid:20624350. <http://dx.doi.org/10.1590/S1984-29612010000200010>.
- Andreotti R, Matos MFC, Gonçalves KN, Oshiro LM, Lima-Junior MSC, Paiva F, et al. Comparison of indirect ELISA based on recombinant protein NcSRS2 and IFAT for detection of *Neospora caninum* antibodies in sheep. *Rev Bras Parasitol Vet* 2009; 18(2): 19-22. PMid:19602311. <http://dx.doi.org/10.4322/rbpv.01802004>.
- Andreotti R, Oliveira JM, Silva EA, Oshiro LM, Matos MFC. Occurrence of *Neospora caninum* in dogs and its correlation with visceral leishmaniasis in the urban area of Campo Grande, Mato Grosso do Sul, Brazil. *Vet Parasitol* 2006; 135(3-4): 375-379. PMid:16310954. <http://dx.doi.org/10.1016/j.vetpar.2005.10.011>.
- Andreotti R, Pinckney RD, Pires PP, Silva EAE. Evidence of *Neospora caninum* in beef cattle and dogs in the state of Mato Grosso do Sul, center-western region, Brazil. *Rev Bras Parasitol Vet* 2004; 13(3): 129-131.
- Antonello AM, Pivoto FL, Camillo G, Braunig P, Sangioni LA, Pompermayer E, et al. The importance of vertical transmission of *Neospora* sp. in naturally infected horses. *Vet Parasitol* 2012; 187(3-4): 367-370. PMid:22436425. <http://dx.doi.org/10.1016/j.vetpar.2012.02.005>.
- Araeas-Santos AI, Araújo AC, Guimarães MF, Santos JR, Pena HFJ, Gennari SM, et al. Seroprevalence of anti-*Toxoplasma gondii* and anti-*Neospora caninum* antibodies in domestic mammals from two distinct regions in the semi-arid region of Northeastern Brazil. *Vet Parasit: Reg Stud Rep* 2016; 5: 14-18.
- Azevedo SS, Batista CSA, Vasconcellos SA, Aguiar DM, Ragazzo AMA, Rodrigues AAR, et al. Seroepidemiology of *Toxoplasma gondii* and *Neospora caninum* in dogs from the state of Paraíba, Northeast region of Brazil. *Res Vet Sci* 2005; 79(1): 51-56. PMid:15894024. <http://dx.doi.org/10.1016/j.rvsc.2004.10.001>.
- Azevedo SS, Pena HFJ, Alves CJ, Guimarães AAM, Oliveira RM, Maksimov P, et al. Prevalence of anti-*Toxoplasma gondii* and anti-*Neospora caninum* antibodies in swine from Northeastern Brazil. *Rev Bras Parasitol*

Vet 2010; 19(2): 80-84. PMid:20624342. <http://dx.doi.org/10.1590/S1984-29612010000200002>.

Balthazar LMC, Leal PDS, Teixeira Filho WL, Lopes CWG. Cães sororreagentes a *Neospora caninum* (Apicomplexa: Toxoplasmatinae) atendidos em uma clínica veterinária na cidade do Rio de Janeiro, RJ. Rev Bras Med Vet 2013; 35(S2): 48-51.

Bandini LA, Neto AFA, Pena HFJ, Cavalcante GT, Schares G, Nishi SM, et al. Experimental infection of dogs (*Canis familiaris*) with sporulated oocysts of *Neospora caninum*. Vet Parasitol 2011; 176(2-3): 151-156. PMid:21094584. <http://dx.doi.org/10.1016/j.vetpar.2010.10.047>.

Benetti AH, Schein FB, Santos TR, Toniollo GH, Costa AJ, Mineo JR, et al. Pesquisa de anticorpos anti-*Neospora caninum* em bovinos leiteiros, cães e trabalhadores rurais da região Sudoeste do Estado de Mato Grosso. Rev Bras Parasitol Vet 2009;18(Suppl 1): 29-33. PMid:20040187. <http://dx.doi.org/10.4322/rbpv.018e1005>.

Benetti AH, Toniollo GH, Santos TR, Gennari SM, Costa AJ, Dias RA. Ocorrência de anticorpos anti-*Neospora caninum* em cães no município de Cuiabá, Mato Grosso. Ciênc Anim Bras 2008; 9(1): 177-180.

Bezerra MA, Pereira LM, Bononi A, Biella CA, Baroni L, Pollo-Oliveira L, et al. Constitutive expression and characterization of a surface SRS (NCsRS67) protein of *Neospora caninum* with no orthologue in *Toxoplasma gondii*. Parasitol Int 2017; 66(2): 173-180. PMid:28108401. <http://dx.doi.org/10.1016/j.parint.2017.01.010>.

Boas RV, Pacheco TA, Melo ALT, Oliveira ACS, Aguiar DM, Pacheco RC. Infection by *Neospora caninum* in dairy cattle belonging to family farmers in the northern region of Brazil. Rev Bras Parasitol Vet 2015; 24(2): 204-208. PMid:26154960. <http://dx.doi.org/10.1590/S1984-29612015035>.

Boaventura CM, Oliveira VSF, Melo DPG, Borges LMF, Silva AC. Prevalência de *Neospora caninum* em cães de Goiânia. Rev Patol Trop 2008; 37(1): 15-22. <http://dx.doi.org/10.5216/rpt.v37i1.4027>.

Braga MSCO, Andre MR, Jusi MMG, Freschi CR, Teixeira MCA, Machado RZ. Occurrence of anti-*Toxoplasma gondii* and anti-*Neospora caninum* antibodies in cats with outdoor access in São Luís, Maranhão, Brazil. Rev Bras Parasitol Vet 2012; 21(2): 107-111. PMid:22832749. <http://dx.doi.org/10.1590/S1984-29612012000200007>.

Brasil AWL, Parentoni RN, Feitosa TF, Bezerra CS, Vilela VLR, Pena HFJ, et al. Risk factors for *Toxoplasma gondii* and *Neospora caninum* seropositivity in buffaloes in Paraíba State, Brazil. Rev Bras Parasitol Vet 2015; 24(4): 459-463. PMid:26689181. <http://dx.doi.org/10.1590/S1984-29612015066>.

Bresciani KDS, Costa AJ, Nunes CM, Serrano ACM, Moura AB, Stobbe NS, et al. Ocorrência de anticorpos contra *Neospora caninum* e *Toxoplasma gondii* e estudo de fatores de risco em cães de Araçatuba - SP. Ars Vet 2007b; 23: 40-46.

Bresciani KDS, Gennari SM, Serrano ACM, Rodrigues AAR, Ueno T, Franco LG, et al. Antibodies to *Neospora caninum* and *Toxoplasma gondii* in domestic cats from Brazil. Parasitol Res 2007a; 100(2): 281-285. PMid:16941188. <http://dx.doi.org/10.1007/s00436-006-0262-4>.

Brom PRF, Regidor-Cerrillo J, Collantes-Fernández E, Ortega-Mora LM, Guimarães MS, Silva AC. Genetic characterisation of *Neospora caninum* strains from clinical samples of zebuine foetuses obtained in abattoirs in Goiás, Brazil. Vet Parasitol 2014; 204(3-4): 381-387. PMid:24893690. <http://dx.doi.org/10.1016/j.vetpar.2014.05.011>.

Bruhn FRP, Daher DO, Lopes E, Barbieri JM, Rocha CMBM, Guimarães AM. Factors associated with seroprevalence of *Neospora caninum* in dairy

cattle in southeastern Brazil. Trop Anim Health Prod 2013; 45(5): 1093-1098. PMid:23212838. <http://dx.doi.org/10.1007/s11250-012-0330-y>.

Bruhn FRP, Figueiredo VC, Andrade GS, Costa-Júnior LM, Rocha CMBM, Guimarães AM. Occurrence of anti-*Neospora caninum* antibodies in dogs in rural areas in Minas Gerais, Brazil. Rev Bras Parasitol Vet 2012; 21(2): 161-164. PMid:22832759. <http://dx.doi.org/10.1590/S1984-29612012000200017>.

Cabral AD, Camargo CN, Galleti NTC, Okuda LH, Pituco EM, Del Fava C. Diagnosis of *Neospora caninum* in bovine fetuses by histology, immunohistochemistry, and nested-PCR. Rev Bras Parasitol Vet 2009; 18(4): 14-19. PMid:20040203. <http://dx.doi.org/10.4322/rbpv.01804003>.

Cañón-Franco WA, Bergamaschi DP, Labruna MB, Camargo LMA, Souza SLP, Silva JCR, et al. Prevalence of antibodies to *Neospora caninum* in dogs from Amazon, Brazil. Vet Parasitol 2003; 115(1): 71-74. PMid:12860070. [http://dx.doi.org/10.1016/S0304-4017\(03\)00131-6](http://dx.doi.org/10.1016/S0304-4017(03)00131-6).

Cañón-Franco WA, Yai LEO, Souza SLP, Santos LC, Farias NAR, Ruas J, et al. Detection of antibodies to *Neospora caninum* in two species of wild canids, *Lycalopex gymnocercus* and *Cerdocyon thous* from Brazil. Vet Parasitol 2004; 123(3-4): 275-277. PMid:15325054. <http://dx.doi.org/10.1016/j.vetpar.2004.06.004>.

Cardoso JMS, Amaku M, Araújo AJUS, Gennari SM. A longitudinal study of *Neospora caninum* infection on three dairy farms in Brazil. Vet Parasitol 2012a; 187(3-4): 553-557. PMid:22309800. <http://dx.doi.org/10.1016/j.vetpar.2012.01.019>.

Cardoso JMS, Amaku M, Araújo AJUS, Gennari SM. *Neospora caninum*: analysis of reproductive parameters in dairy herds in Brazil. Braz J Vet Res Anim Sci 2012b; 49(6): 459-464. <http://dx.doi.org/10.11606/issn.1678-4456.v49i6p459-464>.

Cardoso JMS, Nishi SM, Funada MR, Amaku M, Guimarães JS Jr, Gennari SM. Antibody dynamics during gestation in cows naturally infected with *Neospora caninum* from four dairy herds in Brazil. Braz J Vet Res Anim Sci 2009; 46(5): 395-399. <http://dx.doi.org/10.11606/issn.1678-4456.bjvras.2009.26789>.

Cavalcante GT, Monteiro RM, Soares RM, Nishi SM, Alves Neto AF, Esmerin PO, et al. Shedding of *Neospora caninum* oocysts by dogs fed different tissues from naturally infected cattle. Vet Parasitol 2011; 179(1-3): 220-223. PMid:21450407. <http://dx.doi.org/10.1016/j.vetpar.2011.02.026>.

Cavalcante GT, Soares RM, Nishi SM, Hagen SCF, Vannucchi CI, Maiorka PC, et al. Experimental infection with *Neospora caninum* in pregnant bitches. Rev Bras Parasitol Vet 2012; 21(3): 232-236. PMid:23070432. <http://dx.doi.org/10.1590/S1984-29612012000300010>.

Chahan B, Gaturaga I, Huang X, Liao M, Fukumoto S, Hirata H, et al. Serodiagnosis of *Neospora caninum* infection in cattle by enzyme-linked immunosorbent assay with recombinant truncated NCsAG1. Vet Parasitol 2003; 118(3-4): 177-185. PMid:14729165. <http://dx.doi.org/10.1016/j.vetpar.2003.10.010>.

Chiebao DP, Valadas SYOB, Minervino AHH, Castro V, Romaldini AHCN, Calhau AS, et al. Variables associated with infections of cattle by *Brucella abortus*, *Leptospira* spp. and *Neospora* spp. in Amazon Region in Brazil. Transbound Emerg Dis 2015; 62(5): e30-e36. PMid:26302373. <http://dx.doi.org/10.1111/tbed.12201>.

Chryssafidis AL, Cantón G, Chianini F, Innes EA, Madureira EH, Gennari SM. Pathogenicity of Nc-Bahia and Nc-1 strains of *Neospora caninum* in experimentally infected cows and buffaloes in early pregnancy. Parasitol Res 2014; 113(4): 1521-1528. PMid:24562816. <http://dx.doi.org/10.1007/s00436-014-3796-x>.

- Chryssafidis AL, Cantón G, Chianini E, Innes EA, Madureira EH, Soares RM, et al. Abortion and foetal lesions induced by *Neospora caninum* in experimentally infected water buffalos (*Bubalus bubalis*). *Parasitol Res* 2015; 114(1): 193-199. PMid:25324135. <http://dx.doi.org/10.1007/s00436-014-4178-0>.
- Chryssafidis AL, Soares RM, Rodrigues AAR, Carvalho NAT, Gennari SM. Evidence of congenital transmission of *Neospora caninum* in naturally infected water buffalo (*Bubalus bubalis*) fetus from Brazil. *Parasitol Res* 2011; 108(3): 741-743. PMid:21181191. <http://dx.doi.org/10.1007/s00436-010-2214-2>.
- Coelho WMD, Amarante AFT, Apolinário JC, Coelho NMD, Lima VMF, Perri SHV, et al. Seroepidemiology of *Toxoplasma gondii*, *Neospora caninum*, and *Leishmania* spp. infections and risk factors for cats from Brazil. *Parasitol Res* 2011; 109(4): 1009-1013. PMid:21626423. <http://dx.doi.org/10.1007/s00436-011-2461-x>.
- Constantino C, Pellizzaro M, Paula EFE, Vieira TSWJ, Brandão APD, Ferreira F, et al. Serosurvey for *Leishmania* spp., *Toxoplasma gondii*, *Trypanosoma cruzi* and *Neospora caninum* in neighborhood dogs in Curitiba-Paraná, Brazil. *Rev Bras Parasitol Vet* 2016; 25(4): 504-510. PMid:27925057. <http://dx.doi.org/10.1590/s1984-29612016062>.
- Corbellini LG, Colodel EM, Driemeier D. Granulomatous encephalitis in a neurologically impaired goat kid associated with degeneration of *Neospora caninum* tissue cysts. *J Vet Diagn Invest* 2001; 13(5): 416-419. PMid:11580064. <http://dx.doi.org/10.1177/104063870101300509>.
- Corbellini LG, Driemeier D, Cruz C, Dias MM. Aborto bovino por *Neospora caninum* no Rio Grande do Sul. *Cienc Rural* 2000; 30(5): 863-868. <http://dx.doi.org/10.1590/S0103-84782000000500021>.
- Corbellini LG, Driemeier D, Cruz CFE, Gondim LFP, Wald V. Neosporosis as a cause of abortion in dairy cattle in Rio Grande do Sul, southern Brazil. *Vet Parasitol* 2002; 103(3): 195-202. PMid:11750112. [http://dx.doi.org/10.1016/S0304-4017\(01\)00600-8](http://dx.doi.org/10.1016/S0304-4017(01)00600-8).
- Corbellini LG, Smith DR, Pescador CA, Schmitz M, Correa A, Steffen DJ, et al. Herd-level risk factors for *Neospora caninum* seroprevalence in dairy farms in southern Brazil. *Prev Vet Med* 2006; 74(2-3): 130-141. PMid:16343669. <http://dx.doi.org/10.1016/j.prevetmed.2005.11.004>.
- Costa GHN, Cabral DD, Varandas NP, Sobral EA, Borges FA, Castagnoli KC. Frequênciade anticorpos anti-*Neospora caninum* e anti-*Toxoplasma gondii* em soros de bovinos pertencentes aos estados de São Paulo e de Minas Gerais. *Semina Cienc Agrar* 2001; 22(1): 61-66. <http://dx.doi.org/10.5433/1679-0359.2001v22n1p61>.
- Costa HF, Stachissini AVM, Langoni H, Padovani CR, Gennari SM, Modolo JR. Reproductive failures associated with antibodies against caprine arthritis-encephalitis virus, *Toxoplasma gondii* and *Neospora caninum* in goats in the state of São Paulo, Brazil. *Braz J Vet Res Anim Sci* 2012; 49(1): 67-72. <http://dx.doi.org/10.11606/issn.2318-3659.v49i1p67-72>.
- Costa KS, Santos SL, Uzeda RS, Pinheiro AM, Almeida MAO, Araújo FR, et al. Chickens (*Gallus domesticus*) are natural intermediate hosts of *Neospora caninum*. *Int J Parasitol* 2008; 38(2): 157-159. PMid:18054356. <http://dx.doi.org/10.1016/j.ijpara.2007.10.008>.
- Cringoli G, Rinaldi L, Capuano F, Baldi L, Veneziano V, Capelli G. Serological survey of *Neospora caninum* and *Leishmania infantum* co-infection in dogs. *Vet Parasitol* 2002; 106(4): 307-313. PMid:12079736. [http://dx.doi.org/10.1016/S0304-4017\(02\)00114-0](http://dx.doi.org/10.1016/S0304-4017(02)00114-0).
- Cunha NA Fo, Lucas AS, Pappen FG, Ragozo AMA, Gennari SM, Lucia Jr T, et al. Fatores de risco e prevalência de anticorpos anti-*Neospora caninum* em cães urbanos e rurais do Rio Grande do Sul, Brasil. *Rev Bras Parasitol Vet* 2008; 17(Suppl 1): 301-306. PMid:20059865.
- Dubey JP, Hemphill A, Calero-Bernal R, Schares G. *Neosporosis in animals*. Boca Raton: CRC Press; 2017.
- Dubey JP, Kerber CE, Granstrom DE. Serologic prevalence of *Sarcocystis neurona*, *Toxoplasma gondii*, and *Neospora caninum* in horses in Brazil. *J Am Vet Med Assoc* 1999; 215(7): 970-972. PMid:10511862.
- Dubey JP, Lindsay DS, Hill D, Romand S, Thulliez P, Kwok OCH, et al. Prevalence of antibodies to *Neospora caninum* and *Sarcocystis neurona* in sera of domestic cats from Brazil. *J Parasitol* 2002; 88(6): 1251-1252. PMid:12537122. [http://dx.doi.org/10.1645/0022-3395\(2002\)088\[1251:POATNC\]2.0.CO;2](http://dx.doi.org/10.1645/0022-3395(2002)088[1251:POATNC]2.0.CO;2).
- Faria EB, Cavalcanti EF, Medeiros ES, Pinheiro JW Jr, Azevedo SS, Athayde AC, et al. Risk factors associated with *Neospora caninum* seropositivity in sheep from the State of Alagoas, in the northeast region of Brazil. *J Parasitol* 2010; 96(1): 197-199. PMid:19799489. <http://dx.doi.org/10.1645/GE-2176.1>.
- Faria EB, Gennari SM, Pena HFJ, Athayde ACR, Silva MLCR, Azevedo SS. Prevalence of anti-*Toxoplasma gondii* and anti-*Neospora caninum* antibodies in goats slaughtered in the public slaughterhouse of Patos city, Paraíba State, Northeast region of Brazil. *Vet Parasitol* 2007; 149(1-2): 126-129. PMid:17706359. <http://dx.doi.org/10.1016/j.vetpar.2007.07.009>.
- Fávero JF, Silva AS, Campigotto G, Machado G, Barros LD, Garcia JL, et al. Risk factors for *Neospora caninum* infection in dairy cattle and their possible cause-effect relation for disease. *Microb Pathog* 2017; 110: 202-207. PMid:28666842. <http://dx.doi.org/10.1016/j.micpath.2017.06.042>.
- Fehlberg HF, Maciel BM, Albuquerque GR. Identification and discrimination of *Toxoplasma gondii*, *Sarcocystis* spp., *Neospora* spp., and *Cryptosporidium* spp. by high-resolution melting analysis. *PLoS One* 2017; 12(3): e0174168. PMid:28346485. <http://dx.doi.org/10.1371/journal.pone.0174168>.
- Feitosa TF, Vilela VLR, Dantas ES, Souto DVO, Pena HFJ, Athayde ACR, et al. *Toxoplasma gondii* and *Neospora caninum* in domestic cats from the Brazilian semi-arid: seroprevalence and risk factors. *Arg Bras Med Vet Zootec* 2014a; 66(4): 1060-1066. <http://dx.doi.org/10.1590/1678-6696>.
- Feitosa TF, Vilela VLR, Melo LRB, Almeida JL No, Souto DVO, Morais DF, et al. *Toxoplasma gondii* and *Neospora caninum* in slaughtered pigs from Northeast, Brazil. *Vet Parasitol* 2014b; 202(3-4): 305-309. PMid:24703253. <http://dx.doi.org/10.1016/j.vetpar.2014.03.015>.
- Fernandes BCTM, Gennari SM, Souza SLP, Carvalho JM, Oliveira WG, Cury MC. Prevalence of anti-*Neospora caninum* antibodies in dogs from urban, periurban and rural areas of the city of Uberlândia, Minas Gerais - Brazil. *Vet Parasitol* 2004; 123(1-2): 33-40. PMid:15265569. <http://dx.doi.org/10.1016/j.vetpar.2004.05.016>.
- Ferreira MST, Vogel FSF, Sangioni LA, Weber A, Bräunig P, Vaz MAB, et al. Oral infection of neonate gerbils by *Neospora caninum* tachyzoites. *Cienc Rural* 2016; 46(4): 654-659. <http://dx.doi.org/10.1590/0103-8478cr20150475>.
- Figliuolo LPC, Kasai N, Ragozo AMA, Paula VSO, Dias RA, Souza SLP, et al. Prevalence of anti-*Toxoplasma gondii* and anti-*Neospora caninum* antibodies in ovine from São Paulo State, Brazil. *Vet Parasitol* 2004a; 123(3-4): 161-166. PMid:15325042. <http://dx.doi.org/10.1016/j.vetpar.2004.06.006>.
- Figliuolo LPC, Rodrigues AAR, Viana RB, Aguiar DM, Kasai N, Gennari SM. Prevalence of anti-*Toxoplasma gondii* and anti-*Neospora caninum* antibodies in goat from São Paulo State, Brazil. *Small Rumin Res* 2004b; 55(1-3): 29-32. <http://dx.doi.org/10.1016/j.smallrumres.2003.12.013>.

- Figueredo LA, Dantas-Torres F, de Faria EB, Gondim LFP, Simões-Mattos L, Brandão-Filho SP, et al. Occurrence of antibodies to *Neospora caninum* and *Toxoplasma gondii* in dogs from Pernambuco, Northeast Brazil. *Vet Parasitol* 2008; 157(1-2): 9-13. PMid:18723288. <http://dx.doi.org/10.1016/j.vetpar.2008.07.009>.
- Filho PCGA, Oliveira JMB, Andrade MR, Silva JG, Kim PCP, Almeida JC, et al. Incidence and vertical transmission rate of *Neospora caninum* in sheep. *Comp Immunol Microbiol Infect Dis* 2017; 52: 19-22. PMid:28673457. <http://dx.doi.org/10.1016/j.cimid.2017.05.006>.
- Fujii TU, Kasai N, Nishi SM, Dubey JP, Gennari SM. Seroprevalence of *Neospora caninum* in female water buffaloes (*Bubalus bubalis*) from the southeastern region of Brazil. *Vet Parasitol* 2001a; 99(4): 331-334. PMid:11511420. [http://dx.doi.org/10.1016/S0304-4017\(01\)00474-5](http://dx.doi.org/10.1016/S0304-4017(01)00474-5).
- Fujii TU, Kasai N, Vasconcellos SA, Richtzenhain LJ, Cortez A, Souza SLP, et al. Anticorpos anti-*Neospora caninum* e contra outros agentes de abortamentos em búfalas da região do Vale do Ribeira, São Paulo, Brasil. *Arg Inst Biol (Sao Paulo)* 2001b; 68(2): 5-9.
- Furuta PI, Mineo TWP, Carrasco AOT, Godoy GS, Pinto AA, Machado RZ. *Neospora caninum* infection in birds: experimental infections in chicken and embryonated eggs. *Parasitology* 2007; 134(Pt14): 1931-1939. PMid:17686190. <http://dx.doi.org/10.1017/S0031182007003344>.
- Galvão CMMQ, Rezende-Gondim MM, Chaves ACR, Schares G, Ribas JRL, Gondim LFP. Brazilian donkeys (*Equus asinus*) have a low exposure to *Neospora* spp. *Rev Bras Parasitol Vet* 2015; 24(3): 340-344. PMid:26444065. <http://dx.doi.org/10.1590/S1984-29612015057>.
- García-Melo DP, Regidor-Cerrillo J, Ortega-Mora LM, Collantes-Fernández E, Oliveira VSF, Oliveira MAP, et al. Isolation and biological characterisation of a new isolate of *Neospora caninum* from an asymptomatic calf in Brazil. *Acta Parasitol* 2009; 54(2): 180-185. <http://dx.doi.org/10.2478/s11686-009-0018-2>.
- Gennari SM, Cañón-Franco WA, Feitosa MM, Ikeda FA, Lima VMF, Amaku M.. Presence of anti-*Neospora caninum* and *Toxoplasma gondii* antibodies in dogs with visceral leishmaniosis from the region of Araçatuba, São Paulo, Brazil. *Braz J Vet Res Anim Sci* 2006; 43(5): 613-619. <http://dx.doi.org/10.11606/issn.1678-4456.bjvras.2006.26569>.
- Genari SM, Pena HFJ, Lindsay DS, Lopes MG, Soares HS, Cabral AD, et al. Prevalence of antibodies against *Neospora* spp. and *Sarcocystis neurona* in donkeys from northeastern Brazil. *Rev Bras Parasitol Vet* 2016; 25(1): 109-111. PMid:26982557. <http://dx.doi.org/10.1590/S1984-29612016003>.
- Genari SM, Rodrigues AAR, Viana RB, Cardoso EC. Occurrence of anti-*Neospora caninum* antibodies in water buffaloes (*Bubalus bubalis*) from the northern region of Brazil. *Vet Parasitol* 2005; 134(1-2): 169-171. PMid:16051440. <http://dx.doi.org/10.1016/j.vetpar.2005.05.064>.
- Genari SM, Yai LEO, D'Áuria SNR, Cardoso SMS, Kwok OCH, Jenkins MC, et al. Occurrence of *Neospora caninum* antibodies in sera from dogs of the city of São Paulo, Brazil. *Vet Parasitol* 2002; 106(2): 177-179. PMid:12031819. [http://dx.doi.org/10.1016/S0304-4017\(02\)00052-3](http://dx.doi.org/10.1016/S0304-4017(02)00052-3).
- Giraldi JH, Bracarense APFRL, Vidotto O, Tudury EA, Navarro IT, Batista TN. Sorologia e histopatologia de *Toxoplasma gondii* e *Neospora caninum* em cães portadores de distúrbios neurológicos. *Semina Cienc Agrar* 2002; 23(1): 9-14. <http://dx.doi.org/10.5433/1679-0359.2002v23n1p9>.
- Gonçalves IN, Uzêda RS, Lacerda GA, Moreira RRN, Araújo FR, Oliveira RHM, et al. Molecular frequency and isolation of cyst-forming coccidia from free ranging chickens in Bahia State, Brazil. *Vet Parasitol* 2012; 190(1-2): 74-79. PMid:22673105. <http://dx.doi.org/10.1016/j.vetpar.2012.05.007>.
- Gondim LFP, Laski P, Gao L, McAllister MM. Variation of the internal transcribed spacer 1 sequence within individual strains and among different strains of *Neospora caninum*. *J Parasitol* 2004; 90(1): 119-122. PMid:15040677. <http://dx.doi.org/10.1645/GE-134R>.
- Gondim LFP, Pinheiro AM, Almeida MAO. Frequência de anticorpos anti-*Neospora caninum* em búfalos (*Bubalus bubalis*) criados no estado da Bahia. *Rev Bras Saúde Prod Anim* 2007; 8(2): 92-96.
- Gondim LFP, Pinheiro AM, Santos POM, Jesus EEV, Ribeiro MB, Fernandes HS, et al. Isolation of *Neospora caninum* from the brain of a naturally infected dog, and production of encysted bradyzoites in gerbils. *Vet Parasitol* 2001; 101(1): 1-7. PMid:11587828. [http://dx.doi.org/10.1016/S0304-4017\(01\)00493-9](http://dx.doi.org/10.1016/S0304-4017(01)00493-9).
- Gondim LFP, Sartor IF, Hasegawa M, Yamane I. Seroprevalence of *Neospora caninum* in dairy cattle in Bahia, Brazil. *Vet Parasitol* 1999a; 86(1): 71-75. PMid:10489205. [http://dx.doi.org/10.1016/S0304-4017\(99\)00129-6](http://dx.doi.org/10.1016/S0304-4017(99)00129-6).
- Gondim LFP, Sartor IF, Monteiro LA Jr, Haritani M. *Neospora caninum* infection in an aborted bovine foetus in Brazil. *N Z Vet J* 1999b; 47(1): 35. PMid:16032066. <http://dx.doi.org/10.1080/00480169.1999.36106>.
- Gondim LSQ, Abe-Sandes K, Uzêda RS, Silva MSA, Santos SL, Mota RA, et al. *Toxoplasma gondii* and *Neospora caninum* in sparrows (*Passer domesticus*) in the Northeast of Brazil. *Vet Parasitol* 2010; 168(1-2): 121-124. PMid:19879051. <http://dx.doi.org/10.1016/j.vetpar.2009.09.055>.
- Greca H, Silva AV, Langoni H. Associação entre a presença de anticorpos anti-*Leishmania* sp. e anti-*Neospora caninum* cães de Bauru, SP. *Arg Bras Med Vet Zootec* 2010; 62(1): 224-227. <http://dx.doi.org/10.1590/S0102-09352010000100032>.
- Guedes MHP, Guimarães AM, Rocha CMBM, Hirsch C. Frequência de anticorpos anti-*Neospora caninum* em vacas e fetos provenientes de municípios do sul de Minas Gerais. *Rev Bras Parasitol Vet* 2008; 17(4): 189-194. PMid:19265576. <http://dx.doi.org/10.1590/S1984-29612008000400004>.
- Guimarães AM, Rocha CMBM, Oliveira TMFS, Rosado IR, Moraes LG, Santos RRD. Fatores associados à soropositividade para *Babesia*, *Toxoplasma*, *Neospora* e *Leishmania* em cães atendidos em nove clínicas veterinárias do município de Lavras, MG. *Rev Bras Parasitol Vet* 2009; 18(Suppl 1): 49-53. PMid:20040191. <http://dx.doi.org/10.4322/rbpv.018e1009>.
- Guimarães JS Jr, Souza SLP, Bergamaschi DP, Genari SM. Prevalence of *Neospora caninum* antibodies and factors associated with their presence in dairy cattle of the north of Paraná state, Brazil. *Vet Parasitol* 2004; 124(1-2): 1-8. PMid:15350656. <http://dx.doi.org/10.1016/j.vetpar.2004.07.002>.
- Hasegawa MY, Sartor IF, Canavessi AMO, Pinckney RD. Ocorrência de anticorpos anti-*Neospora caninum* em bovinos de corte e em cães rurais da região de Avaré, Estado de São Paulo, Brasil. *Semina Cienc Agrar* 2004; 25(1): 45-50.
- Hein HE, Machado G, Miranda ICS, Costa EF, Pellegrini DCP, Driemeier D, et al. Neosporose bovina: avaliação da transmissão vertical e fração atribuível de aborto em uma população de bovinos no Estado do Rio Grande do Sul. *Pesq Vet Bras* 2012; 32(5): 396-400. <http://dx.doi.org/10.1590/S0100-736X2012000500006>.
- Higa AC, Machado RZ, Tinucci-Costa M, Domingues LM, Malheiros EB. Evaluation of cross-reactivity of *Toxoplasma gondii* and *Neospora caninum* antigens in dogs sera. *Rev Bras Parasitol Vet* 2000; 9(2): 91-95.
- Hoane JS, Genari SM, Dubey JP, Ribeiro MG, Borges AS, Yai LEO, et al. Prevalence of *Sarcocystis neurona* and *Neospora* spp. infection in horses from Brazil based on presence of serum antibodies to parasite surface

- antigen. *Vet Parasitol* 2006; 136(2): 155-159. PMid:16310955. <http://dx.doi.org/10.1016/j.vetpar.2005.10.023>.
- Hoffmann Kormann DCS, Locatelli-Dittrich R, Richartz RRTB, Antunes J, Dittrich JR, Patrício LFL. Soroprevalência e cinética mensal de anticorpos anti-*Neospora* sp. em éguas gestantes. *Rev Bras Parasitol Vet* 2008;17(Suppl 1): 335-338. PMid:20059873.
- Jesus EEV, Santos POM, Barbosa MVF, Pinheiro AM, Gondim LFP, Guimarães JE, et al. Frequência de anticorpos anti-*Neospora caninum* em cães nos municípios de Salvador e Lauro de Freitas, Estado da Bahia - Brasil. *Braz J Vet Res Anim Sci* 2006; 43(1): 5-10. <http://dx.doi.org/10.11606/issn.1678-4456.bjvras.2006.26511>.
- Klauck V, Machado G, Pazinato R, Radavelli WM, Santos DS, Berwaguer JC, et al. Relation between *Neospora caninum* and abortion in dairy cows: Risk factors and pathogenesis of disease. *Microb Pathog* 2016; 92: 46-49. PMid:26747583. <http://dx.doi.org/10.1016/j.micpath.2015.12.015>.
- Langoni H, Fornazari F, Silva RC, Monti ET, Villa FB. Prevalence of antibodies against *Toxoplasma gondii* and *Neospora caninum* in dogs. *Braz J Microbiol* 2014; 44(4): 1327-1330. PMid:24688530. <http://dx.doi.org/10.1590/S1517-83822013000400043>.
- Langoni H, Greca H Jr, Guimarães FF, Ullmann LS, Gaio FC, Uehara RS, et al. Serological profile of *Toxoplasma gondii* and *Neospora caninum* infection in commercial sheep from São Paulo State, Brazil. *Vet Parasitol* 2011; 177(1-2): 50-54. PMid:21256676. <http://dx.doi.org/10.1016/j.vetpar.2010.11.024>.
- Langoni H, Matteucci G, Medici B, Camossi LG, Richini-Pereira VB, Silva RC. Detection and molecular analysis of *Toxoplasma gondii* and *Neospora caninum* from dogs with neurological disorders. *Rev Soc Bras Med Trop* 2012; 45(3): 365-368. PMid:22760137. <http://dx.doi.org/10.1590/S0037-86822012000300016>.
- Laskoski LM, Muraro LS, Dittrich RL, Abreu RA, Koch MO, Silva FT, et al. Occurrence of anti-*Neospora caninum* and anti-*Toxoplasma gondii* antibodies in horses in the Pantanal of Mato Grosso, Brazil. *Semina Cienc Agrar* 2015; 36(2): 895-900. <http://dx.doi.org/10.5433/1679-0359.2015v36n2p895>.
- Lima JTR, Ahid SMM, Barrêto RA Jr, Pena HFJ, Dias RA, Gennari SM. Prevalência de anticorpos anti-*Toxoplasma gondii* e anti-*Neospora caninum* em rebanhos caprinos do município de Mossoró, Rio Grande do Norte. *Braz J Vet Res Anim Sci* 2008; 45(2): 81-86. <http://dx.doi.org/10.11606/issn.1678-4456.bjvras.2008.26703>.
- Lobato J, Silva DAO, Mineo TWP, Amaral JDHF, Segundo GRS, Costa-Cruz JM, et al. Detection of immunoglobulin G antibodies to *Neospora caninum* in humans: high seropositivity rates in patients who are infected by human immunodeficiency virus or have neurological disorders. *Clin Vaccine Immunol* 2006; 13(1): 84-89. PMid:16426004. <http://dx.doi.org/10.1128/CVI.13.1.84-89.2006>.
- Locatelli-Dittrich R, Dittrich JR, Richartz RRTB, Gasino-Joineau ME, Antunes J, Pinckney RD, et al. Investigation of *Neospora* sp. and *Toxoplasma gondii* antibodies in mares and in precolostral foals from Paraná State, Southern Brazil. *Vet Parasitol* 2006; 135(3-4): 215-221. PMid:16289863. <http://dx.doi.org/10.1016/j.vetpar.2005.10.010>.
- Locatelli-Dittrich R, Machado PC Jr, Fridlund-Plugge N, Richartz RRTB, Montiani-Ferreira F, Patrício LFL, et al. Determinação e correlação de anticorpos anti-*Neospora caninum* em bovinos e cães do Paraná, Brasil. *Rev Bras Parasitol Vet* 2008;17(Suppl 1): 191-196. PMid:20059847.
- Locatelli-Dittrich R, Soccol VT, Richartz RRTB, Gasino-Joineau ME, Vinne R, Pinckney RD. Serological diagnosis of neosporosis in a herd of dairy cattle in southern Brazil. *J Parasitol* 2001; 87(6): 1493-1494. PMid:11780849. [http://dx.doi.org/10.1645/0022-3395\(2001\)087\[1493:SDONIA\]2.0.CO;2](http://dx.doi.org/10.1645/0022-3395(2001)087[1493:SDONIA]2.0.CO;2).
- Locatelli-Dittrich R, Soccol VT, Richartz RRTB, Gasino-Joineau ME, van der Vinne R, Pinckney RD. Isolamento de *Neospora caninum* de feto bovino de rebanho leiteiro no Paraná. *Rev Bras Parasitol Vet* 2004; 13(3): 103-109.
- Locatelli-Dittrich R, Sousa RS, Leite LC, Thomaz-Soccol V, Richartz RRTB, Joineau MEG, et al. Isolation of *Neospora caninum* from a blind calf in Paraná, southern Brazil. *Vet Rec* 2003; 153(12): 366-367. PMid:14533770. <http://dx.doi.org/10.1136/vr.153.12.366>.
- Lopes MG, Mendonça IL, Fortes KP, Amaku M, Pena HFJ, Gennari SM. Presence of antibodies against *Toxoplasma gondii*, *Neospora caninum* and *Leishmania infantum* in dogs from Piauí. *Rev Bras Parasitol Vet* 2011; 20(2): 111-114. PMid:21722484. <http://dx.doi.org/10.1590/S1984-29612011000200004>.
- Machado GP, Kikuti M, Langoni H, Paes AC. Seroprevalence and risk factors associated with neosporosis in sheep and dogs from farms. *Vet Parasitol* 2011; 182(2-4): 356-358. PMid:21676548. <http://dx.doi.org/10.1016/j.vetpar.2011.05.021>.
- Mann TR, Cadore GC, Camillo G, Vogel FSF, Schmidt C, Andrade CM. Canine cutaneous neosporosis in Brazil. *Vet Dermatol* 2016; 27(3): 195-197. PMid:26949109. <http://dx.doi.org/10.1111/vde.12294>.
- Marques FAC, Headley AS, Figueiredo-Pereira V, Taroda A, Barros LD, Cunha IAL, et al. *Neospora caninum*: evaluation of vertical transmission in slaughtered beef cows (*Bos indicus*). *Parasitol Res* 2011; 108(4): 1015-1019. PMid:21063729. <http://dx.doi.org/10.1007/s00436-010-2146-x>.
- Martins AA, Zamprogna TO, Lucas TM, Cunha IAL, Garcia JL, Silva AV. Frequency and risk factors for infection by *Neospora caninum* in dairy farms of Umuarama, PR, Brazil. *Arq Ciênc Vet Zool* 2012; 15(2): 137-142.
- Martins NÉX, Freschi CR, Baptista F, Machado RZ, Freitas FLC, Almeida KS. Ocorrência de anticorpos anti-*Neospora caninum* em vacas lactantes do município de Araguaína, estado do Tocantins, Brasil. *Rev Patol Trop* 2011; 40(3): 231-238.
- Mello RC, Andreotti R, Barros JC, Tomich RGP, Mello AKM, Campolim AI, et al. Levantamento epidemiológico de *Neospora caninum* em bovinos de assentamentos rurais em Corumbá, MS. *Rev Bras Parasitol Vet* 2008;17(Suppl 1): 311-316. PMid:20059867.
- Melo CB, Leite RC, Leite FSC, Leite RC. Serological surveillance on South American wild canids for *Neospora caninum*. *Arg Bras Med Vet Zootec* 2002; 54(4): 444-447. <http://dx.doi.org/10.1590/S0102-09352002000400018>.
- Melo CB, Leite RC, Lobato ZIP, Leite RC. Infection by *Neospora caninum* associated with bovine herpesvirus 1 and bovine viral diarrhea virus in cattle from Minas Gerais State, Brazil. *Vet Parasitol* 2004; 119(2-3): 97-105. PMid:14746970. <http://dx.doi.org/10.1016/j.vetpar.2003.12.002>.
- Melo CB, Leite RC, Souza GN, Leite RC. Frequência de infecção por *Neospora caninum* em dois diferentes sistemas de produção de leite e fatores predisponentes à infecção em bovinos em Minas Gerais. *Rev Bras Parasitol Vet* 2001; 10(2): 67-74.
- Melo DPG, Silva AC, Ortega-Mora LM, Bastos SA, Boaventura CM. Prevalência de anticorpos anti-*Neospora caninum* em bovinos das microrregiões de Goiânia e Anápolis, Goiás, Brasil. *Rev Bras Parasitol Vet* 2006; 15(3): 105-109. PMid:16978474.
- Meneses ID, Andrade MR, Uzeda RS, Bittencourt MV, Lindsay DS, Gondim LFP. Frequency of antibodies against *Sarcocystis neurona* and *Neospora caninum* in domestic cats in the state of Bahia, Brazil. *Rev Bras*

*Parasitol Vet* 2014; 23(4): 526-529. PMid:25517534. <http://dx.doi.org/10.1590/S1984-29612014080>.

Mesquita LP, Nogueira CI, Costa RC, Orlando DR, Bruhn FR, Lopes PFR, et al. Antibody kinetics in goats and conceptuses naturally infected with *Neospora caninum*. *Vet Parasitol* 2013; 196(3-4): 327-333. PMid:23537945. <http://dx.doi.org/10.1016/j.vetpar.2013.03.002>.

Mineo TWP, Alenius S, Näslund K, Montassier HJ, Björkman C. Distribution of antibodies against *Neospora caninum*, BVDV and BHV-1 among cows in Brazilian dairy herds with reproductive disorders. *Rev Bras Parasitol Vet* 2006; 15(4): 188-192. PMid:17196123.

Mineo TWP, Carrasco AOT, Marciano JA, Werther K, Pinto AA, Machado RZ. Pigeons (*Columba livia*) are a suitable experimental model for *Neospora caninum* infection in birds. *Vet Parasitol* 2009; 159(2): 149-153. PMid:19027237. <http://dx.doi.org/10.1016/j.vetpar.2008.10.024>.

Mineo TWP, Carrasco AOT, Raso TF, Werther K, Pinto AA, Machado RZ. Survey for natural *Neospora caninum* infection in wild and captive birds. *Vet Parasitol* 2011; 182(2-4): 352-355. PMid:21680099. <http://dx.doi.org/10.1016/j.vetpar.2011.05.022>.

Mineo TWP, Silva DAO, Costa GHN, von Ancken ACB, Kasper LH, Souza MA, et al. Detection of IgG antibodies to *Neospora caninum* and *Toxoplasma gondii* in dogs examined in a veterinary hospital from Brazil. *Vet Parasitol* 2001; 98(4): 239-245. PMid:11423182. [http://dx.doi.org/10.1016/S0304-4017\(01\)00441-1](http://dx.doi.org/10.1016/S0304-4017(01)00441-1).

Mineo TWP, Silva DAO, Näslund K, Björkman C, Uggla A, Mineo JR. *Toxoplasma gondii* and *Neospora caninum* serological status of different canine populations from Uberlândia, Minas Gerais. *Arq Bras Med Vet Zootec* 2004; 56(3): 414-417. <http://dx.doi.org/10.1590/S0102-09352004000300022>.

Minervino AHH, Cassinelli ABM, Lima JTR, Soares HS, Malheiros AF, Marcili A, et al. Prevalence of anti-*Neospora caninum* and anti-*Toxoplasma gondii* antibodies in dogs from two different indigenous communities in the Brazilian Amazon Region. *J Parasitol* 2012; 98(6): 1276-1278. PMid:22551468. <http://dx.doi.org/10.1645/GE-3151.1>.

Minervino AHH, Ragozo AMA, Monteiro RM, Ortolani EL, Gennari SM. Prevalence of *Neospora caninum* antibodies in cattle from Santarém, Pará, Brazil. *Res Vet Sci* 2008; 84(2): 254-256. PMid:17619028. <http://dx.doi.org/10.1016/j.rvsc.2007.05.003>.

Modolo JR, Stachissini AVM, Gennari SM, Dubey JP, Langoni H, Padovani CR, et al. Freqüência de anticorpos anti-*Neospora caninum* em soros de caprinos do estado de São Paulo e sua relação com o manejo dos animais. *Pesq Vet Bras* 2008; 28(12): 597-600. <http://dx.doi.org/10.1590/S0100-736X2008001200006>.

Moraes CCG, Megid J, Pituco EM, Okuda LH, Del Fava C, Stefano E, et al. Ocorrência de anticorpos anti-*Neospora caninum* em cães da microrregião da Serra de Botucatu, Estado de São Paulo, Brasil. *Rev Bras Parasitol Vet* 2008; 17(1): 1-6. PMid:18554432. <http://dx.doi.org/10.1590/S1984-29612008000100001>.

Moraes LMB, Raimundo JM, Guimarães A, Santos HA, Macedo GL Jr, Massard CL, et al. Occurrence of anti-*Neospora caninum* and anti-*Toxoplasma gondii* IgG antibodies in goats and sheep in western Maranhão, Brazil. *Rev Bras Parasitol Vet* 2011; 20(4): 312-317. PMid:22166386. <http://dx.doi.org/10.1590/S1984-29612011000400010>.

Morikawa VM, Zimpel CK, Paploski IAD, Lara MCCSH, Villalobos EMC, Romaldini AHNC, et al. Occurrences of anti-*Toxoplasma gondii* and anti-*Neospora caninum* antibodies in Barbary sheep at Curitiba zoo, southern Brazil. *Rev Bras Parasitol Vet* 2014; 23(2): 255-259. PMid:25054509. <http://dx.doi.org/10.1590/S1984-296120140434>.

Moura AB, Silva MO, Farias JA, Vieira-Neto A, Souza AP, Sartor AA, et al. *Neospora* spp. antibodies in horses from two geographical regions of the state of Santa Catarina, Brazil. *Rev Bras Parasitol Vet* 2013; 22(4): 597-601. PMid:24473888. <http://dx.doi.org/10.1590/S1984-29612013000400023>.

Munhoz AD, Amaral TF, Gonçalves LR, Moraes VMB, Machado RZ. *Gallus gallus domesticus* are resistant to infection with *Neospora caninum* tachyzoites of the NC-1 strain. *Vet Parasitol* 2014; 206(3-4): 123-128. PMid:25468016. <http://dx.doi.org/10.1016/j.vetpar.2014.10.009>.

Munhoz AD, Flausino W, Silva RT, Almeida CRR, Lopes CWG. Distribuição de anticorpos contra *Neospora caninum* em vacas leiteiras dos municípios de Resende e Rio Claro, Estado do Rio de Janeiro, Brasil. *Rev Bras Parasitol Vet* 2006; 15(3): 101-104. PMid:16978473.

Munhoz AD, Mineo TWP, Alessi AC, Lopes CWG, Machado RZ. Assessment of experimental infection for dogs using *Gallus gallus* chorioallantoic membranes inoculated with *Neospora caninum*. *Rev Bras Parasitol Vet* 2013; 22(4): 565-570. PMid:24473883. <http://dx.doi.org/10.1590/S1984-29612013000400018>.

Munhoz AD, Pereira MJS, Flausino W, Lopes CWG. *Neospora caninum* seropositivity in cattle breeds in the South Fluminense Paraíba Valley, state of Rio de Janeiro. *Pesq Vet Bras* 2009; 29(1): 29-32. <http://dx.doi.org/10.1590/S0100-736X2009000100004>.

Muradian V, Ferreira LR, Lopes EG, Esmerini PO, Pena HFJ, Soares RM, et al. A survey of *Neospora caninum* and *Toxoplasma gondii* infection in urban rodents from Brazil. *J Parasitol* 2012; 98(1): 128-134. PMid:21790367. <http://dx.doi.org/10.1645/GE-2817.1>.

Nascimento COM, Silva MLCR, Kim PCP, Gomes AAB, Gomes ALV, Maia RCC, et al. Occurrence of *Neospora caninum* and *Toxoplasma gondii* DNA in brain tissue from hoary foxes (*Pseudalopex vetulus*) in Brazil. *Acta Trop* 2015; 146: 60-65. PMid:25746974. <http://dx.doi.org/10.1016/j.actatropica.2015.02.016>.

Nascimento EE, Sammi AS, Santos JR, Nino BSL, Bogado ALG, Taroda A, et al. Anti-*Neospora caninum* antibody detection and vertical transmission rate in pregnant zebu beef cows (*Bos indicus*): *Neospora caninum* in pregnant beef cows (*Bos indicus*). *Comp Immunol Microbiol Infect Dis* 2014; 37(4): 267-270. PMid:25193073. <http://dx.doi.org/10.1016/j.cimid.2014.08.002>.

Nogueira CI, Mesquita LP, Abreu CC, Nakagaki KYR, Seixas JN, Bezerra PS, et al. Risk factors associated with seroprevalence of *Neospora caninum* in dogs from urban and rural areas of milk and coffee production in Minas Gerais state, Brazil. *Epidemiol Infect* 2013; 141(11): 2286-2293. PMid:23419686. <http://dx.doi.org/10.1017/S0950268813000162>.

Norlander E. *Seroprevalence of Toxoplasma gondii and Neospora spp. in equids from three municipalities in Pardá, Brazil* [Dissertation]. Uppsala: Swedish University of Agricultural Sciences; 2014.

Ogawa L, Freire RL, Vidotto O, Gondim LFP, Navarro IT. Occurrence of antibodies to *Neospora caninum* and *Toxoplasma gondii* in dairy cattle from the northern region of the Paraná State, Brazil. *Arq Bras Med Vet Zootec* 2005; 57(3): 312-316. <http://dx.doi.org/10.1590/S0102-09352005000300006>.

Oliveira JM, Matos MFC, Oshiro LM, Andreotti R. Prevalence of anti-*Neospora caninum* antibodies in dogs in the urban area of Campo Grande, MS, Brazil. *Rev Bras Parasitol Vet* 2004; 13(4): 155-158.

Oliveira S, Soares RM, Aizawa J, Soares HS, Chiebilo DP, Ortega-Mora LM, et al. Isolation and biological and molecular characterization of *Neospora caninum* (NC-SP1) from a naturally infected adult asymptomatic cattle (*Bos taurus*) in the state of São Paulo, Brazil. *Parasitology* 2017; 144(6): 707-711. PMid:28073388. <http://dx.doi.org/10.1017/S0031182016002481>.

- Oliveira UV, Magalhães VCS, Almeida CP, Santos IA, Mora DA, Macêdo LS, et al. Quails are resistant to infection with *Neospora caninum* tachyzoites. *Vet Parasitol* 2013; 198(1-2): 209-213. PMid:24041486. <http://dx.doi.org/10.1016/j.vetpar.2013.08.009>.
- Onuma SSM, Melo ALT, Kantek DLZ, Crawshaw-Junior PG, Morato RG, May-Júnior JA, et al. Exposure of free-living jaguars to *Toxoplasma gondii*, *Neospora caninum* and *Sarcocystis neurona* in the Brazilian Pantanal. *Rev Bras Parasitol Vet* 2014; 23(4): 547-553. PMid:25517539. <http://dx.doi.org/10.1590/S1984-29612014077>.
- Oshiro LM, Matos MFC, Oliveira JM, Monteiro LARC, Andreotti R. Prevalence of anti-*Neospora caninum* antibodies in cattle from the state of Mato Grosso do Sul, Brazil. *Rev Bras Parasitol Vet* 2007; 16(3): 133-138. PMid:18078599. <http://dx.doi.org/10.1590/S1984-29612007000300004>.
- Oshiro LM, Motta-Castro ARC, Freitas SZ, Cunha RC, Dittrich RL, Meirelles ACF, et al. *Neospora caninum* and *Toxoplasma gondii* serodiagnosis in human immunodeficiency virus carriers. *Rev Soc Bras Med Trop* 2015; 48(5): 568-572. PMid:26516966. <http://dx.doi.org/10.1590/0037-8682-0151-2015>.
- Paiz LM, Silva RC, Menozzi BD, Langoni H. Antibodies to *Neospora caninum* in sheep from slaughterhouses in the state of São Paulo, Brazil. *Rev Bras Parasitol Vet* 2015; 24(1): 95-100. PMid:25909261. <http://dx.doi.org/10.1590/S1984-296120150009>.
- Paulan SC, Lins AGS, Tenório MS, Silva DT, Pena HFJ, Machado RZ, et al. Seroprevalence rates of antibodies against *Leishmania infantum* and other protozoan and rickettsial parasites in dogs. *Rev Bras Parasitol Vet* 2013; 22(1): 162-166. PMid:24252965. <http://dx.doi.org/10.1590/S1984-29612013000100031>.
- Paz GF, Leite RC, Rocha MA. Associação entre sorologia para *Neospora caninum* e taxa de prenhez em vacas receptoras de embriões. *Arg Bras Med Vet Zootec* 2007; 59(5): 1323-1325. <http://dx.doi.org/10.1590/S0102-09352007000500034>.
- Pena HFJ, Soares RM, Ragozo AMA, Monteiro RM, Yai LEO, Nishi SM, et al. Isolation and molecular detection of *Neospora caninum* from naturally infected sheep from Brazil. *Vet Parasitol* 2007; 147(1-2): 61-66. PMid:17451882. <http://dx.doi.org/10.1016/j.vetpar.2007.03.002>.
- Pescador CA, Corbellini LG, Oliveira EC, Raymundo DL, Driemeier D. Histopathological and immunohistochemical aspects of *Neospora caninum* diagnosis in bovine aborted fetuses. *Vet Parasitol* 2007; 150(1-2): 159-163. PMid:17904290. <http://dx.doi.org/10.1016/j.vetpar.2007.08.028>.
- Pessoa GA, Martini AP, Trentin JM, Dalcin VC, Leonardi CEP, Vogel FSF, et al. Impact of spontaneous *Neospora caninum* infection on pregnancy loss and subsequent pregnancy in grazing lactating dairy cows. *Theriogenology* 2016; 85(3): 519-527. PMid:26542136. <http://dx.doi.org/10.1016/j.theriogenology.2015.09.034>.
- Pivoto FL, Macêdo AG Jr, Silva MV, Ferreira FB, Silva DAO, Pompermayer E, et al. Serological status of mares in parturition and the levels of antibodies (IgG) against protozoan family Sarcocystidae from their pre colostral foals. *Vet Parasitol* 2014; 199(1-2): 107-111. PMid:24183649. <http://dx.doi.org/10.1016/j.vetpar.2013.10.001>.
- Plugge NF, Ferreira FM, Richartz RRTB, Siqueira A, Dittrich RL. Occurrence of antibodies against *Neospora caninum* and/or *Toxoplasma gondii* in dogs with neurological signs. *Rev Bras Parasitol Vet* 2011; 20(3): 202-206. PMid:21961748. <http://dx.doi.org/10.1590/S1984-29612011000300004>.
- Plugge NF, Montiani-Ferreira F, Richartz RRTB, Dal Pizzol J, Machado PC Jr, Patrício LFL, et al. Frequency of antibodies against *Neospora caninum* in stray and domiciled dogs from urban, periurban and rural areas from Paraná State, Southern Brazil. *Rev Bras Parasitol Vet* 2008; 17(4): 222-226. PMid:19265582. <http://dx.doi.org/10.1590/S1984-29612008000400010>.
- Portella LP, Cadore GC, Sangioni LA, Pellegrini LFV, Fighera R, Ramos F, et al. Antibodies against Apicomplexa protozoa and absence sarcocysts in heart tissues from horses in southern Brazil. *Rev Bras Parasitol Vet* 2017; 26(1): 100-103. PMid:28327879. <http://dx.doi.org/10.1590/S1984-29612016068>.
- Ragozo AMA, Paula VSO, Souza SLP, Bergamaschi DP, Gennari SM. Ocorrência de anticorpos anti-*Neospora caninum* em soros bovinos procedentes de seis estados brasileiros. *Rev Bras Parasitol Vet* 2003; 12(1): 33-37.
- Raimundo JM, Guimarães A, Moraes LMB, Santos LA, Nepomuceno LL, Barbosa SM, et al. *Toxoplasma gondii* and *Neospora caninum* in dogs from the state of Tocantins: serology and associated factors. *Rev Bras Parasitol Vet* 2015; 24(4): 475-481. PMid:26689184. <http://dx.doi.org/10.1590/S1984-29612015068>.
- Ramos IAS, Silva RJ, Maciel TA, Silva JABA, Fidelis OL, Soares PC, et al. Assessment of transplacental transmission of *Neospora caninum* in dairy cattle in the Agreste region of Pernambuco. *Rev Bras Parasitol Vet* 2016; 25(4): 516-522. PMid:27737368. <http://dx.doi.org/10.1590/S1984-29612016055>.
- Regidor-Cerrillo J, Díez-Fuertes F, García-Culebras A, Moore DP, González-Warleta M, Cuevas C, et al. Genetic diversity and geographic population structure of bovine *Neospora caninum* determined by microsatellite genotyping analysis. *PLoS One* 2013; 8(8): e72678. PMid:23940816. <http://dx.doi.org/10.1371/journal.pone.0072678>.
- Rezende-Gondim MM, Silva AV, Schares G, Gondim LFP. In contrast to *Toxoplasma gondii*, *Neospora caninum* tachyzoites did not sustain multiplication *in vitro* at increased incubation temperatures. *Vet Parasitol* 2017; 234: 19-24. PMid:28115178. <http://dx.doi.org/10.1016/j.vetpar.2016.12.013>.
- Rodrigues AAR, Gennari SM, Aguiar DM, Sreekumar C, Hill DE, Miska KB, et al. Shedding of *Neospora caninum* oocysts by dogs fed tissues from naturally infected water buffaloes (*Bubalus bubalis*) from Brazil. *Vet Parasitol* 2004; 124(3-4): 139-150. PMid:15381294. <http://dx.doi.org/10.1016/j.vetpar.2004.07.007>.
- Romanelli PR, Freire RL, Vidotto O, Marana ERM, Ogawa L, Paula VSO, et al. Prevalence of *Neospora caninum* and *Toxoplasma gondii* in sheep and dogs from Guarapuava farms, Paraná State, Brazil. *Res Vet Sci* 2007; 82(2): 202-207. PMid:17266999. <http://dx.doi.org/10.1016/j.rvsc.2006.04.001>.
- Rossi GF, Cabral DD, Ribeiro DP, Pajuaba ACAM, Corrêa RR, Moreira RQ, et al. Evaluation of *Toxoplasma gondii* and *Neospora caninum* infections in sheep from Uberlândia, Minas Gerais State, Brazil, by different serological methods. *Vet Parasitol* 2011; 175(3-4): 252-259. PMid:21075529. <http://dx.doi.org/10.1016/j.vetpar.2010.10.017>.
- Salaberry SRS, Okuda LH, Nassar AFC, Castro JR, Lima-Ribeiro AMC. Prevalence of *Neospora caninum* antibodies in sheep flocks of Uberlândia county, MG. *Rev Bras Parasitol Vet* 2010; 19(3): 148-151. PMid:20943017. <http://dx.doi.org/10.1590/S1984-29612010000300004>.
- Santos APME, Navarro IT, Freire RL, Vidotto O, Bracarense APFRL. *Neospora caninum* in dairy cattle in Paraná State, Brazil: histological and immunohistochemical analysis in fetuses. *Semina Cienc Agrar* 2005; 26(4): 559-562. <http://dx.doi.org/10.5433/1679-0359.2005v26n4p559>.
- Sartor IF, Garcia A Fo, Vianna LC, Pituco EM, Dal Pai V, Sartor R. Ocorrência de anticorpos anti-*Neospora caninum* em bovinos leiteiros e

de corte da região de Presidente Prudente, SP. *Arg Inst Biol (Sao Paulo)* 2005; 72(4): 413-418.

Sartor IF, Hasegawa MY, Canavessi AMO, Pinckney RD. Ocorrência de anticorpos de *Neospora caninum* em vacas leiteiras avaliados pelos métodos ELISA e RIFI no município de Avaré, SP. *Semina Cienc Agrar* 2003; 24(1): 3-10. <http://dx.doi.org/10.5433/1679-0359.2003v24n1p3>.

Seabra NM, Pereira VF, Kuwassaki MV, Benassi JC, Oliveira TMFS. *Toxoplasma gondii*, *Neospora caninum*, *Leishmania* spp. serology and *Leishmania* spp. PCR in dogs from Pirassununga, SP. *Rev Bras Parasitol Vet* 2015; 24(4): 454-458. PMid:26689180. <http://dx.doi.org/10.1590/S1984-29612015046>.

Sicupira PML, Magalhães VCS, Galvão GS, Pereira MJS, Gondim LFP, Munhoz AD. Factors associated with infection by *Neospora caninum* in dogs in Brazil. *Vet Parasitol* 2012; 185(2-4): 305-308. PMid:22015062. <http://dx.doi.org/10.1016/j.vetpar.2011.09.029>.

Silva DAO, Lobato J, Mineo TWP, Mineo JR. Evaluation of serological tests for the diagnosis of *Neospora caninum* infection in dogs: optimization of cut off titers and inhibition studies of cross-reactivity with *Toxoplasma gondii*. *Vet Parasitol* 2007; 143(3-4): 234-244. PMid:16973287. <http://dx.doi.org/10.1016/j.vetpar.2006.08.028>.

Silva JB, Nicolino RR, Fagundes GM, Bomjardim HA, Reis ASB, Lima DHS, et al. Serological survey of *Neospora caninum* and *Toxoplasma gondii* in cattle (*Bos indicus*) and water buffaloes (*Bubalus bubalis*) in ten provinces of Brazil. *Comp Immunol Microbiol Infect Dis* 2017; 52: 30-35. PMid:28673459. <http://dx.doi.org/10.1016/j.cimid.2017.05.005>.

Silva JB, Santos PN, Castro GNS, Fonseca AH, Barbosa JD. Prevalence survey of selected bovine pathogens in water buffaloes in the north region of Brazil. *J Parasitol Res* 2014; 2014: 603484. PMid:24563780. <http://dx.doi.org/10.1155/2014/603484>.

Silva MIS, Almeida MAO, Mota RA, Pinheiro-Junior JW, Rabelo SSA. Fatores de riscos associados à infecção por *Neospora caninum* em matrizes bovinas leiteiras em Pernambuco. *Cienc Anim Bras* 2008; 9(2): 455-461.

Silva MSA, Uzêda RS, Costa KS, Santos SL, Macedo ACC, Abe-Sandes K, et al. Detection of *Hammondia heydorni* and related coccidia (*Neospora caninum* and *Toxoplasma gondii*) in goats slaughtered in Bahia, Brazil. *Vet Parasitol* 2009; 162(1-2): 156-159. PMid:19278786. <http://dx.doi.org/10.1016/j.vetpar.2009.02.007>.

Soares HS, Ahid SMM, Bezerra ACDS, Pena HFJ, Dias RA, Gennari SM. Prevalence of anti-*Toxoplasma gondii* and anti-*Neospora caninum* antibodies in sheep from Mossoró, Rio Grande do Norte, Brazil. *Vet Parasitol* 2009a; 160(3-4): 211-214. PMid:19091473. <http://dx.doi.org/10.1016/j.vetpar.2008.10.102>.

Soares HS, Ramos VN, Osava CF, Oliveira S, Szabó MPJ, Piovezan U, et al. Occurrence of antibodies against *Neospora caninum* in wild pigs (*Sus scrofa*) in the Pantanal, Mato Grosso do Sul, Brazil. *Braz J Vet Res Anim Sci* 2016; 53(1): 112-116. <http://dx.doi.org/10.11606/issn.1678-4456.v53i1p112-116>.

Soares RM, Cortez LRPB, Gennari SM, Sercundes MK, Keid LB, Pena HFJ. Crab-eating fox (*Cerdocyon thous*), a South American canid, as a definitive host for *Hammondia heydorni*. *Vet Parasitol* 2009b; 162(1-2): 46-50. PMid:19303215. <http://dx.doi.org/10.1016/j.vetpar.2009.02.003>.

Sousa KCM, Herrera HM, Domingos IH, Campos JBV, Santos IMC, Neves HH, et al. Serological detection of *Toxoplasma gondii*, *Leishmania infantum* and *Neospora caninum* in cats from an area endemic for leishmaniasis in Brazil. *Rev Bras Parasitol Vet* 2014; 23(4): 449-455. PMid:25517522. <http://dx.doi.org/10.1590/S1984-29612014078>.

Sousa ME, Porto WJN, Albuquerque PPE, Souza OL No, Pinheiro JW, Mota RA. Seroprevalence of antibodies to *Neospora caninum* in dogs in the state of Alagoas, Brazil. *Rev Bras Parasitol Vet* 2012; 21(3): 287-290. PMid:23070441. <http://dx.doi.org/10.1590/S1984-29612012000300019>.

Souza LM, Nascimento AA, Furuta PI, Basso LMS, Silveira DM, Costa AJ. Detecção de anticorpos contra *Neospora caninum* e *Toxoplasma gondii* em soros de bubalinos (*Bubalus bubalis*) no Estado de São Paulo, Brasil. *Semina Cienc Agrar* 2001; 22(1): 39-48. <http://dx.doi.org/10.5433/1679-0359.2001v22n1p39>.

Souza SLP, Guimarães JS, Ferreira F, Dubey JP, Gennari SM. Prevalence of *Neospora caninum* antibodies in dogs from dairy cattle farms in Paraná, Brazil. *J Parasitol* 2002; 88(2): 408-409. PMid:12054023. [http://dx.doi.org/10.1645/0022-3395\(2002\)088\[0408:PONCAI\]2.0.CO;2](http://dx.doi.org/10.1645/0022-3395(2002)088[0408:PONCAI]2.0.CO;2).

Stelmann UJP, Ullmann LS, Langoni H, Amorim RM. Equine neosporosis: search for antibodies in cerebrospinal fluid and sera from animals with history of ataxia. *Rev Bras Med Vet* 2011; 33(2): 99-102.

Taques IIGG, Barbosa TR, Martini AC, Pitchenin LC, Braga ÍA, Melo ALT, et al. Molecular assessment of the transplacental transmission of *Toxoplasma gondii*, *Neospora caninum*, *Brucella canis* and *Ehrlichia canis* in dogs. *Comp Immunol Microbiol Infect Dis* 2016; 49: 47-50. PMid:27865263. <http://dx.doi.org/10.1016/j.cimid.2016.09.002>.

Teixeira WC, Silva MIS, Pereira JG, Pinheiro AM, Almeida MAO, Gondim LFP. Freqüência de cães reagentes para *Neospora caninum* em São Luís, Maranhão. *Arg Bras Med Vet Zootec* 2006; 58(4): 685-687. <http://dx.doi.org/10.1590/S0102-09352006000400038>.

Teixeira WC, Uzêda RS, Gondim LFP, Silva MIS, Pereira HM, Alves LC, et al. Prevalência de anticorpos anti-*Neospora caninum* (Apicomplexa: Sarcocystidae) em bovinos leiteiros de propriedades rurais em três microrregiões no estado do Maranhão. *Pesq Vet Bras* 2010; 30(9): 729-734. <http://dx.doi.org/10.1590/S0100-736X20100090004>.

Tembue AASM, Ramos RAN, Sousa TR, Albuquerque AR, Costa AJ, Meunier IMJ, et al. Serological survey of *Neospora caninum* in small ruminants from Pernambuco State, Brazil. *Rev Bras Parasitol Vet* 2011; 20(3): 246-248. PMid:21961757. <http://dx.doi.org/10.1590/S1984-29612011000300013>.

Tiemann JCH, Rodrigues AAR, Souza SLP, Duarte JMB, Gennari SM. Occurrence of anti-*Neospora caninum* antibodies in Brazilian cervids kept in captivity. *Vet Parasitol* 2005a; 129(3-4): 341-343. PMid:15845290. <http://dx.doi.org/10.1016/j.vetpar.2004.12.016>.

Tiemann JCH, Souza SLP, Rodrigues AAR, Duarte JMB, Gennari SM. Environmental effect on the occurrence of anti-*Neospora caninum* antibodies in pampas-deer (*Ozotoceros bezoarticus*). *Vet Parasitol* 2005b; 134(1-2): 73-76. PMid:16112811. <http://dx.doi.org/10.1016/j.vetpar.2005.07.015>.

Topazio JP, Weber A, Camillo G, Vogel FF, Machado G, Ribeiro A, et al. Seroprevalence and risk factors for *Neospora caninum* in goats in Santa Catarina state, Brazil. *Rev Bras Parasitol Vet* 2014; 23(3): 360-366. PMid:25271457. <http://dx.doi.org/10.1590/S1984-29612014062>.

Toscan G, Vogel FSF, Cadore GC, Cezar AS, Sangioni LA, Pereira RCF, et al. Occurrence of antibodies anti-*Neospora* spp. in cart horses and Crioula breed horses from Rio Grande do Sul State. *Arg Bras Med Vet Zootec* 2011; 63(1): 258-261. <http://dx.doi.org/10.1590/S0102-09352011000100038>.

Truppel JH, Montiani-Ferreira F, Lange RR, Vilani RGOC, Reifur L, Boerger W, et al. Detection of *Neospora caninum* DNA in capybaras and

- phylogenetic analysis. *Parasitol Int* 2010; 59(3): 376-379. PMid:20470895. <http://dx.doi.org/10.1016/j.parint.2010.05.001>.
- Ueno TEH, Gonçalves VSP, Heinemann MB, Dilli TLB, Akimoto BM, Souza SLP, et al. Prevalence of *Toxoplasma gondii* and *Neospora caninum* infections in sheep from Federal District, central region of Brazil. *Trop Anim Health Prod* 2009; 41(4): 547-552. PMid:18726165. <http://dx.doi.org/10.1007/s11250-008-9220-8>.
- Uzêda RS, Costa KS, Santos SL, Pinheiro AM, Almeida MAO, McAllister MM, et al. Loss of infectivity of *Neospora caninum* oocysts maintained for a prolonged time. *Korean J Parasitol* 2007a; 45(4): 295-299. PMid:18165712. <http://dx.doi.org/10.3347/kjp.2007.45.4.295>.
- Uzêda RS, Pinheiro AM, Fernández SY, Ayres MCC, Gondim LFP, Almeida MAO. Seroprevalence of *Neospora caninum* in dairy goats from Bahia, Brazil. *Small Rumin Res* 2007b; 70(2-3): 257-259. <http://dx.doi.org/10.1016/j.smallrumres.2006.04.003>.
- Valadas S, Gennari SM, Yai LEO, Rosypal AC, Lindsay DS. Prevalence of antibodies to *Trypanosoma cruzi*, *Leishmania infantum*, *Encephalitozoon cuniculi*, *Sarcocystis neurona*, and *Neospora caninum* in capybara, *Hydrochoerus hydrochaeris*, from São Paulo State, Brazil. *J Parasitol* 2010a; 96(3): 521-524. PMid:20020808. <http://dx.doi.org/10.1645/GE-2368.1>.
- Valadas S, Minervino AHH, Lima VMF, Soares RM, Ortolani EL, Gennari SM. Occurrence of antibodies anti-*Neospora caninum*, anti-*Toxoplasma gondii*, and anti-*Leishmania chagasi* in serum of dogs from Pará State, Amazon, Brazil. *Parasitol Res* 2010b; 107(2): 453-457. PMid:20445991. <http://dx.doi.org/10.1007/s00436-010-1890-2>.
- Varandas NP, Rached PA, Costa GHN, Souza LM, Castagnoli KC, Costa AJ. Freqüência de anticorpos anti-*Neospora caninum* e anti-*Toxoplasma gondii* em cães da região nordeste do Estado de São Paulo. Correlação com neuropatias. *Semina Cienc Agrar* 2001; 22(1): 105-111. <http://dx.doi.org/10.5433/1679-0359.2001v22n1p105>.
- Varaschin MS, Hirsch C, Wouters F, Nakagaki KY, Guimarães AM, Santos DS, et al. Congenital neosporosis in goats from the State of Minas Gerais, Brazil. *Korean J Parasitol* 2012; 50(1): 63-67. PMid:22451736. <http://dx.doi.org/10.3347/kjp.2012.50.1.63>.
- Viana RB, Del Fava C, Moura ACB, Cardoso EC, Araújo CV, Monteiro BM, et al. Ocorrência de anticorpos anti-*Neospora caninum*, *Brucella* sp. e *Leptospira* spp. em búfalos (*Bubalus bubalis*) criados na Amazônia. *Arg Inst Biol (Sao Paulo)* 2009; 76(3): 453-457.
- Villalobos EMC, Furman KE, Lara MCCSH, Cunha EMS, Finger MA, Busch APB, et al. Detection of *Neospora* sp. antibodies in cart horses from urban areas of Curitiba, Southern Brazil. *Rev Bras Parasitol Vet* 2012; 21(1): 68-70. PMid:22534949. <http://dx.doi.org/10.1590/S1984-29612012000100014>.
- Villalobos EMC, Ueno TEH, Souza SLP, Cunha EMS, Lara MCCSH, Gennari SM, et al. Association between the presence of serum antibodies against *Neospora* spp. and fetal loss in equines. *Vet Parasitol* 2006; 142(3-4): 372-375. PMid:16962708. <http://dx.doi.org/10.1016/j.vetpar.2006.07.016>.
- Vitaliano SN, Silva DAO, Mineo TWP, Ferreira RA, Bevilacqua E, Mineo JR. Seroprevalence of *Toxoplasma gondii* and *Neospora caninum* in captive maned wolves (*Chrysocyon brachyurus*) from southeastern and midwestern regions of Brazil. *Vet Parasitol* 2004; 122(4): 253-260. PMid:15262003. <http://dx.doi.org/10.1016/j.vetpar.2004.04.004>.
- Vogel FSF, Arenhart S, Bauermann FV. Anticorpos anti-*Neospora caninum* em bovinos, ovinos e bubalinos no Estado do Rio Grande do Sul. *Cienc Rural* 2006; 36(6): 1948-1951. <http://dx.doi.org/10.1590/S0103-84782006000600048>.
- Yai LEO, Cañón-Franco WA, Geraldi VC, Summa MEL, Camargo MCGO, Dubey JP, et al. Seroprevalence of *Neospora caninum* and *Toxoplasma gondii* antibodies in the South American opossum (*Didelphis marsupialis*) from the city of São Paulo, Brazil. *J Parasitol* 2003; 89(4): 870-871. PMid:14533710. <http://dx.doi.org/10.1645/GE-83R>.
- Yai LEO, Ragozo AMA, Cañón-Franco WA, Dubey JP, Gennari SM. Occurrence of *Neospora caninum* antibodies in capybaras (*Hydrochoerus hydrochaeris*) from São Paulo State, Brazil. *J Parasitol* 2008; 94(3): 766. PMid:18605804. <http://dx.doi.org/10.1645/GE-1468.1>.
- Zimpel CK, Grazziotin AL, Barros IR Fo, Guimarães AMS, Santos LC, Moraes W, et al. Occurrence of antibodies anti-*Toxoplasma gondii*, *Neospora caninum* and *Leptospira interrogans* in a captive deer herd in Southern Brazil. *Rev Bras Parasitol Vet* 2015; 24(4): 482-487. PMid:26689185. <http://dx.doi.org/10.1590/S1984-29612015065>.