

# New morphological data and molecular diagnostic of *Henneguya friderici* (Myxozoa: Myxobolidae), a parasite of *Leporinus friderici* (Osteichthyes: Anostomidae) from southeastern Brazil

Novos dados morfológicos e diagnóstico molecular de *Henneguya friderici* (Myxozoa: Myxobolidae), parasito de *Leporinus friderici* (Osteichthyes: Anostomidae) do sudeste do Brasil

Letícia Poblete Vidal<sup>1</sup>; José Luis Luque<sup>2\*</sup>

<sup>1</sup> Programa de Pós-graduação em Ciências Veterinárias, Universidade Federal Rural do Rio de Janeiro – UFRRJ, Seropédica, RJ, Brasil

<sup>2</sup> Departamento de Parasitologia Animal, Universidade Federal Rural do Rio de Janeiro – UFRRJ, Seropédica, RJ, Brasil

Received December 20, 2016

Accepted February 16, 2017

## Abstract

The myxozoan *Henneguya friderici* is a parasite of the gills, intestine, kidney and liver of *Leporinus friderici*, a characiform fish belonging to the family Anostomidae. Forty-two specimens of *L. friderici* that had been caught in the Mogi Guaçu River, state of São Paulo, were studied. Elongated white plasmodia were found in the gill filaments of 10 host specimens (24%). The mature spores had an ellipsoidal body with polar capsules of equal size and caudal length greater than body length. This study also described 18S rDNA sequencing of *H. friderici* infecting the gill filaments. This produced a sequence of 1050 bp that demonstrated significant genetic differences with previously described species of *Henneguya*. Similarity analysis using sequences from species that clustered closest to those produced by this study showed that the species with greatest genetic similarity to *H. friderici* was *H. leporinicola*, with 94% similarity.

**Keywords:** Myxosporea, Characiformes, 18S rDNA, phylogeny.

## Resumo

O myxozoa *Henneguya friderici* é um parasito encontrado nas brânquias, fígado, intestino e rins de *Leporinus friderici*, (Characiformes: Anostomidae). Foram capturados e examinados quarenta e dois espécimes de *L. friderici* oriundos do Rio Mogi Guaçu, estado de São Paulo. Cistos alongados e brancos foram encontrados nos filamentos branquiais de 10 (24%) hospedeiros. Os esporos maduros apresentaram o corpo alongado com as cápsulas polares em tamanhos iguais e o comprimento caudal maior do que o comprimento corporal. Com isso, o presente trabalho, descreve o sequenciamento de 1050 pb do gene 18S rDNA de *H. friderici* infectando os filamentos branquiais, o que demonstrou diferenças genéticas significativas em comparação com espécies previamente descritas de *Henneguya*. A análise de similaridade utilizando as sequencias de espécies que se agruparam mais próximas às produzidas por este estudo mostrou que a espécie com maior semelhança genética com *H. friderici* foi *H. leporinicola*, com 94% de similaridade.

**Paravras-chave:** Myxosporea, Characiformes, 18S rDNA, filogenia.

## Introduction

The diversity of known myxozoans has grown greatly since the early work of Kudo (1919). Around 2.200 species have now been described (LOM & DYKOVÁ, 2006) and these represent around 18% of cnidarian species diversity, as far as is currently known (OKAMURA et al., 2015). *Henneguya* Thélohan, 1892,

is one of the most diverse genera of Myxosporea and currently includes more than 200 known and described species (LOM & DYKOVÁ, 2006). This widespread genus includes typical coelozoic and histozoic species and predominantly infects marine and freshwater fish (EIRAS & ADRIANO, 2012).

Currently, more than 44 species of *Henneguya* are known to infect South America fish (EIRAS, 2002; EIRAS & ADRIANO, 2012; CARRIERO et al., 2013; NALDONI et al., 2014).

\*Corresponding author: José Luis Luque. Departamento de Parasitologia Animal, Universidade Federal Rural do Rio de Janeiro – UFRRJ, CP 74540, CEP 23851-970, Seropédica, RJ, Brasil. e-mail: luqueufri@gmail.com

Of these, around 28 have been found to infect fish species of the order Characiformes.

Identification of the species in this genus, like those in other genera of myxozoans, is based almost exclusively on spore morphology. In the class Myxosporea, morphology has been the main criterion for classification of species (KUDO, 1933; MOLNÁR, 1994). In fact, this method has always failed to identify highly similar species that are found in the same infection site and host and which only have subtle differences in spore structures (YE et al., 2012). Fortunately, this problem has been solved through molecular approaches (SMOTHERS et al., 1994; ANDREE et al., 1999; HOLZER et al., 2004). 18S rDNA is the molecular marker that has most commonly been used for detection, identification and phylogenetic analysis on myxozoans (HOLZER et al., 2006). The difficulties of relying on spore morphology for species identification have led authors to recommend that SSU rDNA sequencing should be included when new species are described (ANDREE et al., 1999; KENT et al., 2001; LOM & DYKOVÁ, 2006).

*Leporinus friderici* (Bloch, 1794) is a characiform fish belonging to the family Anostomidae that is, popularly known in Brazil as “piau”. It is widely distributed in the Amazon and Paraguay river basins (FROESE & PAULY, 2016). Among the species of *Henneguya*, only *Henneguya friderici* (CASAL et al., 2003) has been reported from *L. friderici*.

*Henneguya friderici* was found infecting the gills, intestine, kidney and liver of “piau” from an estuarine region of the Amazon River, in the state of Pará, Brazil. Relative organelle preservation occurred in the liver tissue and, in some cases, development of the parasite caused gradual and generalized degeneration in the intestine, gills and kidney (CASAL et al., 2003).

The present paper supplements the original description of *H. friderici* with new data on morphology and 18S rDNA sequencing on samples from gill filaments of *L. friderici* from the Mogi Guaçú River, state of São Paulo, Brazil. The new data support the original diagnosis by Casal et al. (2003).

## Materials and Methods

Forty-two specimens of *L. friderici* were caught by local fishermen with nets and hooks in the Mogi Guaçú River near Pirassununga, state of São Paulo, Brazil ( $21^{\circ}55'36''$  S;  $47^{\circ}22'6''$  W), between January 2014 and January 2016. Gills extracted from the fish were placed in Petri dishes with tap water and were examined for myxozoans using a dissecting microscope. Infected gill filaments were preserved using two different methods: frozen (for spore measurements) and in 95% ethanol (for DNA analysis).

Parasitological examinations were conducted using standard methods with the aid of an optical microscope (Olympus BX51) with differential interference contrast (DIC). Images were captured using a 3.2 mp UC30 digital camera and were analyzed by means of photomicrography software (Cell<sup>D</sup> 3.4, Olympus Soft Imaging Solutions GmbH, Germany). At least 30 measurements were made for each relevant spore dimension, following the guidelines of Lom & Arthur (1989).

Gill filaments from three hosts were used for DNA extraction by means of the DNeasy Blood & Tissue Kit, following the manufacturer's instructions (QIAGEN Inc., California, USA). The polymerase chain reaction (PCR) was performed as described by Whipps et al. (2015) in 50 µl reaction volumes of the Quick-Load Taq 23 Master Mix (New England Biolabs, Ipswich, Massachusetts, USA), with 0.5 µM of each primer and 3 µl of template DNA. A first round of amplification targeting the small subunit (SUU) rDNA was performed using the primers 18E and 18R (WHIPPS et al., 2003), followed by a second round of PCR with 18E and Myxgen2R (KENT et al., 2000) or with 18R and Myxgen3F (KENT et al., 2000). The amplifications were performed in a C1000<sup>TM</sup> thermal cycler (BioRad Laboratories, Hercules, California, USA) with initial denaturation at 95 °C for 3 min, followed by 35 cycles of 94 °C for 30 s, 56 °C for 45 s and 68 °C for 90 s, and a final extension at 72 °C for 7 min. Product amplification was evaluated by observation on 1% agarose gel, and the remainder of the sample was purified using the E.Z.N.A. Cycle Pure Kit (Omega Bio-Tek, Norcross, Georgia, USA). DNA was quantified using a DNA spectrophotometer (NanoDrop Technologies, Wilmington, Delaware, USA). Sequencing reactions were carried out by means of the ABI BigDye Terminator Cycle Sequencing Ready Reaction Kit version 3.1, using the ABI3730xl Genetic Analyzer (Applied Biosystems, Foster City, California, USA). Contiguous sequences were assembled in Geneious (Geneious version 9, created by Biomatters, available from (<http://www.geneious.com/>) and were deposited in GenBank (Table 1).

Alignments were subjected to maximum likelihood (ML) and Bayesian inference (BI) (rates = invgamma) analyses; additionally, Tamura & Nei (TRN) distance values were performed using Geneious. ML and BI trees were calculated under the TRN + I + G model for the sequences of the rDNA 18S, using PHYML (GUINDON & GASCUEL, 2003) and MrBayes (HUELSENBECK & RONQUIST, 2001) Geneious plug-ins for ML and BI, respectively. These models were selected using jModelTest2 (DARRIBA et al., 2012). Nucleotide frequencies were estimated from the data (A = 0.2824, C = 0.1634, G = 0.2826, T = 0.2715). Six rates of nucleotide substitution were (AC) = 1.0000, (AG) = 3.2212, (AT) = 1.0000, (CG) = 1.0000, (CT) = 6.0419, (GT) = 1.0000; proportion of invariable sites = 0.1460; gamma distribution = 0.3960 estimated with 4 rate categories. ML nodal support was estimated by 1000 nonparametric bootstrap replications. Bayesian posterior probability were determined running the Markov chains (two runs and four chains) for  $4 \times 10^6$  generations, discarding the initial 1/4 of sampled trees (trees sample every  $4 \times 10^3$  generations) as burn in fraction. Phylogenetic trees were rooted using *Ceratonoova shasta* (Noble, 1950) as outgroup based upon previous Myxobolidae phylogenies (ADRIANO et al., 2009; CAPODIFOGLIO et al., 2015; NALDONI et al., 2015).

## Results

*Henneguya friderici* cysts were found in gill filaments from ten piau specimens, i.e. 24% of the total examined. The elongated white plasmodia measured approximately 2 mm in length. The mature spores were ellipsoid in frontal view and the valves were symmetrical


**Table 1.** List of myxozoan whose sequences were used for analyses and the obtained in the present study.

Parasite	GenBank accession No.	Host	Country	Reference
<i>Ceratomyxa shasta</i>	AF001579	<i>Oncorhynchus mykiss</i>	USA	Bartholomew et al. (1997)
<i>Henneguya adiposa</i>	EU492929	<i>Ictalurus punctatus</i>	USA	Griffin et al. (2009)
<i>Henneguya bulbosus</i>	KM000055	<i>Ictalurus punctatus</i>	USA	Rosser et al. (2014)
<i>Henneguya. cerebralis</i>	JX131380	<i>Thymallus nigrescens</i>	Mongolia	Batueva et al. (2013)
<i>Henneguya. corruscans</i>	JQ654971	<i>Pseudoplatystoma corruscans</i>	Brazil	Adriano et al. (2012)
<i>Henneguya creplini</i>	EU732597	<i>Zingel zingel</i>	Hungary	Eszterbauer et al. (2006)
<i>Henneguya cuniculator</i>	KF732840	<i>Pseudoplatystoma corruscans</i>	Brazil	Naldoni et al. (2014)
<i>Henneguya cutanea</i>	AY676460	<i>Abramis brama</i>	Hungary	Kallert et al. (2005)
<i>Henneguya dogieli</i>	KJ725078	<i>Siniperca chuatsi</i>	China	Unpublished
<i>Henneguya doneci</i>	LC011456	<i>Carassius gibelio</i>	China	Li et al. (2015)
<i>Henneguya doneci</i>	EU344898	<i>Carassius auratus</i>	China	Unpublished
<i>Henneguya doneci</i>	HM146129	<i>Carassius gibelio</i>	China	Ye et al. (2012)
<i>Henneguya doori</i>	HDU37549	<i>Perca fluviatilis</i>	Canada	Siddall et al. (1995)
<i>Henneguya exilis</i>	AF021881	<i>Ictalurus punctatus</i>	USA	Lin et al. (1999)
<i>Henneguya friderici</i>	KY315824	<i>Leporinus friderici</i>	Brazil	Present study
<i>Henneguya gurlei</i>	DQ673465	<i>Ameiurus nebulosus</i>	USA	Iwanowicz et al. (2008)
<i>Henneguya ictaluri</i>	AF195510	<i>Ictalurus punctatus</i>	USA	Pote et al. (2000)
<i>Henneguya jocu</i>	KF264964	<i>Lutjanus jocu</i>	Portugal	Azevedo et al. (2014)
<i>Henneguya leporinicola</i>	KP980550	<i>Leporinus macrocephalus</i>	Brazil	Capodifoglio et al. (2015)
<i>Henneguya lobosa</i>	EU732600	<i>Esox lucius</i>	Germany	Eszterbauer et al. (2006)
<i>Henneguya maculosus</i>	KF296344	<i>Pseudoplatystoma corruscans</i>	Brazil	Carrierio et al. (2013)
<i>Henneguya mississippiensis</i>	KP404438	<i>Ictalurus punctatus</i>	USA	Rosser et al. (2015)
<i>Henneguya multiplasmoidalis</i>	JQ654969	<i>Pseudoplatystoma corruscans</i>	Brazil	Adriano et al. (2012)
<i>Henneguya pellis</i>	FJ468488	<i>Ictalurus punctatus</i>	USA	Griffin et al. (2009)
<i>Henneguya pellucida</i>	KF296352	<i>Piaractus mesopotamicus</i>	Brazil	Carrierio et al. (2013)
<i>Henneguya piaractus</i>	KF597016	<i>Piaractus mesopotamicus</i>	Brazil	Müller et al. (2013)
<i>Henneguya pseudoplatystoma</i>	KP981638	<i>Pseudoplatystoma corruscans</i>	Brazil	Milanin et al. (2015)
<i>Henneguya pseudorhinogobii</i>	AB447994	<i>Rhinogobius</i> sp.	Japan	Kageyama et al. (2009)
<i>Henneguya psorospermica</i>	EU732602	<i>Esox lucius</i>	Germany	Eszterbauer et al. (2006)
<i>Henneguya rhinogobii</i>	AB447992	<i>Rhinogobius</i> sp.	Japan	Kageyama et al. (2009)
<i>Henneguya rotunda</i>	KJ416130	<i>Salminus brasiliensis</i>	Brazil	Moreira et al. (2014a)
<i>Henneguya salminicola</i>	AF031411	<i>Oncorhynchus kisutch</i>	Canada	Hervio et al. (1997)
<i>Henneguya</i> sp.	JQ411297	<i>Oncorhynchus masou masou</i>	Japan	Yokoyama et al. (2012)
<i>Henneguya</i> sp.	KR704889	<i>Cirrhinus mrigala</i>	India	Unpublished
<i>Henneguya</i> sp.	EU732601	<i>Esox lucius</i>	Hungary	Eszterbauer et al. (2006)
<i>Henneguya</i> sp.	EU732599	<i>Perca fluviatilis</i>	Hungary	Eszterbauer et al. (2006)
<i>Henneguya</i> sp.	JQ690355	<i>Carassius auratus</i>	China	Unpublished
<i>Henneguya sutherlandi</i>	EF191200	<i>Ictalurus punctatus</i>	USA	Griffin et al. (2008)
<i>Henneguya visibilis</i>	KC771143	<i>Leporinus obtusidens</i>	Brazil	Moreira et al. (2014b)
<i>Henneguya zikawaiensis</i>	KR020026	<i>Carassius auratus</i>	China	Zhang et al. (2015)
<i>Henneguya zschorkei</i>	HZU13827	<i>Prosopium williamsonii</i>	USA	Smathers et al. (1994)
<i>Henneguya zschorkei</i>	AF378344	<i>Prosopium williamsonii</i>	Canada	Kent et al. (2001)

and convex in lateral view. The polar capsules were elongated and equal in size and occupied a little less than half of the spore body (Figure 1). Spores ( $N = 30$ ) were  $12.8 \pm 2.1$  (7.4-14.8)  $\mu\text{m}$  in length,  $4.4 \pm 0.4$  (3.4-5.2)  $\mu\text{m}$  in width and  $32.8 \pm 2.6$  (2.49-40)  $\mu\text{m}$  in total length. The bifurcated caudal processes were cylindrical, equal in size,  $19.6 \pm 2.2$  (16.1-24.4)  $\mu\text{m}$  in length, and extended behind the spore. Two equal capsules were pyriform, tapering toward their anterior end and occupying nearly half of the spore, and they measured  $5.1 \pm 0.5$  (3.7-5.9)  $\mu\text{m}$  in length and  $1.5 \pm 0.1$  (1.2-1.8)  $\mu\text{m}$  in width (Figure 1). Table 2 provides a comparison

between the data on the spore dimensions, infection sites and host of *H. friderici* obtained in this study and the data from the original descriptions.

The 18S rDNA sequencing on *H. friderici* spores resulted in a sequence containing 1050 bp, which was deposited in the GenBank database under accession number KY315824. This sequence was used for phylogenetic analysis. A BLAST comparison between the sequence obtained and other myxosporean sequences available in GenBank revealed that the 18S rDNA sequence of *H. friderici* had 92% similarity to that of *Henneguya leporinicola* Martins,



**Figure 1.** (a) Mature spore of *Henneguya friderici* parasite of gill filaments of *Leporinus friderici* in frontal view with Nomarski interference contrast; (b) schematic of *Henneguya friderici* myxospore demonstrating the polar capsule, spore capsule, and caudal processes. Scale bar 10 µm.


Souza, Moraes & Moraes, 1999 (KP980550) and 89% similarity to that of *H. bulbosus* Rosser, Griffin, Quiniu, Khoo & Pote, 2014 (KM000055).

Similarity analysis using sequences from species that clustered closest to those produced by the present study showed that the species with greatest genetic similarity to *H. friderici* was *H. leporinicola*, with 94% similarity. The ML and BI phylogenetic tree (Figure 2) showed that *H. friderici* appears as a sister species of *H. leporinicola* in a subclade composed mainly of myxosporean parasites of Characiformes and Esociformes.

## Discussion

*Henneguya friderici* was described by Casal et al. (2003) infecting the gills, intestine, kidney and liver of *L. friderici* in the Amazon River, near Belém, state of Pará, Brazil. Its description was based on morphological and ultrastructural data. This was, in the past, the main method for characterization and identification of myxosporeans (MOLNÁR, 2002). However, Kent et al. (2001) and Lom & Dyková (2006) suggested that amplification of 18S rDNA is fundamental for describing new species of myxosporeans, because of the difficulties of characterizing the spores morphologically.

The present study provided 18S rDNA sequencing on *H. friderici* that was found infecting the gill filaments of host caught in the Mogi Guaçú River in the state of São Paulo. This enabled phylogenetic




**Figure 2.** Maximum Likelihood from phylogenetic analysis of the sequences of 18S rDNA gene of *Henneguya friderici* associated with the closest species indicated by the analysis of Max Score by BLAST of the NCBI platform. First number of nodal support is from maximum likelihood bootstrap (1000 replications), the second number shows Bayesian posterior probability (for  $4 \times 10^6$  generations; burn-in =  $4 \times 10^3$ ). Sample from the present study is in **bold**.

**Table 2.** Comparison of the characteristics of *Henneguya friderici* with similar species.

Species	LS	WS	AL	TL	PCL	PCW	Host	Site	Reference
<i>Henneguya leporincola</i>	7.6(5.5-8.7)	4.2 (3.6-4.9)	21.8 (12.9-32.2)	-	3.0 (2.0-3.6)	1.6 (1.2-2.0)	<i>Leporinus macrocephalus</i>	Gills	Martins et al. (1999)
<i>Henneguya azvedoi</i>	12.0(11-13)	3.2 (3-4)	39.4 (37-40)	56.4 (52-58)	6.3 (6-7)	2.1 (2-3)	<i>Leporinus obtusidens</i>	Gill lamellae	Barassa et al. (2012)
<i>Henneguya canticula</i>	11.3 (11-12)	5.4 (5-6)	3.4 (3-4)	14.7 (14-16)	3.7 (3-4)	1.5	<i>Leporinus lacustris</i>	Gill filament	Eiras et al. (2008)
<i>Henneguya friderici</i>	10.4 (9.6-11.8)	5.7 (4.8-6.6)	23.3 (19.1-28.7)	33.8 (28.7-39.3)	4.9 (4.2-5.9)	2.1 (1.5-2.6)	<i>Leporinus friderici</i>	Gills	Casal et al. (2003)
<i>Henneguya schizodon</i>	13.1 (12-14)	3.3 (3.4)	16.3 (15-17)	28.9 (27-30)	5.4 (5-6)	1.3 (1-1.5)	<i>Schizodon fasciatus</i>	Kidney	Eiras et al. (2004)
<i>Henneguya visibilis</i>	10.8 ± 0.6	3.9 ± 0.2	18 ± 1.2	26.8 ± 1.1	4.9 ± 0.3	1.4 ± 0.1	<i>Leporinus obtusidens</i>	Connective tissue	Moreira et al. (2014b)
<i>Henneguya friderici</i>	12.8 ± 2.1 (7.4-14.8)	4.4 ± 0.4 (3.4-5.2)	19.6 ± 2.2 (16.1-24.4)	32.8 ± 2.6 (24.9-40)	5.1 ± 0.5 (3.7-5.9)	1.5 ± 0.1 (1.2-1.8)	<i>Leporinus friderici</i>	Gill filament	Present study

LS: length of the spore; WS: width of the spore; AL: length of the tail; TL: total length of the spore; PCL: polar capsules length; PCW: polar capsules width.



**Figure 3.** Plasmodia of *Henneguya friderici* infecting the gill filaments of *Leporinus friderici*. Scale bar = 10 mm.

analysis on this parasite. The 18S rDNA gene is used in molecular systematics for determining relationships among myxozoans because it is highly variable between very closely related species (KENT et al., 2001). The morphometric and morphological data obtained in the present study clearly confirmed the identification of the species as *H. friderici*, which was originally described by Casal et al. (2003) (Table 2).

Molnár (2002) divided the formation of gill-located myxosporean plasmodia into three types: (1) lamellar; (2) filamental; and (3) gill arch. Among these, the filamental type is subdivided into four types: (1) vascular; (2) epithelial; (3) intrachondral; and (4) basifilamental. In the present study, the *H. friderici* plasmodia developed on the filamentous epithelium of the gills and deformed the gill filaments (Figure 3).

The prevalence of *H. friderici* in piau was 24%. This was close to the 30% reported by Casal et al. (2003), considering all the infected organs of *L. friderici*. However, in fish from the Mogi Guaçu River, infection was only observed in the gill filaments. Furthermore, these results corroborated data from other studies conducted in South America in which species of *Henneguya* were found at the same infection site (NALDONI et al., 2009, 2014).

These supplementary data on the morphology, 18S rDNA sequencing and phylogeny of *H. friderici* may facilitate accurate diagnoses and better understanding of the phylogenetic relationships of this parasite. Fiala (2006) indicated that host preference is very important and that myxosporean species could group together according to fish host species. Although host geographical origin is particularly important, tissue tropism in myxosporean evolution has also been revealed in numerous phylogenetic studies (ANDREE et al., 1999; KENT et al., 2001; ESZTERBAUER, 2004; FIALA, 2006).

## Acknowledgements

To M.Sc. Júlio Cesar C. de Aguiar (CEPTA/IBAMA) for assistance in the development of the work. Letícia G. P. Vidal was supported by a Doctoral fellowship from CNPq (Conselho Nacional

de Desenvolvimento Científico e Tecnológico do Brasil), and José L. Luque was supported by a Researcher fellowship from CNPq.

## References

- Adriano EA, Arana S, Alves AL, Silva MRM, Ceccarelli PS, Henrique-Silva F, et al. *Myxobolus cordeiroi* n. sp., a parasite of *Zungaro jahu* (Siluriformes: Pimelodiidae) from Brazilian Pantanal: morphology, phylogeny and histopathology. *Vet Parasitol* 2009; 162(3-4): 221-229. PMid:19372007. <http://dx.doi.org/10.1016/j.vetpar.2009.03.030>.
- Adriano EA, Carriero MM, Maia AAM, Silva MRM, Naldoni J, Ceccarelli PS, et al. Phylogenetic and host-parasite relationship analysis of *Henneguya multiplasmodialis* n. sp. infecting *Pseudoplatystoma* spp. in Brazilian Pantanal wetland. *Vet Parasitol* 2012; 185(2-4): 110-120. PMid:22051071. <http://dx.doi.org/10.1016/j.vetpar.2011.10.008>.
- Andree KB, Székely C, Molnár K, Gresoviac SJ, Hedrick RP. Relationships among members of the Genus *Myxobolus* (Myxozoa: Bivalvidea) based on small subunit ribosomal DNA sequence. *J Parasitol* 1999; 85(1): 68-74. PMid:10207366. <http://dx.doi.org/10.2307/3285702>.
- Azevedo C, Rocha S, Matos P, Matos E, Oliveira E, Al-Quraishi S, et al. Morphology and phylogeny of *Henneguya jocu* n. sp. (Myxosporea, Myxobolidae), infecting the gills of the marine fish *Lutjanus jocu*. *Eur J Protistol* 2014; 50(2): 185-193. PMid:24457131. <http://dx.doi.org/10.1016/j.ejop.2013.12.002>.
- Barassa B, Adriano EA, Cordeiro NS, Arana S, Ceccarelli PS. Morphology and host – parasite interaction of *Henneguya azevedoi* n. sp., parasite of gills of *Leporinus obtusidens* from Mogi Guaçu River, Brazil. *Parasitol Res* 2012; 110(2): 887-894. PMid:21842391. <http://dx.doi.org/10.1007/s00436-011-2571-5>.
- Bartholomew JL, Whipple MJ, Stevens DG, Fryer JL. The life cycle of *Ceratomyxa shasta*, a Myxosporean parasite of salmonids, requires a freshwater polychaete as an alternate host. *J Parasitol* 1997; 83(5): 859-868. PMid:9379291. <http://dx.doi.org/10.2307/3284281>.
- Batueva MD, Katokhin AV, Pronina SV, Pronin NM. Supplementary studies and molecular data on *Henneguya cerebralis* Pronin, 1972 (Myxozoa: Myxosporea), a parasite from Kosogol grayling *Thymallus arcticus nigrescens* in Mongolia. *Parasitol Int* 2013; 62(6): 530-534. PMid:23933262. <http://dx.doi.org/10.1016/j.parint.2013.07.014>.
- Capodifoglio KRH, Adriano EA, Silva MRM, Maia AAM. Supplementary data of *Henneguya leporinicola* (Myxozoa, Myxosporea) a parasite of *Leporinus macrocephalus* from fish farms in the state of São Paulo, Brazil. *Acta Parasitol* 2015; 60(3): 451-458. PMid:26204182. <http://dx.doi.org/10.1515/ap-2015-0062>.
- Carriero MM, Adriano EA, Silva MR, Ceccarelli PS, Maia AA. Molecular phylogeny of the *Myxobolus* and *Henneguya* genera with several new South American species. *PLoS One* 2013; 8(9): e73713. PMid:24040037. <http://dx.doi.org/10.1371/journal.pone.0073713>.
- Casal G, Matos E, Azevedo C. Light and electron microscopic study of the myxosporean, *Henneguya friderici* n. sp. from the Amazonian teleostean fish, *Leporinus friderici*. *Parasitol* 2003; 126(4): 313-319. PMid:12741510. <http://dx.doi.org/10.1017/S0031182003002944>.
- Darriba D, Taboada GL, Ramón D, Posada D. jModelTest 2: more models, new heuristics and high-performance computing. *Nat Methods* 2012; 9(8): 772. PMid:22847109. <http://dx.doi.org/10.1038/nmeth.2109>.

- Eiras JC. Synopsis of the species of the genus *Henneguya* Thélohan, 1892 (Myxozoa: Myxosporea: Myxobolidae). *Syst Parasitol* 2002; 52(1): 43-54. PMid:12023561. <http://dx.doi.org/10.1023/A:1015016312195>.
- Eiras JC, Adriano EA. A checklist of new species of *Henneguya* Thélohan, 1892 (Myxozoa: Myxosporea, Myxobolidae) described between 2002 and 2012. *Syst Parasitol* 2012; 83(2): 95-104. PMid:22983797. <http://dx.doi.org/10.1007/s11230-012-9374-7>.
- Eiras JC, Malta JC, Varela A, Pavanelli GC. *Henneguya schizodon* n. sp. (Myxozoa, Myxobolidae), a parasite of the Amazonian Teleost fish *Schizodon fasciatus* (Characiformes, Anostomidae). *Parasite* 2004; 11(2): 169-173. PMid:15224578. <http://dx.doi.org/10.1051/parasite/2004112169>.
- Eiras JC, Takemoto RM, Pavanelli GC. *Henneguya caudicula* n. sp. (Myxozoa, Myxobolidae) a parasite of *Leporinus lacustris* (Osteichthyes, Anostomidae) from the high Paraná River, Brazil, with a revision of *Henneguya* spp. infecting South American fish. *Acta Protozool* 2008; 47(2): 149-154.
- Eszterbauer E. Genetic relationship among gill-infecting *Myxobolus* species (Myxosporea) of cyprinids: molecular evidence of importance of tissue-specificity. *Dis Aquat Organ* 2004; 58(1): 35-40. PMid:15038449. <http://dx.doi.org/10.3354/dao058035>.
- Eszterbauer E, Marton S, Rácz OZ, Letenyei M, Molnár K. Morphological and genetic differences among actinosporean stages of fish-parasitic myxosporeans (Myxozoa): difficulties of species identification. *Syst Parasitol* 2006; 65(2): 97-114. PMid:16676228. <http://dx.doi.org/10.1007/s11230-006-9041-y>.
- Fiala I. The phylogeny of Myxosporea (Myxozoa) based on small subunit ribosomal RNA gene analysis. *Int J Parasitol* 2006; 36(14): 1521-1534. PMid:16904677. <http://dx.doi.org/10.1016/j.ijpara.2006.06.016>.
- Froese R, Pauly D, editors. *FishBase* [online]. 2016. [cited 2016 Nov 20]. Available from: [www.fishbase.org](http://www.fishbase.org)
- Griffin MJ, Khoo LH, Torrants L, Bosworth BG, Quiniou SM, Gaunt PS, et al. New data on *Henneguya pellis* (Myxozoa: Myxobolidae), a parasite of blue catfish *Ictalurus furcatus*. *J Parasitol* 2009; 95(6): 1455-1467. PMid:19575542. <http://dx.doi.org/10.1645/GE-2106.1>.
- Griffin MJ, Pote LM, Wise DJ, Greenway TE, Mael MJ, Camus AC. A novel *Henneguya* species from channel catfish described by morphological, histological, and molecular characterization. *J Aquat Anim Health* 2008; 20(3): 127-135. PMid:18942589. <http://dx.doi.org/10.1577/H07-001.1>.
- Guindon S, Gascuel O. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst Biol* 2003; 52(5): 696-704. PMid:14530136. <http://dx.doi.org/10.1080/10635150390235520>.
- Hervio DML, Kent ML, Khattra J, Sakanari J, Yokoyama H, Devlim RH. Taxonomy of *Kudoa* species (Myxosporea), using a small-subunit ribosomal DNA sequence. *Can J Zool* 1997; 75(12): 2112-2119. <http://dx.doi.org/10.1139/z97-846>.
- Holzer ASC, Sommerville RW, Wootten R. Molecular relationships and phylogeny in a community of myxosporeans and actinosporeans based on their 18S rDNA sequences. *Int J Parasitol* 2004; 34(10): 1099-1111. PMid:15380681. <http://dx.doi.org/10.1016/j.ijpara.2004.06.002>.
- Holzer AS, Sommerville C, Wootten R. Molecular studies on the seasonal occurrence and development of five myxozoans in farmed *Salmo trutta* L. *Parasitology* 2006; 132(2): 193-205. PMid:16216135. <http://dx.doi.org/10.1017/S0031182005008917>.
- Huelsenbeck JP, Ronquist F. MrBayes: Bayesian inference of phylogenetic trees. *Bioinformatics* 2001; 17(8): 754-755. PMid:11524383. <http://dx.doi.org/10.1093/bioinformatics/17.8.754>.
- Iwanowicz LR, Iwanowicz DD, Pote LM, Blazer VS, Schill WB. Morphology and 18S rDNA of *Henneguya gurlei* (Myxosporea) from *Ameiurus nebulosus* (Siluriformes) in North Carolina. *J Parasitol* 2008; 94(1): 46-57. PMid:18372621. <http://dx.doi.org/10.1645/GE-1092.1>.
- Kageyama T, Yanagida T, Ohara K, Yokoyama H. *Henneguya pseudorhinogobii* n. sp. (Myxozoa: Myxosporea) parasitizing the gills of the freshwater goby *Rhinogobius* sp. or from the Nagara River and redescription of *Henneguya rhinogobii*. *Fish Sci* 2009; 75(3): 657-663. <http://dx.doi.org/10.1007/s12562-009-0096-y>.
- Kallert DM, Eszterbauer E, El-Matbouli M, Erséus C, Haas W. The life cycle of *Henneguya nuesslini* Schuberg & Schro 1905 (Myxozoa) involves a triactinomyxon-type actinospore. *J Fish Dis* 2005; 28(2): 71-79. PMid:15705152. <http://dx.doi.org/10.1111/j.1365-2761.2004.00599.x>.
- Kent ML, Andree KB, Bartholomew JL, El-Matbouli M, Desser SS, Devlin RH, et al. Recent advances in our knowledge of the Myxozoa. *J Eukaryot Microbiol* 2001; 48(4): 395-413. PMid:11456316. <http://dx.doi.org/10.1111/j.1550-7408.2001.tb00173.x>.
- Kent ML, Khattra J, Hedrick RP, Devlin RH. *Tetracapsula renicola* n. sp. (Myxozoa: Saccosporidae); the pkx myxozoan: the cause of proliferative kidney disease of salmonid fishes. *J Parasitol* 2000; 86(1): 103-111. [http://dx.doi.org/10.1645/0022-3395\(2000\)086\[0103.PMID:10701572](http://dx.doi.org/10.1645/0022-3395(2000)086[0103.PMID:10701572).
- Kudo R. Studies on Myxosporidia: a synopsis of genera and species of Myxosporidia. *Illinois Biol Monogr* 1919; 5(3): 1-265.
- Kudo R. A taxonomic consideration of Myxosporidia. *Trans Am Microsc Soc* 1933; 52(3): 195-216. <http://dx.doi.org/10.2307/3222254>.
- Li Y, Zhang Y, Siriguleng, Sato H, Siriguleng, Hiroshi S. *Henneguya doneci* (Myxozoa: Bivalvulida) in the gill filaments of Prussian carp *Carassius gibelio* (Bloch) from the upper Yellow River running through Inner Mongolia, China. *J Vet Med Sci* 2015; 77(8): 1001-1005. PMid:25843612. <http://dx.doi.org/10.1292/jvms.14-0666>.
- Lin D, Hanson LA, Pote LM. Small subunit ribosomal RNA sequence of *Henneguya exilis* (Class Myxosporea) identifies the actinosporean stage from an oligochaete host. *J Eukaryot Microbiol* 1999; 46(1): 66-68. PMid:10188262. <http://dx.doi.org/10.1111/j.1550-7408.1999.tb04585.x>.
- Lom J, Arthur JR. A guideline for the preparation of species descriptions in Myxosporea. *J Fish Dis* 1989; 12(2): 151-156. <http://dx.doi.org/10.1111/j.1365-2761.1989.tb00287.x>.
- Lom J, Dyková I. Myxozoan genera: definition and notes on taxonomy, life-cycle terminology and pathogenic species. *Folia Parasitol (Praha)* 2006; 53(1): 1-36. PMid:16696428. <http://dx.doi.org/10.14411/fp.2006.001>.
- Martins ML, Souza VN, Moraes JRE, Moraes FR. Gill infection of *Leporinus macrocephalus* Garavello & Britski, 1988 (Osteichthyes: Anostomidae) by *Henneguya leporinicola* n. sp. (Myxozoa: Myxobolidae). Description, histopathology and treatment. *Rev Bras Biol* 1999; 59(3): 527-534. PMid:10765464. <http://dx.doi.org/10.1590/S0034-71081999000300018>.
- Milanin T, Maia AAM, Silva MR, Carriero MM, Adriano EA. Molecular phylogeny and ultrastructure of *Myxobolus* cf. *cuneus*, a parasite of patinga hybrid and *Henneguya pseudoplatystoma*, a parasite of pintado hybrid. *Acta Parasitol* 2015; 60(3): 442-450. PMid:26204181. <http://dx.doi.org/10.1515/ap-2015-0061>.
- Molnár K. Comment on the host, organ and tissue specificity of fish myxosporeans and on the types of their intrapiscine development. *Parasitol Hung* 1994; 27: 5-20.
- Molnár K. Site preference of fish myxosporeans in the gill. *Dis Aquat Organ* 2002; 48(3): 197-207. PMid:12033706. <http://dx.doi.org/10.3354/dao048197>.

- Moreira GSA, Adriano EA, Silva MRM, Ceccarelli PS, Maia AAM. The morphological and molecular characterization of *Henneguya rotunda* n. sp., a parasite of the gill arch and fins of *Salminus brasiliensis* from the Mogi Guaçu River, Brazil. *Parasitol Res* 2014a; 113(5): 1703-1711. PMid:24535737. <http://dx.doi.org/10.1007/s00436-014-3815-y>.
- Moreira GSA, Adriano EA, Silva MRM, Ceccarelli OS, Maia AAM. Morphology and 18S rDNA sequencing identifies *Henneguya visibilis* n. sp., a parasite of *Leporinus obtusidens* from Mogi Guaçu River, Brazil. *Parasitol Res* 2014b; 113(1): 81-90. PMid:24100607. <http://dx.doi.org/10.1007/s00436-013-3629-3>.
- Müller MI, Adriano EA, Ceccarelli PS, Silva MRM, Maia AAM, Ueta MT. Prevalence, intensity, and phylogenetic analysis of *Henneguya piaractus* and *Myxobolus cf. colossomatis* from farmed *Piaractus mesopotamicus* in Brazil. *Dis Aquat Organ* 2013; 107(2): 129-139. PMid:24334355. <http://dx.doi.org/10.3354/dao02668>.
- Naldoni J, Arana S, Maia AAM, Ceccarelli PS, Tavares LER, Borges FA, et al. *Henneguya pseudoplatystoma* n. sp. causing reduction in epithelial area of gills in the farmed pintado, a South American catfish: Histopathology and ultrastructure. *Vet Parasitol* 2009; 166(1-2): 52-59. PMid:19695782. <http://dx.doi.org/10.1016/j.vetpar.2009.07.034>.
- Naldoni J, Maia AAM, Silva MRM, Adriano EA. *Henneguya cuniculator* sp. nov., a parasite of spotted sorubim *Pseudoplatystoma corruscans* in the São Francisco Basin, Brazil. *Dis Aquat Organ* 2014; 107(3): 211-221. PMid:24429472. <http://dx.doi.org/10.3354/dao02685>.
- Naldoni J, Zatti SA, Capodifoglio KRH, Milanin T, Maia AAM, Silva MRM, et al. Host-parasite and phylogenetic relationships of *Myxobolus filamentum* sp. n. (Myxozoa: Myxosporea), a parasite of *Brycon orthotaenia* (Characiformes: Bryconidae) in Brazil. *Folia Parasitol (Praha)* 2015; 62(1): 4-11. <http://dx.doi.org/10.14411/fp.2015.014>. PMid:25960558.
- Okamura B, Gruhl A, Bartholomew JL. An introduction to Myxozoan evolution, ecology and development. In: Okamura B, Gruhl A, Bartholomew JL. *Myxozoan evolution, ecology and development*. Heidelberg: Springer; 2015. p. 1-20. <http://doi.org/10.1007/978-3-319-14753-6>
- Pote LM, Hanson LA, Shivaji R. Small subunit ribosomal RNA sequences link the cause of proliferative gill disease in channel catfish to *Henneguya* n. sp. (Myxozoa: Myxosporea). *J Aquat Anim Health* 2000; 12(3): 26-34. [http://dx.doi.org/10.1577/1548-8667\(2000\)012<0230:SSRRSL>2.0.CO;2](http://dx.doi.org/10.1577/1548-8667(2000)012<0230:SSRRSL>2.0.CO;2)
- Rosser TG, Griffin MJ, Quiniou SMA, Khoo LH, Pote LM. 18S rRNA gene sequencing identifies a novel species of *Henneguya* parasitizing the gills of the channel catfish (Ictaluridae). *Parasitol Res* 2014; 113(12): 4651-4658. PMid:25270236. <http://dx.doi.org/10.1007/s00436-014-4156-6>.
- Rosser TG, Griffin MJ, Quiniou SMA, Khoo LH, Greenway TE, Wise DJ, et al. Small subunit ribosomal RNA sequence links the myxospore stage of *Henneguya mississippiensis* n. sp. from channel catfish *Ictalurus punctatus* to an actinospore released by the benthic oligochaete *Dero digitata*. *Parasitol Res* 2015; 114(4): 1595-1602. PMid:25716821. <http://dx.doi.org/10.1007/s00436-015-4345-y>.
- Siddall ME, Martin DS, Bridge D, Desser SS, Cone DK. The demise of a Phylum of Protists: Phylogeny of Myxozoa and other parasitic Cnidaria. *J Parasitol* 1995; 81(6): 961-967. PMid:8544072. <http://dx.doi.org/10.2307/3284049>.
- Smother JF, von Dohlen CD, Smith LH Jr, Spall RD. Molecular evidence that the myxozoan Protists are Metazoans. *Science* 1994; 265(5179): 1719-1721. PMid:8085160. <http://dx.doi.org/10.1126/science.8085160>.
- Whipps CM, Adlard RD, Bryant MS, Lester RGJ, Findlav V, Kent ML. First report of three *Kudoa* species from eastern Australia: *Kudoa thrysites* from mahi mahi (*Coryphaena hippurus*), *Kudoa amamiensis* and *Kudoa minithrysites* n. sp. from sweeper (*Pempheris ypsilichnus*). *J Eukaryot Microbiol* 2003; 50(3): 215-219. PMid:12836879. <http://dx.doi.org/10.1111/j.1550-7408.2003.tb00120.x>.
- Whipps CM, Murray KN, Kent ML. Occurrence of a Myxozoan parasite *Myxidium streisingeri* n. sp. in Laboratory Zebrafish *Danio rerio*. *J Parasitol* 2015; 101(1): 86-90. PMid:25277837. <http://dx.doi.org/10.1645/14-613.1>.
- Ye LT, Li WX, Wu SG, Wang GT. Supplementary studies on *Henneguya doneci* Schulman, 1962 (Myxozoa: Myxosporea) infecting the gill filaments of *Carassius auratus gibelio* (Bloch) in China: histologic, ultrastructural, and molecular data. *Parasitol Res* 2012; 110(4): 1509-1516. PMid:21989578. <http://dx.doi.org/10.1007/s00436-011-2655-2>.
- Yokoyama H, Urawa S, Grabner D, Shirakashi S. *Henneguya cartilaginis* n. sp. (Myxozoa: Myxosporea) in the head cartilage of masu salmon *Oncorhynchus masou masou*. *Parasitol Int* 2012; 61(4): 594-598. PMid:22664475. <http://dx.doi.org/10.1016/j.parint.2012.05.013>.
- Zhang J, Mo X, Li N, Chen W, Yang C. Supplementary description of *Henneguya zikawiensis* Sikama, 1938 and its molecular phylogeny. *J Neijiang Norm Univ* 2015; 35(4): 42-46. <http://dx.doi.org/10.13603/j.cnki.51-1621/z.2015.04.010>.