


## Revista da Sociedade Brasileira de Medicina Tropical

Journal of the Brazilian Society of Tropical Medicine








## **Images in Infectious Diseases**

## Giant lung and liver hydatid cyst in a 3-year-old child

Mesut Ozgokce<sup>[1]</sup>, Kemal Ayengin<sup>[2]</sup> and Yener Aydin<sup>[3]</sup>

- [1]. Yuzuncu Yil University, Medical Faculty, Department of Radiology, Van, Turkey.
- [2]. Yuzuncu Yil University, Medical Faculty, Department of Pediatric Surgery, Van, Turkey.
- [3]. Ataturk University, Medical Faculty, Department of Thoracic Surgery, Erzurum, Turkey.








FIGURE 1: Direct radiography shows a radiopaque appearance in the right lower zone of the lung in the paracardial region and the left lower zone of the lung. The homogeneous opacity increase (arrows) pushed on the intestinal loops (A). An axial thoracic computed tomography (CT) mediastinal section shows a cyst in the right middle lobe (arrow) compressing the right atrium and a giant hydatid cyst filling the lower lobe of the left lung (B). An abdominal CT section shows a giant hydatid cyst with similar characteristics in the liver (C).

A 3-year-old girl presented with growth retardation. Radiologically, a radiopacity consistent with a hydatid cyst was noted in both the lungs and the liver (Figure 1). The patient underwent surgical treatment to remove the cyst from the lungs and liver.

Hydatid disease is a parasitic disease transmitted via the fecal-oral route1. In approximately 8% of cases, the hydatid cyst is located in both the liver and lungs<sup>1</sup>. When the hydatid cyst is intact, it appears on direct radiographs as a round radiopacity with smooth margins. In endemic areas, the diagnosis is made by clinical and radiologic findings. Growth retardation rarely occurs in hydatid cysts<sup>1</sup>. Serology may help diagnose, especially if the cyst ruptures and the case becomes complicated<sup>2</sup>. Although hydatid cysts can be large, the coexistence of giant cysts in both the liver and lungs is rare in 3-year-old children<sup>3</sup>. In these cases, the hydatid cyst should be treated immediately.

## **REFERENCES**

- Aydin Y, Ulas AB, Ince I, Korkut E, Ogul H, Eren S, et al. Large Case Series Analysis of Cystic Echinococcosis. Indian J Surg. 2021;83(Suppl 4):S897-S906. Available from: https://doi.org/10.1007/s12262-021-03061-0
- Agudelo Higuita NI, Brunetti E, McCloskey C. Cystic Echinococcosis. J Clin Microbiol. 2016;54(3):518-23.
- Tekin R, Tekin RC, Avcı A. Giant Hydatid Cysts of the Lung and Liver. Rev Soc Bras Med Trop. 2020;53:e20190492.

Corresponding author: Dr. Yener Aydin. e-mail: dryeneraydin@hotmail.com

Authors' contribution: MO: Conception and design of the study, Acquisition of data, Drafting the article, Final approval of the version to be submitted; KA: Conception and design of the study, Analysis and interpretation of data; YA: Conception and design of the study, Analysis and interpretation of data.

Conflict of Interest: The authors declared no conflicts of interest with respect to the authorship and/or publication of this article.

Financial Support: The authors declare that the article content was composed in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

www.scielo.br/rsbmt I www.rsbmt.org.br

Received 6 April 2022 | Accepted 1 July 2022









