Revista da Sociedade Brasileira de Medicina Tropical Journal of the Brazilian Society of Tropical Medicine

Vol.:54:(e0641-2020): 2021



https://doi.org/10.1590/0037-8682-0641-2020

Images in Infectious Diseases

Acute necrotizing encephalopathy with H1N1 virus infection in a child

Berhan Pirimoglu^[1], Halil Keskin^[2] and Huseyin Tan^[3]

[1]. Ataturk University, Medical Faculty, Department of Radiology, Erzurum, Turkey.
[2]. Ataturk University, Medical Faculty, Department of Pediatric Intensive Care Unit, Erzurum, Turkey.
[3]. Ataturk University, Medical Faculty, Department of Pediatric Neurology, Erzurum, Turkey.

A 7-year-old man presented to our radiology unit with high fever, generalized tonic-clonic type seizures, and neurological deterioration requiring intensive care. Cerebrospinal fluid analysis showed increased protein (94 mg/dL) and normal glucose (41 mg/ dL) levels. An initial brain magnetic resonance imaging (MRI) revealed symmetrical and expansile hyperintense lesions in the bilateral thalami and external capsules (Figure 1A and 1B). Restricted diffusion patterns of the involved brain regions were detected (Figure 1C and 1D). These radiological imaging findings suggested acute viral encephalitis, compatible with acute necrotizing encephalopathy (ANEC). Real-time polymerase chain reaction test performed with samples from nasopharyngeal swabs was positive for influenza A virus (H1N1). Therefore, ANEC-associated H1N1 virus infection was diagnosed. The patient underwent antiviral, anticonvulsant, and anti-inflammatory therapy. Fifteen days after therapy, a follow-up brain MRI revealed a decrease in signal intensity in the extent of lesions involving the bilateral thalami and external capsules with decreases in diffusion restriction patterns (Figure 2).

ANEC is an unusual type of encephalopathy characterized by multiple bilateral brain lesions, involving the thalami, putamina, internal and external capsules, cerebellar white matter, and brainstem. It usually develops secondary to viral infections, including influenza virus A and B, parainfluenza, varicella, and




FIGURE 1: (A) Axial T2-weighted image and (B) axial FLAIR image show the symmetrical hyperintense expansile lesions in the bilateral thalami and external capsules (black asterisks). (C) Axial b-1000 image (black asterisk) and (D) axial ADC diffusion image (white asterisk) show the symmetrical diffusion restriction in the bilateral thalami and external capsules.

Corresponding author: Dr. Berhan Pirimoglu. **e-mail**: berhan.dr@gmail.com

https://orcid.org/0000-0002-1608-1189

Received 19 September 2020 **Accepted** 19 October 2020




FIGURE 2: (A) Axial FLAIR image and **(B)** coronal T2-weighted image show the decreased symmetrical hyperintense lesions in the bilateral thalami and external capsules (black asterisks). **(C)** Axial b-1000 image (black asterisk) and **(D)**, axial ADC diffusion image (white asterisk) show the regression of symmetrical diffusion restriction pattern in the bilateral thalami

severe acute respiratory syndrome coronavirus infections. Abnormal signals on MRI consisted of hypointense and hyperintense signals on T1-weighted and T2-weighted images, respectively. Restricted diffusion patterns of the involved regions could be detected on diffusion-weighted images¹⁻³.

AUTHORS' CONTRIBUTION

BP: conceptualization, data curation, resources, software, writing-review, and editing; HK: supervision, validation, writing-original draft, conceptualization, visualization; HT: validation, and writing review and editing.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

REFERENCES

- Skelton BW, Hollingshead MC, Sledd AT, Phillips CD, Castillo M. Acute necrotizing encephalopathy of childhood: typical findings in an atypical disease. Pediatr Radiol. 2008;38: 810-3.
- Poyiadji N, Shahin G, Noujaim D, Stone M, Patel S, Griffith B. COVID-19-associated Acute Hemorrhagic Necrotizing Encephalopathy: Imaging Features. Radiology. 2020;296:E119-20.
- Albayram S, Bilgi Z, Selcuk H, Selcuk D, Cam H, Kocer N, et al. Diffusion-weighted MR imaging findings of acute necrotizing encephalopathy. AJNR Am J Neuroradiol. 2004;25:792–97.