

Synthesis and Expansion of Bicyclic Enol Ether: A Probable Precursor for the Synthesis of Macrolide (\pm)-Pyrenophorin

Maísa B. Costa,^{*,a,#} *Marcos P. Martins*,^a *Hugo C. de Araújo*^b and *Inês S. Resck*^b

^aLaboratório de Síntese, Isolamento e Modificações de Compostos Orgânicos, Universidade Estadual de Goiás, Campus Henrique Santillo, BR 153, No. 3105, Fazenda Barreiro do Meio, Setor Arco Verde, CP 459, 75132-400 Anápolis-GO, Brazil

^bInstituto de Química, Universidade de Brasília, Campus Darcy Ribeiro, CP 4478, 70910-970 Brasília-DF, Brazil

A convenient procedure for the synthesis and expansion of bicyclic rings has been developed for the production of probable precursors of non-racemic pyrenophorin, an antibiotic dilactone. The major highlight for this new synthetic methodology came from the use of a readily available reagent of easy manipulation, 9-oxabicyclo[3.3.1]nonane-2,6-diol, for the preparation of the bicyclic intermediate, which sequentially was subjected to oxidative cleavage with butyl nitrite resulted in an isomeric mixture, a dioximedilactone and diisoxazoledilactone.

Keywords: enol ether, ring expansion, macrolides, pyrenophorin

Introduction

The structural diversity of the macrolides represents an interesting synthetic challenge for the obtainment of molecules with medium or macrocyclic sizes. Following the research conducted in our laboratory to obtain macrocyclic lactones, it was developed the synthesis of a precursor that will be later applied to the racemic synthesis of the natural 16-membered macrolide pyrenophorin¹ (**1**), a macrolactone dimer with high biological potential (Figure 1).




Figure 1. Natural macrolide **1**.

The bioactive metabolite **1**, pyrenophorin, was isolated from the pathogenic fungi *Pyrenophora avenae* and *Stemphylium radicinum*,¹ and the cultures of *Drechslera avenae* acting as a phytotoxin on leaves segments of *Avena sterilis*.²

*e-mail: maisabc@gmail.com, maisa.costa@ueg.br

#Synthetic part of this research was initially developed at the Instituto de Química, Universidade de Brasília, and is related to the doctoral thesis.

In as much of the wide spectrum of biological activities, mainly as antibiotics, several synthetic methodologies for the synthesis of this 16-member macrolide, racemic or chiral, have been reported in the literature. Many are based on the lactonization and the dimeric cyclization of seco- ω -hydroxy-acid derivatives and γ -oxo- α , β -unsaturated derivatives, as well as the formation of intramolecular C–C bonds, in addition to cyclodimerization under Mitsunobu conditions.³⁻⁷

Therefore, in this article, a new procedure is demonstrated to obtain potential synthetic precursors that may lead to the synthesis of (\pm)-pyrenophorin (**1**) through the use of oxidative ring expansion methodology.⁸

Experimental


General methods

All reagents used were of analytical grade. The products of all reactions were monitored by capillary gas chromatography on VARIAN STAR 3400 CX equipment using DB-5 and DB-1 capillary columns of length equal to 30 m \times 25 mm, with a temperature range of 120-190 °C, injector temperature of 250 °C and the detector of 300 °C. The DEX-CB capillary chiral column with a length equal to 25 m \times 25 mm, with a temperature range of 160-200 °C, injector temperature of 250 °C and detector of 300 °C, was

used in specific cases. The products were purified on flash chromatographic column using silica gel (230-400 mesh). The melting points were determined in the K ofler block and were recorded without correction. The ^1H and ^{13}C (single and two-dimensional) nuclear magnetic resonance (NMR) spectra were obtained on the spectrometer. The probes used were ATB and SW, with 5 mm internal diameter, at room temperature with 45° pulse for hydrogen and carbon. Chemical displacements (δ) in ^1H NMR, with deuterated solvents CDCl_3 and $\text{DMSO}-d_6$, were referenced with TMS and the DMSO residue (δ 2.49), respectively. In ^{13}C NMR, they were referenced with δ 77.0 (CDCl_3) and δ 39.7 ($\text{DMSO}-d_6$). The multiplicities were defined in the usual way: s (singlet), d (doublet), dd (double doublet), t (triplet), q (quartet), m (multiplet). Low resolution mass spectra were obtained from the Shimadzu GC-17A (Civil Police, Distrito Federal) equipment. High resolution mass spectra were obtained from VG-auto SPEC (IQ, Unicamp) equipment. Elemental analyzes were obtained from the Analytische Laboratorien Prof. Dr. H. Malissa und G. Reuter GmbH.

Synthesis of 9-oxabicyclo[3.3.1]nonane-2,6-diol (**3**)

The isomeric mixture of 9-oxabicyclo[3.3.1]nonane-2,6-diol (**3**) and 9-oxabicyclo[4.2.1]nonane-2,5-diol (**3a**) was prepared by the oxidation of cycloocta-5-diene (**2**) (1.18 mol) under treatment with formic acid⁹⁻¹¹ (1.06 mol) and under vigorous magnetic stirring and, in an ice bath, 30% H_2O_2 (300 mL, 9.20 mol) was added dropwise. After the addition, the formation of a biphasic mixture was observed, which was maintained at 50°C (internal temperature). The mixture became homogeneous after two days under magnetic stirring at 50°C . Subsequently, 10% Pd/C (0.10 g) was added to decompose the excess of H_2O_2 . This step was monitored with KI/starch indicator paper. The solution was filtered under celite and subjected to liquid-liquid extraction with ethyl acetate for 72 h. The organic phase was concentrated on a rotary evaporator in which provided a yellow solid of melting point equal to $55\text{--}58^\circ\text{C}$ in 85% yield (1.50 mol). The diol **3** was isolated from the isomeric mixture, **3** and **3a**, by successive recrystallizations from ethyl acetate. Chromatographic analysis and NMR confirmed the structure and purity of compound **3** (Scheme 1).




Scheme 1. Reaction conditions for synthesis of diol **3**.

9-Oxabicyclo[3.3.1]nonane-2,6-diol (**3**)

White solid, m.p. $124\text{--}125^\circ\text{C}$; IR (KBr) ν / cm^{-1} 3422, 3282, 2949, 2909, 1483, 1079, 1062, 974; ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 4.78 (d, J 6 Hz, OH), 3.68 (2H), 3.51 (dd, J 5.0, 5.0 Hz, 2H), 2.01-1.88 (m, 2H), 1.74-1.53 (m, 6H); ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) δ 69.4, 67.5, 28.8, 22.0; anal. calcd. for $\text{C}_8\text{H}_{14}\text{O}_3$ (158.0946 g mol⁻¹): C, 54.53; H, 9.15%. Found: C, 54.52; H, 9.11%.

Synthesis of 2,6-dioxo-9-oxabicyclo[3.3.1]nonane (**4**)

The 2-iodoxybenzoic acid¹² (8.40 g; 30.0 mmol) was added to the solution of diol **3** (1.58 g, 10.0 mmol) in ethyl acetate (60 mL) and stirred magnetically at 80°C . After 6 h of reaction, the suspension was filtered through a column of florisil (60-100 mesh) washed with ethyl acetate (2×30 mL) and evaporated. The product obtained, recrystallized from isopropanol, with degree of high purity demonstrated by gas chromatographic was characterized as a white solid (9.80 mmol; 99% m.p. $49\text{--}51^\circ\text{C}$) (Scheme 2).



Scheme 2. Reaction conditions for synthesis of diketone **4**.

2,6-Dioxo-9-oxabicyclo[3.3.1]nonane (**4**)

White solid, m.p. $49\text{--}51^\circ\text{C}$; IR (KBr) ν / cm^{-1} 3409, 2950, 2927, 1715, 1455, 1299, 1242, 1189, 1127; ^1H NMR (300 MHz, CDCl_3) δ 4.38 (m, 2H), 2.80-2.69 (m, 2H), 2.57-2.38 (m, 4H), 2.17-2.04 (m, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ 209.6, 75.7, 33.5, 25.0; anal. calcd. for $\text{C}_8\text{H}_{10}\text{O}_3$ (154.0629 g mol⁻¹): C, 62.33; H, 6.54%. Found: C, 62.20; H, 6.51%.

Synthesis of 2,8-dimethyl-3,4,6,8,9,10,11,12-octahydro-2H,5H-pyrano[2',3',5,6]cycloocta[1,2-b]pyran-6,12-diol (**7**)

Diketone **4** (1.54 g, 10.0 mmol), anhydrous cyclohexane (50 mL), morpholine (4.40 mL, 30.0 mmol) and nafion 417[®] (2.00 mm) were refluxed under stirring on a Dean-Stark for 48 h to convert **4** into enamine **5**. The solvent (cyclohexane) and the residual reactant (morpholine) were removed under reduced pressure (1.00 torr) resulting in a yellowish solid (2.67 g; 9.10 mmol; 90%) which was used in the next step without prior purification.

The solid crude enamine **5** (9.10 mmol) was subjected to alkylation (Michael addition) in anhydrous dioxane (50 mL) and methyl vinyl ketone (2.50 mL, 30.0 mmol) was dripped, under controlled conditions (N_2 atmosphere

and CaCl₂ tube). After 24 h reflux, the residual solvents and reagents were removed under reduced pressure to provide a viscous yellowish liquid. NaBH₄ (0.76 g, 20.0 mmol) was added to the solution of the alkylated enamine **6** in ethanol (25 mL), cooled in an ice bath. The reagent suspension was stirred for 0.5 h at that temperature, then for 2 h at room temperature.

After this, the residual NaBH₄ was decomposed with 20% aqueous HCl solution (ca. 15 mL). Then a further 30 mL of 20% aqueous HCl solution and fourteen hour reflux was added to occur the hydrolysis and cyclization of the reduced product **7** (Scheme 3). The reaction was cooled and then the aqueous phase was extracted with chloroform (3 × 30 mL), and the combined organic phases were washed with saturated NaHCO₃ solution (3 × 30 mL) and brine (3 × 20 mL). After manipulation, a yellowish viscous liquid (1.72 g, 6.10 mmol, 81%) was obtained. Purification by dry flash chromatography column (10% hexane/ethyl acetate) yielded yellowish crystals (1.42 g, 5.00 mmol, 50%) which recrystallized from ethyl acetate or dichloromethane provided white crystals (m.p. 102-103 °C; 41%) with R_t (retention time) 14.18 min (50.95%) and 14.38 min (49.04%).

2,8-Dimethyl-3,4,6,8,9,10,11,12-octahydro-2*H*,5*H*-pyrano[2',3',5,6]cycloocta[1,2-*b*]pyran-6,12-diol (**7**)

White solid, m.p. 102-103 °C; IR (KBr) ν / cm⁻¹ 3393, 2973, 2928, 2875, 1705, 1449, 1365, 1322, 1219, 1186, 1089, 890, 801, 765; ¹H NMR (300 MHz, CDCl₃) δ 4.15-4.05 (m, 3H), 3.97-3.89 (m, 3H), 3.82-3.75 (m, 1H), 2.42-1.54 (m, 22H), 1.27 (d, *J* 6.2 Hz, 3H), 1.22 (d, *J* 6.3 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 145.5, 102.5, 102.0, 72.0, 71.2, 70.8, 68.9, 67.0, 66.8, 30.0, 28.9, 27.9, 25.5, 22.2, 21.2, 19.7; anal. calcd. for C₁₆H₂₄O₄ (281 g mol⁻¹): C, 68.54; H, 8.63%. Found: C, 68.42; H, 8.88%.


Synthesis for the expansion of 2,8-dimethyl-3,4,6,8,9,10,11,12-octahydro-2*H*,5*H*-pyrano[2',3',5,6]cycloocta[1,2-*b*]pyran-6,12-diol (**7**)

Butyl nitrite (0.60 mL, 5.00 mmol) was added to a solution of enol ether **7** (0.28 g, 1.00 mmol) solubilized in ethanol, kept under ice-bath. The reaction flask was carefully sealed with a rubber septum to prevent the loss of nitrous acid formed with the addition of 10% HCl (5 mL). After 1 h of stirring at 0 °C, the reaction was allowed to stand at -18 °C for 24 hours. The aqueous phase was extracted with dichloromethane (5 × 20 mL), and the organic extract was washed with brine (1 × 15 mL) and stirred with solid NaHCO₃. After the usual procedure, a yellowish semi-solid was obtained (0.26 g, 0.72 mmol, 70%) and purified by flash column chromatography (hexane/ethyl acetate 90:10 and 50:50). Column fractions were characterized by spectroscopic analyzes as oxime **8** (R_t 22.8 min), and isoxazol **9** (R_t 13.4 min).


Results and Discussion

The reaction sequence for the preparation of precursors **7**, **8** and **9** was initiated with the classical oxidation of cycloocta-1,5-diene (**2**),⁹⁻¹¹ to obtain diol **3**, which was obtained with its **3a** isomer in 85% yield. The separation of the diol **3** from the isomeric mixture **3** and **3a** was done by successive fractional recrystallizations of ethyl acetate.

High resolution mass spectrometry (MS) confirmed the exact molar mass (158.0946 g mol⁻¹) and provided the characteristic fragments of the molecule. The elemental analysis data (9.11% H, 54.52% C) presented a variation when compared with the theoretical data (8.92% H, 60.74% C).⁹ This difference is related to the presence of water of crystallization in the structure of diol **3**, which



Scheme 3. Reaction conditions for synthesis of enol ether **7**.



Scheme 4. Reaction conditions for expansion of enol ether **7**.

assertion is confirmed when the data obtained are compared with the theoretical data containing water of crystallization in the calculations (9.15% H, 54.53% C).⁹

The conversion of 9-oxabicyclo[3.3.1]nonane-2,6-diol (**3**) to 2,6-dioxo-9-oxabicyclo[3.3.1]nonane (**4**) was successfully achieved in 99% yield from the use of the oxidant 2-iodoxybenzoic acid oxidant.¹² High resolution mass spectrometry (MS) confirmed the exact molar mass (m/z 154.0629) and provided the characteristic fragments of the molecule. Data from the elemental analysis (6.51% H, 62.20% C) were consistent with the theoretical data (6.54% H, 62.33% C).⁹

Gross enamine **5**, obtained from diketone **4**, was alkylated, reduced and hydrolyzed in a same pot and provided precursor **7**, which had its structure elucidated from spectrometric techniques including 1D and 2D NMR.

The infrared spectrum analysis of enol ether **7** showed the OH bands at 3393 cm^{-1} , C=C at 1704 cm^{-1} , and C–O at the region of 1015 to 1200 cm^{-1} , present in the cyclized product **7**. The hydroxyls present proved that the C–O related bonds of the bridging heads of the bicyclic system **4** were broken. The absorption at 1704 cm^{-1} relative to the C=C bond, although relatively high for the tetrasubstituted unsaturated groups, has been reported in the literature for enol ethers.¹³

To verify the chemical purity of product **7**, a low resolution mass spectrometry (MS) analysis was performed, which showed molar mass 281 g mol^{-1} and the characteristic fragmentations. The elemental analysis data (68.42% C, 8.88% H) are consistent with the theoretical data (68.54% C, 8.63% H).

The ¹H NMR spectrum (300 MHz, CDCl₃) of product **7** showed two doublets at δ 1.22 and 1.27 for the methyls of probable isomers. The absorptions in δ 1.54 to 2.42 relative to the CH₂, as well as the **7** hydrogens relative to the CH recorded in the region of δ 3.75 to 4.15, were related to the probable stereoisomers of the product **7**.

The ¹³C NMR spectrum (75.46 MHz, CDCl₃) showed the quaternary carbons of the ether enol **7** at δ 101.5, 102.3, 143.8 and 145.3, absent in the DEPT spectrum.

The absorptions of CH₂ (δ 21.0 to 35.0) and of **7** CH₃ (δ 66.0 to 72.0) confirmed the stereoisomers of the enol ether **7**.

By the 2D NMR techniques, some correlations of ¹H,¹H and ¹H,¹³C were observed, which aided in the structural elucidation of the isomeric mixture of **7**. The gCOSY (correlation spectroscopy) spectrum showed the methyl groups (δ 1.22 and 1.27) coupled with CH (δ 4.05–4.15 and δ 3.75–3.82), respectively.

The gHMBC (heteronuclear multiple bond correlation) experiment confirmed the sp² carbons (δ 144.3, 145.5 and

δ 102.0, 102.5) tetrasubstituted not shown in the gHMBC (heteronuclear multiple quantum correlation) spectrum.

Posteriorly, it was observed in CGC (capillary column gas chromatography) (DB-1) that compound **7**, submitted to successive recrystallizations, was enriched by one of the isomers. The analysis of ¹H NMR (300 MHz, CDCl₃) found different intensities of the two methyl groups.

The expansion of bicyclic ethers can be promoted by oxidative cleavage methods, such as ozonolysis,^{14,15} oxidation with *m*-chloroperbenzoic acid¹⁴ and hydrolytic nitrosation.^{16–20} In this work, to expand ring **7**, ozonolysis reactions and oximation/deoximation reactions were used.

The ozonolysis reaction was studied with substrate **7** under reaction conditions described in the literature.^{14,15} However, the products obtained with this reaction provided mixtures of compounds that could not be elucidated by means of spectroscopic analyzes.

The expansion of bicyclic ether enol by hydrolytic nitrosation with butyl nitrite was developed in our laboratory for the production of macrolide pyrenophorin (**1**),⁸ which resulted in an isomeric mixture (70%) formed mainly by dioximedilactone **8** and diisoxazoledilactone **9** (Scheme 4), in which traces of diketodilactone **9a** could be observed in the NMR and IR spectra.

The CGC analysis (DB-1) showed a mixture of compounds **8** and **9** which were obtained (60–70%) with the butyl nitrite oxidant.

The IR spectra presented the vibrational frequencies of lactone (1726 cm^{-1}) and oxime (1604 cm^{-1}). ¹H NMR (300 MHz, DMSO-*d*₆) and ¹³C NMR (75.46 MHz, DMSO-*d*₆) spectra showed characteristic absorptions of the oxime **8** and ketolactone **9a** (see Supplementary Information).

Subsequently, the product mixture was purified by flash chromatographic column (hexane/ethyl acetate 90:10 and 50:50) to provide enriched fractions of R_t 15.9; 13.4 and 22.8 min, previously observed by CGC.

The solid fractions of R_t 13.4 and 22.8 min observed in the CGC (DB-1) and ¹H NMR (300 MHz, CDCl₃), remained as main components even undergoing successive recrystallizations.

Afterwards this fraction was maintained in CDCl₃ solution and cooling for 2 days, there was enrichment of the R_t compound of 13.4 min (DB-1). ¹H NMR spectroscopic analyzes (300 MHz, CDCl₃) and ¹³C NMR (75.49 MHz, CDCl₃) showed a structure of the compound of R_t 13.4 min, being probably the isoxazole **9**.

In the isoxazole compound **9**, the prominent singlet at δ 5.80, related to CH sp², was characterized by the DEPT, gHMBC, gHMBC techniques.

The chemical displacement of the oxime **8** in δ 10.15

was not observed maybe because of the low concentration of this isomer in the compound mixture. In the ^1H NMR spectra (300 MHz, $\text{DMSO-}d_6$) doublets, with chemical shift δ 5.09, 5.11, 5.84 and 5.89 ppm, were obtained resulting from labile hydrogen interactions of the compounds of the mixture **9** and **9a** with this deuterated solvent.

Conclusions

The key intermediate **7** was prepared from the cheap starting material **3** in three steps and transformed into **8** and **9** in excellent yields. The conversion of the **8** and **9** compounds into dilactone **1** is under investigation and will be reported in due course.

Supplementary Information

Supplementary information associated with this work (NMR spectra (^1H and ^{13}C), infrared and elementary analysis) is available free of charge at <http://jbcs.s bq.org.br> as PDF file.

Acknowledgments

The authors are grateful to the Brazilian agencies CNPq and FINEP (grant CT-INFRA 970/01) for financial support, FAPEG/CAPES and PROBIP-2016/UEG.

References

1. Nozoe, S.; Hirai, K.; Tsuda, K.; Ishibashi, K.; Shirasaka, W.; Grove, J. F.; *Tetrahedron Lett.* **1965**, *51*, 4675.
2. Aliferis, K. A.; Chrysai-Tokousbalides, M.; Fasseas, C.; *Plant Physiol. Biochem.* **2006**, *44*, 851.
3. Kalita, D.; Klan, A. T.; Saikia, A. K.; Bez, G.; Barua, N. C.; *Synthesis* **1998**, *7*, 975.
4. Fürstner, A.; Thiel, O. R.; Ackermann, L.; *Org. Lett.* **2001**, *3*, 449.
5. Lee, C. W.; Grubbs, R. H.; *J. Org. Chem.* **2001**, *66*, 7155.
6. Rao, K. S.; Reddy, D. S.; Mukkanti, K.; Pal, M.; Iqbal, J.; *Tetrahedron Lett.* **2006**, *47*, 6623.
7. Ramakrishna, K.; Sreenivasulu, R.; Vidavalur, S.; Reddy, B. J. M.; *Lett. Org. Chem.* **2016**, *13*, 693.
8. Mahajan, J. R.; Araújo, H. C.; *Synthesis* **1981**, 49.
9. Duthaler, R. O.; Wicker, K.; Ackermann, P.; Ganter, C.; *Helv. Chim. Acta* **1972**, *55*, 1809.
10. Eaton, P. E.; Millikan, R.; *Synthesis* **1990**, 483.
11. Hegemann, K.; Fröhlich, R.; Haufe, G.; *Eur. J. Org. Chem.* **2004**, 2181.
12. More, J. D.; Finney, N. S.; *Org. Lett.* **2002**, *4*, 3001.
13. Mahajan, J. R.; Araújo, H. C.; *Synthesis* **1976**, 111.
14. Mahajan, J. R.; Ferreira, G. A. L.; Araújo, H. C.; Nunes, B. J.; *Synthesis* **1976**, 112.
15. Mahajan, J. R.; Araújo, H. C.; *Synthesis* **1975**, 53.
16. Mahajan, J. R.; Ferreira, G. A. L.; Araújo, H. C.; *J. Chem. Soc., Chem. Commun.* **1972**, 1078.
17. Mahajan, J. R.; Araújo, H. C.; *Synthesis* **1980**, 64.
18. Mahajan, J. R.; Resck, I. S.; *Synthesis* **1980**, 998.
19. Mahajan, J. R.; Monteiro, M. B.; *J. Chem. Res.* **1980**, 264.
20. Mahajan, J. R.; Araújo, H. C.; *Synthesis* **1981**, 49.

Submitted: April 16, 2017

Published online: June 23, 2017

